

Bayesian model averaging is suboptimal for generalization under model misspecification

Andrés Masegosa

*Department of Mathematics
University of Almería
Spain*

The learning problem

- **Notation:**

- $\nu(\mathbf{x})$ is the data generating distribution (**unknown**).
- $p(\mathbf{x}|\theta)$ is a probabilistic model parametrized by θ .
- $\forall \theta \nu(\mathbf{x}) \neq p(\mathbf{x}|\theta)$.

- **Notation:**

- $\nu(\mathbf{x})$ is the data generating distribution (**unknown**).
- $p(\mathbf{x}|\boldsymbol{\theta})$ is a probabilistic model parametrized by $\boldsymbol{\theta}$.
- $\forall \boldsymbol{\theta} \nu(\mathbf{x}) \neq p(\mathbf{x}|\boldsymbol{\theta})$.

- The **predictive posterior distribution** for a given $\rho(\boldsymbol{\theta})$,

$$p(\mathbf{x}) = \int p(\mathbf{x}|\boldsymbol{\theta})\rho(\boldsymbol{\theta})d\boldsymbol{\theta} = \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]$$

- **Notation:**

- $\nu(\mathbf{x})$ is the data generating distribution (**unknown**).
- $p(\mathbf{x}|\boldsymbol{\theta})$ is a probabilistic model parametrized by $\boldsymbol{\theta}$.
- $\forall \boldsymbol{\theta} \nu(\mathbf{x}) \neq p(\mathbf{x}|\boldsymbol{\theta})$.

- The **predictive posterior distribution** for a given $\rho(\boldsymbol{\theta})$,

$$p(\mathbf{x}) = \int p(\mathbf{x}|\boldsymbol{\theta})\rho(\boldsymbol{\theta})d\boldsymbol{\theta} = \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]$$

- The **learning problem** is defined as,

$$\rho^* = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})])$$

- **Notation:**

- $\nu(\mathbf{x})$ is the data generating distribution (**unknown**).
- $p(\mathbf{x}|\boldsymbol{\theta})$ is a probabilistic model parametrized by $\boldsymbol{\theta}$.
- $\forall \boldsymbol{\theta} \nu(\mathbf{x}) \neq p(\mathbf{x}|\boldsymbol{\theta})$.

- The **predictive posterior distribution** for a given $\rho(\boldsymbol{\theta})$,

$$p(\mathbf{x}) = \int p(\mathbf{x}|\boldsymbol{\theta})\rho(\boldsymbol{\theta})d\boldsymbol{\theta} = \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]$$

- The **learning problem** is defined as,

$$\rho^* = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]) = \arg \min_{\rho} \underbrace{\mathbb{E}_{\nu(\mathbf{x})} \left[\ln \frac{1}{\mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]} \right]}_{CE(\rho)}$$

- **Notation:**

- $\nu(\mathbf{x})$ is the data generating distribution (**unknown**).
- $p(\mathbf{x}|\boldsymbol{\theta})$ is a probabilistic model parametrized by $\boldsymbol{\theta}$.
- $\forall \boldsymbol{\theta} \nu(\mathbf{x}) \neq p(\mathbf{x}|\boldsymbol{\theta})$.

- The **predictive posterior distribution** for a given $\rho(\boldsymbol{\theta})$,

$$p(\mathbf{x}) = \int p(\mathbf{x}|\boldsymbol{\theta})\rho(\boldsymbol{\theta})d\boldsymbol{\theta} = \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]$$

- The **learning problem** is defined as,

$$\rho^* = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]) = \arg \min_{\rho} \underbrace{\mathbb{E}_{\nu(\mathbf{x})} \left[\ln \frac{1}{\mathbb{E}_{\rho(\boldsymbol{\theta})}[p(\mathbf{x}|\boldsymbol{\theta})]} \right]}_{CE(\rho)}$$

Learning from a finite dataset

- We do not have access to $\nu(\mathbf{x})$, only to a i.i.d. sample $D = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$.

First-order PAC-Bayes bounds

Remind!

$$\rho^* = \arg \min_{\rho} CE(\rho) = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\theta)}[p(\mathbf{x}|\theta)])$$

Remind!

$$\rho^{\star} = \arg \min_{\rho} CE(\rho) = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\theta)}[p(\mathbf{x}|\theta)])$$

$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}}$$

Remind!

$$\rho^* = \arg \min_{\rho} CE(\rho) = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\theta)}[p(\mathbf{x}|\theta)])$$

$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}} \stackrel{\text{PAC-Bayes}}{\lesssim} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta, D)] + \frac{KL(\rho, \pi) + \ln \frac{1}{\xi} + \psi_{\pi, \nu}(c, n)}{cn}}_{\text{PAC-Bayes bound}}$$

Remind!

$$\rho^* = \arg \min_{\rho} CE(\rho) = \arg \min_{\rho} KL(\nu(\mathbf{x}), \mathbb{E}_{\rho(\theta)}[p(\mathbf{x}|\theta)])$$

$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}} \stackrel{\text{PAC-Bayes}}{\lesssim} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta, D)] + \frac{KL(\rho, \pi) + \ln \frac{1}{\xi} + \psi_{\pi, \nu}(c, n)}{cn}}_{\text{PAC-Bayes bound}}$$

The learning strategy is to **minimize the PAC-Bayes bound**

- ρ^* is the **Bayesian posterior** for $c = 1$ (Germain et al. 2016),

$$\rho^* = p(\theta|D) = \frac{p(D|\theta)\pi(\theta)}{\int p(D|\theta)\pi(\theta)d\theta}$$

- The **Bayesian** learning strategy,

$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}} \stackrel{\text{PAC-Bayes}}{\lesssim} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta, D)] + \frac{KL(\rho, \pi) + \ln \frac{1}{\xi} + \psi_{\pi, \nu}(c, n)}{cn}}_{\text{PAC-Bayes bound}}$$


- The **Bayesian learning strategy**,

$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}} \stackrel{\text{PAC-Bayes}}{\lesssim} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta, D)] + \frac{KL(\rho, \pi) + \ln \frac{1}{\xi} + \psi_{\pi, \nu}(c, n)}{cn}}_{\text{PAC-Bayes bound}}$$

- The **minimum of the Jensen bound is a Dirac-delta distribution** centered around,


$$\theta_{ML}^* = \arg \min_{\theta} KL(\nu(\mathbf{x}), p(\mathbf{x}|\theta))$$

Is the Bayesian approach an optimal learning strategy?




$$CE(\rho)$$

Is the Bayesian approach an optimal learning strategy?




$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}}$$

Is the Bayesian approach an optimal learning strategy?




$$CE(\rho)$$

Is the Bayesian approach an optimal learning strategy?




$$CE(\rho) \stackrel{\text{Jensen}}{\leq} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)]}_{\text{Jensen bound}}$$

Is the Bayesian approach an optimal learning strategy?



$$CE(\rho) \stackrel{(Liao \& Berg, 2019)}{\leq} \underbrace{E_{\rho(\theta)}[L(\theta)] - V(\rho)}_{\text{Second-order Jensen bound}}$$

Is the Bayesian approach an optimal learning strategy?



$$\underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta)] - \mathbb{V}(\rho)}_{\text{Second-order Jensen bound}} \stackrel{\text{PAC-Bayes}}{\lesssim} \underbrace{\mathbb{E}_{\rho(\theta)}[L(\theta, D)] - \hat{\mathbb{V}}(\rho, D) + \frac{KL(\rho, \pi) + \ln \frac{1}{\xi} + \psi_{\pi, \nu}(c, n)}{cn}}_{\text{Second-order PAC-Bayes bound}}$$

PAC²-Variational Inference

- Variational methods for minimizing **second-order PAC-Bayes bounds**,

$$\arg \min_{\rho \in Q} \mathbb{E}_{\rho(\theta)} [L(\theta, D)] - \hat{\mathbb{V}}(\rho, D) + \frac{KL(\rho, \pi)}{n}$$

where Q is a tractable family of densities (i.e. fully factorized Gaussian distribution).

PAC²-Variational Inference

- Variational methods for minimizing **second-order PAC-Bayes bounds**,

$$\arg \min_{\rho \in Q} \mathbb{E}_{\rho(\theta)} [L(\theta, D)] - \hat{\mathbb{V}}(\rho, D) + \frac{KL(\rho, \pi)}{n}$$

where Q is a tractable family of densities (i.e. fully factorized Gaussian distribution).

Variational Inference

- **Standard Variational methods** tries to minimize the first-order PAC-Bayes bound,

$$\arg \min_{\rho \in Q} \mathbb{E}_{\rho(\theta)} [L(\theta, D)] + \frac{KL(\rho, \pi)}{n}$$

Conclusions

- The Bayesian approach does **not** seem to be an **optimal learning strategy**.
- Novel **variational and ensemble learning algorithms**.

`https://github.com/PGM-Lab/PAC2BAYES`