

$$A = \left\{ x \mid x \in \mathbb{H}, \frac{x}{2} < \frac{20}{x} < \frac{x}{1.7} \right\} \Rightarrow A = ?$$

$$\overset{+}{\frac{13}{4}} ? \overset{+}{\frac{11}{6}}$$

$$\underbrace{13 \cdot 6}_{78} > \underbrace{4 \cdot 11}_{44} \Rightarrow \frac{13}{4} > \frac{11}{6}$$

$$-\frac{3}{4} ? -\frac{2}{3} \quad \text{Comparison} \quad \frac{3}{4} ? \frac{2}{3}$$

$$\frac{3 \cdot 3}{9} > \frac{4 \cdot 2}{8} \Rightarrow \frac{3}{4} > \frac{2}{3} \Rightarrow -\frac{3}{4} < -\frac{2}{3}$$

$$x \in \mathbb{H}, x \neq 0 \Rightarrow \frac{x}{2} < \frac{20}{x} \Leftrightarrow x \cdot x < 2 \cdot 20 \Leftrightarrow$$

$$x^2 < 40 \Leftrightarrow \sqrt{x^2} < \sqrt{40} \Leftrightarrow |x| < \sqrt{40} \Leftrightarrow x < \sqrt{40}$$

(because $x > 0 \Rightarrow |x| = x$)

$$40 < 49 \Rightarrow \sqrt{40} < \sqrt{49} \Rightarrow \sqrt{40} < 7 \Rightarrow \boxed{x < 7}$$

$$\overset{+}{\frac{20}{x}} < \overset{+}{\frac{x}{1.7}} \Leftrightarrow 20 \cdot 1.7 < x \cdot x \Leftrightarrow 34 < x^2 \Leftrightarrow$$

$$\sqrt{34} < \sqrt{x^2} \Leftrightarrow \sqrt{34} < |x| \Leftrightarrow \sqrt{34} < x$$

($x > 0$)

$$25 < 34 \Leftrightarrow \sqrt{25} < \sqrt{34} \Leftrightarrow 5 < \sqrt{34} < x \Rightarrow \boxed{5 < x}$$

$$A = \{ x \in \mathbb{H} \mid 5 < x < 7 \} = \{6\}$$

OBS: Dacă $m, m \in \mathbb{N}$ și $m^2 < m < (m+1)^2 \Rightarrow$
 m nu este pătrat perfect.

Între două pătrate perfecte consecutive nu mai avem
niciun pătrat perfect (în \mathbb{N} , \mathbb{N} , naturale).

$$n = 4 \Rightarrow m+1 = 5 \Rightarrow 16, 17, 18, 19, 20, 21, 22, 23, 24, 25$$

4^2 └────────────────────────────────┘ 5^2
nu sunt p.p.

Dem. că $(\forall) m \in \mathbb{N} \Rightarrow m^2 < \underline{m \cdot (m+1)} < (m+1)^2$

Dem: $n^2 = m \cdot m < m \cdot \underline{(m+1)} < (m+1) \cdot \underline{(m+1)} = (m+1)^2$
 $\Rightarrow m^2 < m(m+1) < (m+1)^2$

Dem. că $(\forall) m \in \mathbb{N}^* \Rightarrow \sqrt{m^2 + m} \notin \mathbb{Q}$

$$m^2 + m = m(m+1) \Rightarrow m^2 < m(m+1) < (m+1)^2 \Rightarrow$$

$$\Rightarrow m(m+1) \text{ nu e p.p.} \Rightarrow m^2 + m \text{ nu e p.p.} \Rightarrow$$

$$\underline{\sqrt{m^2 + m} \notin \mathbb{Q}}$$

Dem. că $(\forall) m \in \mathbb{N} \Rightarrow \sqrt{5m+2} \notin \mathbb{Q}$

Dacă $m \in \mathbb{N} \Rightarrow u.c.(m) \in \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\} \Rightarrow$

$$\Rightarrow u.c.(m^2) \in \{0, 1, 4, 9, 6, 5\} \Rightarrow \text{dacă } u.c.(m) \in \{2, 3, 7, 8\} \Rightarrow$$

$$\Rightarrow m \text{ nu e p.p.}$$

$$(\forall) n \in \mathbb{N} \Rightarrow u.c. (5n) \in \{0, 5\} \Rightarrow u.c. (5n+2) \in \{2, 7\}$$

$$\Rightarrow 5n+2 \text{ nu e p.p.} \Rightarrow \sqrt{5n+2} \notin \mathbb{Q}$$

Efectuatii:

$$\mathbb{N} \cap \mathbb{Z} = \mathbb{N}$$

$$\mathbb{N} \cup \mathbb{Q} = \mathbb{Q}$$

$$\mathbb{Z} \cap (\mathbb{R} \setminus \mathbb{Q}) = \emptyset$$

$$\mathbb{N} \setminus \mathbb{Z} = \emptyset$$

$$\mathbb{Q} \setminus (\mathbb{R} \setminus \mathbb{Q}) = \mathbb{Q}$$

$$\mathbb{R} \setminus (\mathbb{R} \setminus \mathbb{Q}) = \mathbb{Q}$$

$$\mathbb{Z} \setminus \mathbb{N} = \{\dots, -4, -3, -2, -1\} = \mathbb{Z}_-$$

$$\text{Stim ca } \mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$$

$$\mathbb{R} = \mathbb{Q} \cup \mathbb{I}$$

$$\mathbb{I} = \mathbb{R} \setminus \mathbb{Q}$$

$$\mathbb{R} \setminus \mathbb{Q} \subset \mathbb{R}$$

$$\underline{A = \{1, 2, 3\}}$$

$$B = \{2\} \quad C = \{3, 4\}$$

$$\underline{D = \{9\}}$$

$$A \cup B = A$$

$$A \cup C = \{1, 2, 3, 4\}$$

$$A \cup D = \{1, 2, 3, 9\}$$

$$A \cap B = B$$

$$A \cap C = \{3\}$$

$$A \cap D = \emptyset$$

$$A - B = \{1, 3\}$$

$$A - C = \{1, 2\}$$

$$A - D = A \quad \text{MULTITII}$$

$$B - A = \emptyset$$

$$C - A = \{4\}$$

$$D - A = D \quad \text{DISJUNCTE}$$

$$\text{Daca } \bar{A} \quad B \subset A \Rightarrow \begin{cases} A \cup B = A \\ B - A = \emptyset \end{cases}$$

$$A \cap B = B$$