

Project Fys3600

Candidate numbers: 1. & 14.

November 8, 2018



Contents

1	Introduction	2
1.1	Theory	3
1.1.1	The Sun:	3
1.1.2	The Magnetosphere:	5
1.1.3	Ionospheric currents	6
2	instruments	8
3	Observations	8
4	Discussion	8
5	Conclusion	8
6	Appendix	8
7	References and acknowledgements	8

1 Introduction

As society moves towards more automatization, accurate, globally available navigational tools are coming in demand. A solution to this is using satellites. There are some problems however. Satellites (and in some cases even earthbound equipment) are susceptible to bad space weather. Having your self- driving car suddenly acting up is not desirable. Predicting when equipment will be in danger of getting hit by high energy particles is therefore of interest.

In this report, we will be looking at data from several instruments measuring properties of the solar wind and the interplanetary magnetic field (IMF) along with auroras and some ionospheric currents. We will also be looking at some potential consequences of the measurements.

1.1 Theory

1.1.1 The Sun:

The root of all our problems is the star in the center of our solar system: the sun.

IMF

The sun, like the earth, has a magnetic field. This magnetic field stretches throughout the solar system, and has fittingly been dubbed the interplanetary magnetic field. What causes this magnetic field is beyond the scope of this report, but it is seemingly quite erratic, changing changing sign and field strength often.

Something more constant about the imf is the general direction of the field lines. Due to the rotation of the sun, the magnetic field lines end up looking like archimedian spirals (or a ballerina skirt in 3 dimensions).




Figure 1: An artistic rendition of the interplanetary magnetic field. With the sun in the middle and rotating, the magnetic field lines end up forming archimedian spirals.

Solar Wind

Accompanying the magnetic field lines is plasma streams. Ejected from the suns corona, these plasma streams of varying velocity and density can also be drawn like figure 1.

Carrying large amount of highly energized ionized particles, the solar wind is of great interest to us. Knowing the density and velocity of the solar wind is necessary in understanding the consequences of its interactions with earth and its satellites.

Coronal mass ejection and solar flares

There are events where the sun sends out relatively huge amounts of plasma

and radiation.

One such event is a solar flare. A solar flare is a massive outburst of radiation happening in the lower corona. Being composed of mostly photons, a solar flare does not follow the magnetic field lines.

Another such event is a coronal mass ejection (CME). During times of magnetic unrest, the sun might eject massive amounts of mass from its corona. Such outbursts are sometimes several times larger than the sun in diameter. The mass ejected follows the magnetic field lines, and ends up causing a spike in the solar wind density, something that can be harmful to instruments on earth.




Figure 2: Artistic size comparison between the sun and a possible cme. As one can see, the cme can be many times larger in size than the sun.

1.1.2 The Magnetosphere:

Like the sun, the earth also has a magnetic field.




Figure 3: The earths magnetosphere. Note that it looks a bit different when the IMF Z component has the same sign as earths (in the figure, they are opposite).

The earths magnetic south pole is currently located on the north pole. This means that the magnetic field goes from the geographical south pole to the geographical north pole. In figure 3, this is indicated by arrows pointing along the field lines.

Magnetic reconnection

We observe many magnetosphere-related events from earth, such as the aurora. As these are not happening all the time, something must happen to trigger them.

When the Z (north/south) direction of the IMF and earths magnetic field points in opposite directions, field lines from earth and field lines from the IMF can break and recouple with eachother. This phenomenon is known as magnetic reconnection.

When magnetic reconnection happens, the field lines "wrap around the earth" in an attempt to regain equilibrium. We can divide this process into different stages of what we call the Dungey cycle.




Figure 5: Some ionospheric currents. Here we see the Birkeland currents stretching out from the polar cusp, the Pedersen current going from dawn to dusk and Hall-currents going from day to night, and back around the cusp. The numbers indicate magnetic local time, with 12 being dayside and 06 being dawn.

One of these is the Birkeland currents which bring plasma in and out of the polar cusps. The cause of this is the interaction between the magnetosphere and the IMF.

As the charged particles in the Birkeland currents approach earth, their density increases. This increases their collisions rate, which in turn causes the ions and electrons to drift apart (since they gyrate in opposite directions). This causes another current in the lower ionosphere: the Pedersen current (the orange and purple lines going along the earth in figure 5).

Since the Pedersen current includes a division in charge, we also end up with an E field. This, along with the B field lines that point towards the polar cusp introduces another current through $E \times B$ drift: the Hall-current which is depicted in green in figure 5.

2 instruments

3 Observations

4 Discussion

We got a bunch of shit here

5 Conclusion

Conclude some shit!

6 Appendix

7 References and acknowledgements