


VOQUEZNA Campaign

Desktop

VOQUEZNA Campaign — Desktop



Learn more about a **FIRST-IN-CLASS PCAB** at ACG



THE POWER TO HELP TREAT GERD¹

GERD=Gastroesophageal Reflux Disease;
PCAB=potassium-competitive acid blocker.
GERD consists of 2 main forms: Non-Erosive GERD and Erosive GERD.²

Dear [Title/First Name/Last Name],

Are your patients suffering from GERD symptoms?

VOQUEZNA[®] (vonoprazan) is a first-in-class PCAB approved for the healing and maintenance of healing of Erosive GERD (Erosive Esophagitis), and now for heartburn relief in Non-Erosive GERD.¹

To learn more about VOQUEZNA, **visit ACG Booth 214**, then head to the Product Theater for an informative clinical presentation by John Erik Pandolfino, MD. Please see details below.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

VOQUEZNA[®] (vonoprazan) is contraindicated in patients with a known hypersensitivity to vonoprazan or any component of VOQUEZNA, or in patients receiving rilpivirine-containing products.

Please see additional Important Safety Information below.

GET THE WHOLE VOQUEZNA STORY AT ACG!

Uncover additional clinical insights at our booth and at the Product Theater presentation by:



John Erik Pandolfino, MD, MSCI
Executive Director, Digestive Health Institute
Division Chief, Gastroenterology and Hepatology
Hans Popper Professor of Medicine
Kenneth C. Griffin Esophageal Center
Northwestern Medicine, Northwestern University


DATE
OCTOBER 27

VENUE
ACG Product Theater
Refreshments provided

TIME
3:45–4:30 pm

[LEARN MORE AT VOQUEZNAPRO.COM](#)

| What makes VOQUEZNA different? See the MOA in action.



| Another difference? Once-daily dosing with or without food.¹



EROSIVE HEALING	EROSIVE MAINTENANCE	NON-EROSIVE HEARTBURN RELIEF
 20 mg once daily For 8 weeks for the healing of Erosive GERD and relief of associated heartburn in adults.	 10 mg once daily For up to 6 months for the maintenance of healing of Erosive GERD and relief of associated heartburn in adults.	 10 mg once daily For 4 weeks for the relief of heartburn associated with Non-Erosive GERD in adults.

Pills not actual size.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS

Presence of Gastric Malignancy: In adults, symptomatic response to therapy with VOQUEZNA does not preclude the presence of gastric malignancy. Consider additional follow-up and diagnostic testing in patients who have a suboptimal response or an early symptomatic relapse after completing treatment with VOQUEZNA. In older patients, also consider endoscopy.

Acute Tubulointerstitial Nephritis: Acute tubulointerstitial nephritis (TIN) has been reported with VOQUEZNA. If suspected, discontinue VOQUEZNA and evaluate patients with suspected acute TIN.

Clostridioides difficile-Associated Diarrhea: Published observational studies suggest that proton pump inhibitors (PPIs) may be associated with an increased risk of *Clostridioides difficile*-associated diarrhea (CDAD), especially in hospitalized patients. VOQUEZNA may also increase the risk of CDAD. Consider CDAD in patients with diarrhea that does not improve. Use the shortest duration of VOQUEZNA appropriate to the condition being treated.

Bone Fracture: Several published observational studies suggest that PPI therapy may be associated with an increased risk for osteoporosis-related fractures of the hip, wrist, or spine, especially in patients receiving high dose (multiple daily doses) and long-term therapy (a year or longer). Bone fracture, including osteoporosis-related fracture, has also been reported with vonoprazan. Use the shortest duration of VOQUEZNA appropriate to the condition being treated. Patients at risk for osteoporosis-related fractures should be managed according to the established treatment guidelines.

Severe Cutaneous Adverse Reactions (SCAR): Severe cutaneous adverse reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported with VOQUEZNA. Discontinue VOQUEZNA at the first signs or symptoms of SCAR or other signs of hypersensitivity and consider further evaluation.

Vitamin B12 (Cobalamin) Deficiency: Long-term use of acid-suppressing drugs can lead to malabsorption of Vitamin B12 caused by hypo- or achlorhydria. Vitamin B12 deficiency has been reported postmarketing with vonoprazan. If clinical symptoms consistent with vitamin B12 deficiency are observed in patients treated with VOQUEZNA, consider further workup.

Hypomagnesemia and Mineral Metabolism: Hypomagnesemia has been reported postmarketing with vonoprazan. Hypomagnesemia may lead to hypocalcemia and/or hypokalemia and may exacerbate underlying hypocalcemia in at-risk patients.

Consider monitoring magnesium levels prior to initiation of VOQUEZNA and periodically in patients expected to be on prolonged treatment, in patients taking drugs that may have increased toxicity in the presence of hypomagnesemia or drugs that may cause hypomagnesemia. Treatment of hypomagnesemia may require magnesium replacement and discontinuation of VOQUEZNA.

Consider monitoring magnesium and calcium levels prior to initiation of VOQUEZNA and periodically while on treatment in patients with a preexisting risk of hypocalcemia. Supplement with magnesium and/or calcium, as necessary. If hypocalcemia is refractory to treatment, consider discontinuing VOQUEZNA.

Interactions with Diagnostic Investigations for Neuroendocrine Tumors: Serum chromogranin A (CgA) levels increase secondary to drug-induced decreases in gastric acidity. The increased CgA level may cause false positive results in diagnostic investigations for neuroendocrine tumors. Temporarily discontinue VOQUEZNA treatment at least 4 weeks before assessing CgA levels and consider repeating the test if initial CgA levels are high.

Fundic Gland Polyps: Use of VOQUEZNA is associated with a risk of fundic gland polyps that increases with long-term use, especially beyond one year. Fundic gland polyps have been reported with vonoprazan in clinical trials and during postmarketing use with PPIs. Most patients who developed fundic gland polyps were asymptomatic and fundic gland polyps were identified incidentally on endoscopy. Use the shortest duration of VOQUEZNA appropriate to the condition being treated.

ADVERSE REACTIONS

Healing of Erosive GERD: The most common adverse reactions (≥2% of patients in the VOQUEZNA arm) include gastritis (3%), diarrhea (2%), abdominal distention (2%), abdominal pain (2%), and nausea (2%).

Maintenance of Healed Erosive GERD: The most common adverse reactions (≥3% of patients in the VOQUEZNA arm) include gastritis (6%), abdominal pain (4%), dyspepsia (4%), hypertension (3%), and urinary tract infection (3%).

Relief of Heartburn Associated with Non-Erosive GERD: The most common adverse reactions (≥2% of patients in the VOQUEZNA arm) include abdominal pain (2%), constipation (2%), diarrhea (2%), nausea (2%), and urinary tract infection (2%).

DRUG INTERACTIONS

VOQUEZNA has the potential for clinically important drug interactions, including interactions with drugs dependent on gastric pH for absorption, drugs that are substrates for certain CYP enzymes, and some diagnostic tests. Avoid concomitant use of VOQUEZNA with atazanavir or nelfinavir. See full Prescribing Information for more details about important drug interactions. Consult the labeling of concomitantly used drugs to obtain further information about interactions with vonoprazan.

USE IN SPECIFIC POPULATIONS

Lactation: Breastfeeding is not recommended during treatment. Because of the potential risk of adverse liver effects shown in animal studies with vonoprazan, advise patients not to breastfeed during treatment with VOQUEZNA.

Renal Impairment: For the healing of Erosive GERD, dosage reduction is recommended in patients with severe renal impairment (eGFR < 30 mL/min).

Hepatic Impairment: For the healing of Erosive GERD, dosage reduction is recommended in patients with moderate to severe hepatic impairment (Child-Pugh Class B and C).

INDICATIONS AND USAGE

VOQUEZNA is a potassium-competitive acid blocker (PCAB) indicated in adults:

- for the healing of all grades of Erosive Esophagitis (Erosive Gastroesophageal Reflux Disease or Erosive GERD) and relief of heartburn associated with Erosive GERD.
- to maintain healing of all grades of Erosive GERD and relief of heartburn associated with Erosive GERD.
- for the relief of heartburn associated with Non-Erosive GERD.

You are encouraged to report suspected adverse reactions by contacting Phathom Pharmaceuticals at 1-888-775-PHAT (7428) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](#).

Please [click here](#) to see full Prescribing Information for VOQUEZNA.

[[Click here](#) to view Rx Drug Pricing Disclosures.]

References: 1. VOQUEZNA. Prescribing Information. Phathom Pharmaceuticals; 2024. 2. Savarino E, de Bortoli N, De Cassan C, et al. The natural history of gastro-esophageal reflux disease: a comprehensive review. *Dis Esophagus*. 2017;30(2):1-9. doi:10.1111/dote.12511

VOQUEZNA Campaign


Mobile

VOQUEZNA Campaign — Mobile



VOQUEZNA
(vonoprazan) tablets 375g

Learn more about a
FIRST-IN-CLASS PCAB at ACG



GERD=Gastroesophageal Reflux Disease;
PCAB=potassium competitive acid blocker
GERD consists of 2 main forms: Non-Erosive GERD and Erosive GERD.¹

Dear [Title/First Name/Last Name],

Are your patients suffering from GERD symptoms?

VOQUEZNA® (vonoprazan) is a first-in-class PCAB approved for the healing and maintenance of healing of Erosive GERD (Erosive Esophagitis), and now for heartburn relief in Non-Erosive GERD.¹

To learn more about VOQUEZNA, visit **ACG Booth 214**, then head to the Product Theater for an informative clinical presentation by John Erik Pandolfino, MD. Please see details below.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

VOQUEZNA® (vonoprazan) is contraindicated in patients with a known hypersensitivity to vonoprazan or any component of VOQUEZNA, or in patients receiving rilpivirine-containing products.

Please see additional Important Safety Information below.

GET THE WHOLE VOQUEZNA STORY
AT ACG!

Uncover additional clinical insights at our booth and at the Product Theater presentation by:



John Erik Pandolfino, MD, MSCI
Executive Director, Digestive Health Institute Division
Chief, Gastroenterology and Hepatology Hans Popper
Professor of Medicine
Kenneth C. Griffin Esophageal Center Northwestern
Medicine, Northwestern University

DATE
OCTOBER 27

PLACE
ACG Product Theater
Philadelphia, Pennsylvania

TIME
3:45–4:30 pm

LEARN MORE AT VOQUEZNAPRO.COM

What makes VOQUEZNA different? See the MOA in action.



Another difference? Once-daily dosing with or without food.¹



EROSIVE HEALING




20 mg once daily
For 8 weeks for the healing of Erosive GERD and relief of associated heartburn in adults.

EROSIVE MAINTENANCE



10 mg once daily
For up to 6 months for the maintenance of healing of Erosive GERD and relief of associated heartburn in adults.

NON-EROSIVE HEARTBURN RELIEF



10 mg once daily
For 4 weeks for the relief of heartburn associated with Non-Erosive GERD in adults.

Pills not actual size.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS

Presence of Gastric Malignancy: In adults, symptomatic response to therapy with VOQUEZNA does not preclude the presence of gastric malignancy. Consider additional follow-up and diagnostic testing in patients who have a suboptimal response or an early symptomatic relapse after completing treatment with VOQUEZNA. In older patients, also consider endoscopy.

Acute Tubulointerstitial Nephritis: Acute tubulointerstitial nephritis (TIN) has been reported with VOQUEZNA. If suspected, discontinue VOQUEZNA and evaluate patients with suspected acute TIN.

Clostridioides difficile-Associated Diarrhea: Published observational studies suggest that proton pump inhibitors (PPIs) may be associated with an increased risk of *Clostridioides difficile*-associated diarrhea (CDAD), especially in hospitalized patients. VOQUEZNA may also increase the risk of CDAD. Consider CDAD in patients with diarrhea that does not improve. Use the shortest duration of VOQUEZNA appropriate to the condition being treated.

Bone Fracture: Several published observational studies suggest that PPI therapy may be associated with an increased risk for osteoporosis-related fractures of the hip, wrist, or spine, especially in patients receiving high dose (multiple daily doses) and long-term therapy (a year or longer). Bone fracture, including osteoporosis-related fracture, has also been reported with vonoprazan. Use the shortest duration of VOQUEZNA appropriate to the condition being treated. Patients at risk for osteoporosis-related fractures should be managed according to the established treatment guidelines.

Severe Cutaneous Adverse Reactions (SCAR): Severe cutaneous adverse reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported with VOQUEZNA. Discontinue VOQUEZNA at the first signs or symptoms of SCAR or other signs of hypersensitivity and consider further evaluation.

Vitamin B12 (Cobalamin) Deficiency: Long-term use of acid-suppressing drugs can lead to malabsorption of Vitamin B12 caused by hypo- or achlorhydria. Vitamin B12 deficiency has been reported postmarketing with vonoprazan. If clinical symptoms consistent with vitamin B12 deficiency are observed in patients treated with VOQUEZNA, consider further workup.

Hypomagnesemia and Mineral Metabolism: Hypomagnesemia has been reported postmarketing with vonoprazan. Hypomagnesemia may lead to hypocalcemia and/or hypokalemia and may exacerbate underlying hypocalcemia in at-risk patients.

Consider monitoring magnesium levels prior to initiation of VOQUEZNA and periodically in patients expected to be on prolonged treatment, in patients taking drugs that may have increased toxicity in the presence of hypomagnesemia or drugs that may cause hypomagnesemia. Treatment of hypomagnesemia may require magnesium replacement and discontinuation of VOQUEZNA.

Consider monitoring magnesium and calcium levels prior to initiation of VOQUEZNA and periodically while on treatment in patients with a pre-existing risk of hypocalcemia. Supplement with magnesium and/or calcium, as necessary. If hypocalcemia is refractory to treatment, consider discontinuing VOQUEZNA.

Interactions with Diagnostic Investigations for Neuroendocrine Tumors: Serum chromogranin A (CgA) levels increase secondary to drug-induced decreases in gastric acidity. The increased CgA level may cause false positive results in diagnostic investigations for neuroendocrine tumors. Temporarily discontinue VOQUEZNA treatment at least 4 weeks before assessing CgA levels and consider repeating the test if initial CgA levels are high.

Fundic Gland Polyps: Use of VOQUEZNA is associated with a risk of fundic gland polyps that increases with long-term use, especially beyond one year. Fundic gland polyps have been reported with vonoprazan in clinical trials and during postmarketing use with PPIs. Most patients who developed fundic gland polyps were asymptomatic and fundic gland polyps were identified incidentally on endoscopy. Use the shortest duration of VOQUEZNA appropriate to the condition being treated.

ADVERSE REACTIONS

Healing of Erosive GERD: The most common adverse reactions (≥2% of patients in the VOQUEZNA arm) include gastritis (3%), diarrhea (2%), abdominal distention (2%), abdominal pain (2%), and nausea (2%).

Maintenance of Healed Erosive GERD: The most common adverse reactions (≥3% of patients in the VOQUEZNA arm) include gastritis (6%), abdominal pain (4%), dyspepsia (4%), hypertension (3%), and urinary tract infection (3%).

Relief of Heartburn Associated with Non-Erosive GERD: The most common adverse reactions (≥2% of patients in the VOQUEZNA arm) include abdominal pain (2%), constipation (2%), diarrhea (2%), nausea (2%), and urinary tract infection (2%).

DRUG INTERACTIONS

VOQUEZNA has the potential for clinically important drug interactions, including interactions with drugs dependent on gastric pH for absorption, drugs that are substrates for certain CYP enzymes, and some diagnostic tests. Avoid concomitant use of VOQUEZNA with atazanavir or nelfinavir. See full Prescribing Information for more details about important drug interactions. Consult the labeling of concomitantly used drugs to obtain further information about interactions with vonoprazan.

USE IN SPECIFIC POPULATIONS

Lactation: Breastfeeding is not recommended during treatment. Because of the potential risk of adverse liver effects shown in animal studies with vonoprazan, advise patients not to breastfeed during treatment with VOQUEZNA.

Renal Impairment: For the healing of Erosive GERD, dosage reduction is recommended in patients with severe renal impairment (eGFR < 30 mL/min).

Hepatic Impairment: For the healing of Erosive GERD, dosage reduction is recommended in patients with moderate to severe hepatic impairment (Child-Pugh Class B and C).

INDICATIONS AND USAGE

VOQUEZNA is a potassium-competitive acid blocker (PCAB) indicated in adults:

- for the healing of all grades of Erosive Esophagitis (Erosive Gastroesophageal Reflux Disease or Erosive GERD) and relief of heartburn associated with Erosive GERD,
- to maintain healing of all grades of Erosive GERD and relief of heartburn associated with Erosive GERD,
- for the relief of heartburn associated with Non-Erosive GERD.

You are encouraged to report suspected adverse reactions by contacting Phathom Pharmaceuticals at 1-888-775-PHAT (7428) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please click [here](#) to see full Prescribing Information for VOQUEZNA.

[Click here](#) to view Rx Drug Pricing Disclosures.

References: 1. VOQUEZNA Prescribing Information. Phathom Pharmaceuticals; 2024. 2. Savarino E, de Bortoli N, De Cassan C, et al. The natural history of gastro-esophageal reflux disease: a comprehensive review. *Dis Esophagus*. 2017;30(2):1-9. doi:10.1111/dote.12511

VOQUEZNA, Phathom Pharmaceuticals, and their respective logos are registered trademarks of Phathom Pharmaceuticals, Inc.
© 2025 Phathom Pharmaceuticals. All rights reserved. XX/XX US-VPZ-XX-XXXX
100 Campus Drive, Suite 102
Florham Park, NJ 07932

Unsubscribe

Contact Us

Privacy Policy

Terms and Conditions

Phathom
PHARMACEUTICALS