

Deep Generative Models

Lecture 5

Roman Isachenko



AI Masters

2024, Summer

Recap of previous lecture

Bayes theorem

$$p(\mathbf{t}|\mathbf{x}) = \frac{p(\mathbf{x}|\mathbf{t})p(\mathbf{t})}{p(\mathbf{x})} = \frac{p(\mathbf{x}|\mathbf{t})p(\mathbf{t})}{\int p(\mathbf{x}|\mathbf{t})p(\mathbf{t})d\mathbf{t}}$$

- ▶ \mathbf{x} – observed variables, \mathbf{t} – unobserved variables (latent variables/parameters);
- ▶ $p(\mathbf{x}|\mathbf{t})$ – likelihood;
- ▶ $p(\mathbf{x}) = \int p(\mathbf{x}|\mathbf{t})p(\mathbf{t})d\mathbf{t}$ – evidence;
- ▶ $p(\mathbf{t})$ – prior distribution, $p(\mathbf{t}|\mathbf{x})$ – posterior distribution.

Posterior distribution

$$p(\theta|\mathbf{X}) = \frac{p(\mathbf{X}|\theta)p(\theta)}{p(\mathbf{X})} = \frac{p(\mathbf{X}|\theta)p(\theta)}{\int p(\mathbf{X}|\theta)p(\theta)d\theta}$$

Recap of previous lecture

Latent variable models (LVM)

$$p(\mathbf{x}|\boldsymbol{\theta}) = \int p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})d\mathbf{z} = \int p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})p(\mathbf{z})d\mathbf{z}.$$

MLE problem for LVM

$$\begin{aligned}\boldsymbol{\theta}^* &= \arg \max_{\boldsymbol{\theta}} \log p(\mathbf{X}|\boldsymbol{\theta}) = \arg \max_{\boldsymbol{\theta}} \sum_{i=1}^n \log p(\mathbf{x}_i|\boldsymbol{\theta}) = \\ &= \arg \max_{\boldsymbol{\theta}} \sum_{i=1}^n \log \int p(\mathbf{x}_i|\mathbf{z}_i, \boldsymbol{\theta})p(\mathbf{z}_i)d\mathbf{z}_i.\end{aligned}$$

Naive Monte-Carlo estimation

$$p(\mathbf{x}|\boldsymbol{\theta}) = \int p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})p(\mathbf{z})d\mathbf{z} = \mathbb{E}_{p(\mathbf{z})}p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) \approx \frac{1}{K} \sum_{k=1}^K p(\mathbf{x}|\mathbf{z}_k, \boldsymbol{\theta}),$$

where $\mathbf{z}_k \sim p(\mathbf{z})$.

Recap of previous lecture

ELBO derivation 1 (inequality)

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \log \int p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta}) d\mathbf{z} \geq \mathbb{E}_q \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} = \mathcal{L}(q, \boldsymbol{\theta})$$

ELBO derivation 2 (equality)

$$\begin{aligned} \mathcal{L}(q, \boldsymbol{\theta}) &= \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} d\mathbf{z} = \int q(\mathbf{z}) \log \frac{p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}) p(\mathbf{x}|\boldsymbol{\theta})}{q(\mathbf{z})} d\mathbf{z} = \\ &= \log p(\mathbf{x}|\boldsymbol{\theta}) - KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \end{aligned}$$

Variational decomposition

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta}) + KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(q, \boldsymbol{\theta}).$$

Recap of previous lecture

Variational lower Bound (ELBO)

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta}) + KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) \geq \mathcal{L}(q, \boldsymbol{\theta}).$$

$$\mathcal{L}(q, \boldsymbol{\theta}) = \int q(\mathbf{z}) \log \frac{p(\mathbf{x}, \mathbf{z}|\boldsymbol{\theta})}{q(\mathbf{z})} d\mathbf{z} = \mathbb{E}_q \log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) - KL(q(\mathbf{z})||p(\mathbf{z}))$$

Log-likelihood decomposition

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathbb{E}_q \log p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) - KL(q(\mathbf{z})||p(\mathbf{z})) + KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$

- ▶ Instead of maximizing incomplete likelihood, maximize ELBO

$$\max_{\boldsymbol{\theta}} p(\mathbf{x}|\boldsymbol{\theta}) \rightarrow \max_{q, \boldsymbol{\theta}} \mathcal{L}(q, \boldsymbol{\theta})$$

- ▶ Maximization of ELBO by variational distribution q is equivalent to minimization of KL

$$\arg \max_q \mathcal{L}(q, \boldsymbol{\theta}) \equiv \arg \min_q KL(q(\mathbf{z})||p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$


Outline

1. Variational autoencoder (VAE)
2. Normalizing flows as VAE model

Outline

1. Variational autoencoder (VAE)
2. Normalizing flows as VAE model

Generative models zoo



Variational autoencoder (VAE)

Final EM-algorithm

- ▶ pick random sample $\mathbf{x}_i, i \sim U[1, n]$.
- ▶ compute the objective:

$$\epsilon^* \sim r(\epsilon); \quad \mathbf{z}^* = \mathbf{g}_\phi(\mathbf{x}, \epsilon^*);$$

$$\mathcal{L}(\phi, \theta) \approx \log p(\mathbf{x}|\mathbf{z}^*, \theta) - KL(q(\mathbf{z}^*|\mathbf{x}, \phi)||p(\mathbf{z}^*)).$$

- ▶ compute a stochastic gradients w.r.t. ϕ and θ


$$\begin{aligned}\nabla_\phi \mathcal{L}(\phi, \theta) &\approx \nabla_\phi \log p(\mathbf{x}|\mathbf{g}_\phi(\mathbf{x}, \epsilon^*), \theta) - \nabla_\phi KL(q(\mathbf{z}|\mathbf{x}, \phi)||p(\mathbf{z})); \\ \nabla_\theta \mathcal{L}(\phi, \theta) &\approx \nabla_\theta \log p(\mathbf{x}|\mathbf{z}^*, \theta).\end{aligned}$$

- ▶ update θ, ϕ according to the selected optimization method (SGD, Adam, etc):

$$\begin{aligned}\phi &:= \phi + \eta \cdot \nabla_\phi \mathcal{L}(\phi, \theta), \\ \theta &:= \theta + \eta \cdot \nabla_\theta \mathcal{L}(\phi, \theta).\end{aligned}$$


Variational autoencoder (VAE)

- ▶ VAE learns stochastic mapping between \mathbf{x} -space, from complicated distribution $\pi(\mathbf{x})$, and a latent \mathbf{z} -space, with simple distribution.
- ▶ The generative model learns a joint distribution $p(\mathbf{x}, \mathbf{z}|\theta) = p(\mathbf{z})p(\mathbf{x}|\mathbf{z}, \theta)$, with a prior distribution $p(\mathbf{z})$, and a stochastic decoder $p(\mathbf{x}|\mathbf{z}, \theta)$.
- ▶ The stochastic encoder $q(\mathbf{z}|\mathbf{x}, \phi)$ (inference model), approximates the true but intractable posterior $p(\mathbf{z}|\mathbf{x}, \theta)$ of the generative model.



Variational Autoencoder

$$\mathcal{L}(\phi, \theta) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \left[\log p(\mathbf{x}|\mathbf{z}, \theta) - \log \frac{q(\mathbf{z}|\mathbf{x}, \phi)}{p(\mathbf{z})} \right] \rightarrow \max_{\phi, \theta}.$$



Variational autoencoder (VAE)

- ▶ Encoder $q(\mathbf{z}|\mathbf{x}, \phi) = \text{NN}_e(\mathbf{x}, \phi)$ outputs $\mu_\phi(\mathbf{x})$ and $\sigma_\phi(\mathbf{x})$.
- ▶ Decoder $p(\mathbf{x}|\mathbf{z}, \theta) = \text{NN}_d(\mathbf{z}, \theta)$ outputs parameters of the sample distribution.




image credit:

<https://lilianweng.github.io/lil-log/2018/08/12/from-autoencoder-to-beta-vae.html>

Outline

1. Variational autoencoder (VAE)
2. Normalizing flows as VAE model

VAE vs Normalizing flows

	VAE	NF
Objective	ELBO \mathcal{L}	Forward KL/MLE
Encoder	stochastic $\mathbf{z} \sim q(\mathbf{z} \mathbf{x}, \phi)$	deterministic $\mathbf{z} = \mathbf{f}_{\theta}(\mathbf{x})$ $q(\mathbf{z} \mathbf{x}, \theta) = \delta(\mathbf{z} - \mathbf{f}_{\theta}(\mathbf{x}))$
Decoder	stochastic $\mathbf{x} \sim p(\mathbf{x} \mathbf{z}, \theta)$	deterministic $\mathbf{x} = \mathbf{g}_{\theta}(\mathbf{z})$ $p(\mathbf{x} \mathbf{z}, \theta) = \delta(\mathbf{x} - \mathbf{g}_{\theta}(\mathbf{z}))$
Parameters	ϕ, θ	$\theta \equiv \phi$

Theorem

MLE for normalizing flow is equivalent to maximization of ELBO for VAE model with deterministic encoder and decoder:

$$p(\mathbf{x}|\mathbf{z}, \theta) = \delta(\mathbf{x} - \mathbf{f}_{\theta}^{-1}(\mathbf{z})) = \delta(\mathbf{x} - \mathbf{g}_{\theta}(\mathbf{z}));$$

$$q(\mathbf{z}|\mathbf{x}, \theta) = p(\mathbf{z}|\mathbf{x}, \theta) = \delta(\mathbf{z} - \mathbf{f}_{\theta}(\mathbf{x})).$$

Normalizing flow as VAE

Proof

1. Dirac delta function property

$$\mathbb{E}_{\delta(\mathbf{x}-\mathbf{y})} \mathbf{f}(\mathbf{x}) = \int \delta(\mathbf{x}-\mathbf{y}) \mathbf{f}(\mathbf{x}) d\mathbf{x} = \mathbf{f}(\mathbf{y}).$$

2. CoV theorem and Bayes theorem:

$$p(\mathbf{x}|\boldsymbol{\theta}) = p(\mathbf{z}) |\det(\mathbf{J}_{\mathbf{f}})|;$$

$$p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}) = \frac{p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) p(\mathbf{z})}{p(\mathbf{x}|\boldsymbol{\theta})}; \quad \Rightarrow \quad p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}) |\det(\mathbf{J}_{\mathbf{f}})|.$$

3. Log-likelihood decomposition

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta}) || p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) = \mathcal{L}(\boldsymbol{\theta}).$$

Normalizing flow as VAE

Proof

ELBO objective:

$$\begin{aligned}\mathcal{L} &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x},\theta)} \left[\log p(\mathbf{x}|\mathbf{z}, \theta) - \log \frac{q(\mathbf{z}|\mathbf{x}, \theta)}{p(\mathbf{z})} \right] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x},\theta)} \left[\log \frac{p(\mathbf{x}|\mathbf{z}, \theta)}{q(\mathbf{z}|\mathbf{x}, \theta)} + \log p(\mathbf{z}) \right].\end{aligned}$$

1. Dirac delta function property:

$$\mathbb{E}_{q(\mathbf{z}|\mathbf{x},\theta)} \log p(\mathbf{z}) = \int \delta(\mathbf{z} - \mathbf{f}_\theta(\mathbf{x})) \log p(\mathbf{z}) d\mathbf{z} = \log p(\mathbf{f}_\theta(\mathbf{x})).$$

2. CoV theorem and Bayes theorem:

$$\mathbb{E}_{q(\mathbf{z}|\mathbf{x},\theta)} \log \frac{p(\mathbf{x}|\mathbf{z}, \theta)}{q(\mathbf{z}|\mathbf{x}, \theta)} = \mathbb{E}_{q(\mathbf{z}|\mathbf{x},\theta)} \log \frac{p(\mathbf{z}|\mathbf{x}, \theta) |\det(\mathbf{J}_\mathbf{f})|}{q(\mathbf{z}|\mathbf{x}, \theta)} = \log |\det \mathbf{J}_\mathbf{f}|.$$

3. Log-likelihood decomposition

$$\log p(\mathbf{x}|\theta) = \mathcal{L}(\theta) = \log p(\mathbf{f}_\theta(\mathbf{x})) + \log |\det \mathbf{J}_\mathbf{f}|.$$

Summary

- ▶ The VAE model is an LVM with two neural network: stochastic encoder $q(\mathbf{z}|\mathbf{x}, \phi)$ and stochastic decoder $p(\mathbf{x}|\mathbf{z}, \theta)$.
- ▶ NF models could be treated as VAE model with deterministic encoder and decoder.