

# Productivity estimation in waterfowl using a non-invasive method

Adrien Tableau\*    Alain Caizergues<sup>†</sup>    Iain Henderson<sup>‡</sup>    Sébastien Reeber<sup>§</sup>  
Matthieu Guillemain<sup>¶</sup>

29 septembre, 2021

---

\*Office Français de la Biodiversité, France - adrien.tableau@ofb.gouv.fr - contribution in data analyses & writing

<sup>†</sup>Office Français de la Biodiversité, France - alain.caizergues@ofb.gouv.fr - contribution in writing

<sup>‡</sup>Animal and Plant Health Agency, United Kingdom - iain.henderson@apha.gov.uk - contribution in data collection

<sup>§</sup>Société Nationale pour la Protection de la Nature, France - sebastien.reeber@snpn.fr - contribution in data collection

<sup>¶</sup>Office Français de la Biodiversité, France - matthieu.guillemain@ofb.gouv.fr - contribution in writing

# Abstract

The response of a waterfowl population to a harvest pressure depends on its capacity to renew. The recruitment, i.e. the number of young adults reproducing for the first time, is a key indicator to describe the renewal of a population, and therefore an essential tool in population management. The productivity, i.e. the number of recruits produced per breeder, is even more informative because it is independent of the population size and allows a comparison over time and between species. The proportion of young adults in a waterfowl population is often estimated from game-hunting samples. However, this proportion, which is a main step in estimating productivity, is not accessible in years without harvest, or with a low harvest rate. Moreover, the age-structure in the harvest samples, commonly called hunting bag, does not necessarily reflect the underlying age-structure of the population. It is often skewed towards juveniles and can lead to an overestimation of the productivity. In waterfowl, adult males usually display brighter colours than juveniles and females. This dichromatism can be characterized and monitored from count surveys, a non-invasive method. Such information can be used to infer the proportion of young adults, and consequently the productivity. In using two populations of ruddy duck, *Oxyura oxyura*, this study develops a Bayesian method to estimate the productivity from count data. To judge the accuracy of this approach, the results are compared to productivity estimates from samples. The adult survival is commonly assessed by capture mark recapture and published in the literature. Since the survival can also be estimated from both the counting and the sampling methods, the consistency of the two approaches is discussed by comparing the survival estimates with the literature values.

## 1 Introduction

### 1.1 Definitions

One defines the productivity of a breeding population as the average number of recruits produced per each breeder (ref). If one considers a species reaching the sexual maturity as early as its first year, all the population is mature just before the reproduction season. The productivity is thus the ratio of the number of young adults at the reproduction season in year  $t$ , i.e. the recruits, to the number of all individuals before the reproduction season in year  $t - 1$ , i.e. the recruits and the adults older than one year.

The population size in year  $t$  is equal to the number of breeders that survived over year  $t - 1$  plus the juveniles produced in year  $t - 1$  that survived until the reproduction period of year  $t$ , i.e. the recruitment in year  $t$ . The growth rate of a population is thus the sum of the productivity and the adult survival rate.

The productivity parameter is complex because it is composed of two sub-parameters: the reproduction success and the juvenile survival.

### The use of productivity in population management

The management of an exploited population generally aims at...

### The variability of the productivity

A waterfowl species that is released in a favourable habitat will naturally expand (malthus). The growth of such a population can be explained by its productivity parameter that is higher than its adult mortality rate. This expansion is limited by the carrying capacity of the habitat (ref). When a waterfowl species is endemic of an ecosystem and evolves in stable environmental conditions, the population size varies around the carrying capacity because the individuals compete for space or/and food (ref), which prevents the population from expanding. This competition commonly induces a lower reproduction success or/and a higher youth mortality (ref). The productivity then decreases because it is dampened by density-dependent effects (holt).

For an endemic species evolving in stable environmental conditions, the productivity and the adult mortality rate are balanced, which explains the stabilization of the number of individuals.

Hunting waterfowl induces adult and juvenile mortality (ref). Paradoxically, this juvenile mortality does not always induce a decrease of productivity (ref). Indeed, one expects that a population with a high adult mortality rate combined to a low productivity would rapidly extinct. However, hunting waterfowl has a long history, and most of these species have persisted over time. The reason is that harvest pressure reduces the competition for space and food. The density-dependent effects affecting the productivity in case of high population density are dampened by the harvest mortality. The exploitation of a population thus increases adult mortality, but also productivity. A balance between these two parameters can be reached if the population is exploited to a constant and moderate harvest rate. The size of a newly exploited population in stable environmental conditions should thus reach a new equilibrium, which is expected to be lower than the size without exploitation.

Productivity is thus a varying parameter that reaches its highest values when no density-dependent effects occurs. Exploring the gradient of productivity, and defining the maximum productivity to predict the response of a population to different harvest rates is a challenge in population management.

## **What part of the productivity gradient should we know for population management?**

### **What is the ideal biological model to explore the gradient of the productivity?**

Hunting waterfowl has a long history, and few species have observed long-term moratorium, so few opportunities to explore the response of the productivity to a gradient of pressure. Alien species colonizing new territories and then heavily controlled are a good model to overcome density-dependent effects and explore the gradient productivity. These species are often introduced during involuntary release events. Since its presence might impact the balance of the colonized ecosystem, it happens that managers take control measures to restrict the growth of these populations or to eradicate them.

In Europe, a population following its maximum growth rate is a perfect study model

### **What are the limitations of the sampling method to estimate the productivity?**

- We cannot explore the productivity when there is no harvest, however, it is crucial to explore what happens when there is no harvest in a scenario where we suggest a moratorium to let the population recover.
- Bias because hunting like any predation action is selective. Often towards the weakest individuals. So overestimation of productivity, and potentially maximum sustainable harvest rate overestimated and consequently depletion of the managed population.

### **Why counting data is a potential tool to estimate the productivity?**

Long time series, easy access, no need of samples. Can only provide subestimation of productivity because if some male recruits have already their male colours, they are considered as adult and not a recruit.

### **What is the methodology of this study?**

Nichols -> il faut compter de la façon dont on demande

idée: estimer l'âge moyen de la pop grâce la vulnérabilité et le sexe ratio chez les adultes

breeding/reproduction success good but not enough because variable survival of juveniles

Introduction:

productivity often defined as  $J_t/At$  but more realistic to defined as  $J_t/At-1$

## 2 Materials & methods

### Few words on the species

The ruddy duck is a species introduced in the United Kingdom in the 50's. The first reproduction was observed in the 60's. The population rapidly grew until reaching about more than 5000 individuals spread over the entire country. A new population set up in France from the 90's, likely because of an arrival of few individuals from the United Kingdom. This population set up around a single wintering spot, the Grand Lieu Lake. Since this species is considered as a major threat to the white-headed duck because of the risk of hybridization (AEWA report), a European plan of eradication has been concluded (EU report). Control measures were taken in both countries from 1999 in the UK and 2004 in France.

### Dichromatism characterization

Pictures of male juvenile in winter Pictures of male in winter

### Time series

The French and the UK wintering populations have been monitored since the presence of the first individuals was validated.

Name	Description
$N_t$	Number of individuals in year $t$
$D_t$	Number of individuals removed in year $t$
$R_t$	Number of recruits in year $t$
$S_t$	Adult survival rate, i.e. proportion of breeder in year $t - 1$ still alive in year $t$
$P_t$	Number of recruits in year $t$ produced per breeder in year $t - 1$

The proportion of males in the adult part of a population at its equilibrium is inferred from the samples.

Indeed, the available French samples were insufficient to get Since the . Age in samples to 2003 to 2020 for the United Kingdom and

$$N_t = N_{t-1} - D_{t-1} + R_t \quad (1)$$

$$N_t = N_{t-1} - (1 - S_t).N_{t-1} + P_t.N_{t-1}$$

$$N_t = S_t.N_{t-1} + P_t.N_{t-1}$$

$$\frac{N_t}{N_{t-1}} = S_t + P_t$$

$$\lambda_t = S_t + P_t \quad (2)$$

$PR_{i,t}$  estimation by the counting method




Figure 1: Evolution of the size of the two Ruddy duck populations in light of the harvest pressure; on time periods without harvest, population growth rates can be estimated (slopes in red for the UK and in blue for France)




Figure 2: Evolution of the size of the two Ruddy duck populations in light of the data time series; on restricted time periods with available data, population growth rates can be estimated (slopes in red for the UK and in blue for France)

## Model

Name	Class	Description
$SM$	Data	Total number of adult males sampled
$SF$	Data	Total number of adult females sampled
$PM$	Parameter	Proportion of males in the adult part of population at its equilibrium
$S_{i,t}$	Data	Number of individuals sampled in population $i$ in year $t$
$SJ_{i,t}$	Data	Number of juveniles sampled in population $i$ in year $t$
$CM_{i,t}$	Data	Cumulated number of type-male individuals counted in population $i$ in year $t$
$C_{i,t}$	Data	Cumulated number of individuals counted in population $i$ in year $t$
$Cmax_{i,t}$	Data	Maximum number of individuals counted in population $i$ in year $t$
$PR_{i,t}$	Parameter	Proportion of recruits in population $i$ in year $t$
$R_{i,t}$	Parameter	Number of recruits in population $i$ in year $t$
$P_{i,t}$	Parameter	Number of recruits in population $i$ in year $t$ per breeder in year $t - 1$
$N0_j$	Parameter	Intercept of the regression model, the $j$ index refers to a restricted time period for a single population
$SD_j$	Parameter	Standard deviation of the regression model, the $j$ index refers to a restricted time period for a single population

Name	Class	Description
$\lambda_j$	Parameter	Population growth rate (in $year^{-1}$ ), the $j$ index refers to a restricted time period for a single population
$S_j$	Parameter	Adult survival rate (in $year^{-1}$ ), the $j$ index refers to a restricted time period for a single population

The proportion of males in the adult part of a population at its equilibrium is inferred from the samples. Indeed, the available French samples were insufficient to get Since the . Age in samples to 2003 to 2020 for the United Kingdom and

$$PM \sim \text{Beta}(SM, SF) \quad (3)$$

$PR_{i,t}$  estimation by the counting method

$$CM_{i,t} \sim \text{Binom}(PM.(1 - PR_{i,t}), C_{i,t}) \quad (4)$$

$PR_{i,t}$  estimation by the sampling method

$$SJ_{i,t} \sim \text{Binom}(PR_{i,t}, S_{i,t}) \quad (5)$$

From  $PR_{i,t}$  estimates,  $R_{i,t}$  and  $P_{i,t}$  are expressed as:

$$R_{i,t} = PR_{i,t}.Cmax_{i,t} \quad (6)$$

$$P_{i,t} = \frac{PR_{i,t}.Cmax_{i,t}}{Cmax_{i,t-1}} \quad (7)$$

The index  $j$  refers to a single population  $i$  and a restricted time period for  $t$ . The evolution of the population size can be related to the time to infer the population growth rate. It is a linear regression on the logarithm scale:

$$\log(Cmax_{i,t}) \sim \text{Norm}(N0_j + \log(\lambda_j).t, SD_j) \quad (8)$$

$$S_j = \lambda_j - \overline{P_{i,t}} \quad (9)$$

## Survival estimation

develop the counting method :

2 ways to check the method :

sampling method? Ok but few data as poorly harvested

realistic survival? if  $> 1$ , estimated productivity is too low to support the maximum growth rate in the range of similar species in litterature -> satisfying

blabla

### 3 Results

blabla




Figure 3: (A) Estimates of productivity, i.e. average number of recruits produced per breeder; (B) Comparison of the productivity values between the two estimation methods; bars define the 95% confidence intervals




Figure 4: (A) Estimates of the population growth rate without harvest; two estimates were produced for the UK population because the growth rate dropped significantly when its size reached more than 1000 individuals; (B) Growth rate estimates, bars define the 95% confidence intervals






Figure 5: (A) Estimates of the population growth rate on time intervals with data availability. This aims at producing survival estimates to discuss the reliability of the methods to estimate productivity; (B) Growth rate estimates, bars define the 95% confidence intervals




Figure 6: (A) Average productivity estimates on the time intervals defined for survival analysis

## 4 Discussion

Productivité: Stabilité grosse pop, variabilité petite pop.

Productivity: drop from 2009

blabla

Nichols -> il faut compter de la façon dont on demande

## Aknowledgements

This work was partly funded by the LIFE Oxyura project (LIFE17 NAT/FR/000942) through the LIFE program. This work was carried on with the impulse of Jean-François Maillard, and Jean-Baptiste Mouronval from the Office Français de la Biodiversité, and Jean-Marc Gillier from the Société Nationale pour la Protection de la Nature. The authors aknowledge all the contributors of the data collection in the UK and in France, especially: **UK names**, Vincent Fontaine, Denis Lacourpaille, Jules Joly, Justin Potier, Valentin Boniface, Alexis Laroche, and Médéric Lortion, **other names**.

## References

## Supplement 1




Figure 7: (A1) Estimates of the number of recruits, i.e. the young adults reproducing for their first time; (B1) estimates of the proportion of recruits in the population at the reproduction period; (A2) & (B2) comparison of the values between the two estimation methods; bars define the 95% confidence intervals

Comments?