

The quest for 3 magic squares of perfect square numbers

Adrian Suter

December 23, 2017

1 The maths behind it

Each component s_i , $i = 1, 2, 3, \dots, 9$ of the magic square has to be a perfect square number. Say $s_i = n_i^2$ for $n_i \in \mathbb{Z}$. From [1] we know that the components need to be such that

$$n_i^2 \equiv 1 \pmod{24} \quad (1)$$

Looking at the smallest perfect square numbers and their modulo 24 (see table 1) we can see some sort of a pattern every 12-th integer. Indeed, as

$$(x + 12)^2 \pmod{24} = (x^2 + 24x + 144) \pmod{24} \quad (2)$$

for any integer x and as

$$(24x + 144) \pmod{24} = 0 \quad (3)$$

we can see that

$$x^2 \pmod{24} = (x + 12)^2 \pmod{24} \quad (4)$$

For $x \in [0, 11]$ there are exactly four perfect squares fulfilling the condition (1). Therefore we can build four infinite series \mathbb{S}_j , $j = 1, 2, 3, 4$ for which all members fulfill the condition. Let there be t any integer, then

$$\mathbb{S}_1 = 1 + 12t = \{1, 13, 25, \dots\} \quad (5)$$

$$\mathbb{S}_2 = 5 + 12t = \{5, 17, 29, \dots\} \quad (6)$$

$$\mathbb{S}_3 = 7 + 12t = \{7, 19, 31, \dots\} \quad (7)$$

$$\mathbb{S}_4 = 11 + 12t = \{11, 23, 35, \dots\} \quad (8)$$

The square numbers of the members of all four sets are the only perfect square numbers that fulfill the condition (1). There are no others.

x	x^2	$x^2 \pmod{24}$	x	x^2	$x^2 \pmod{24}$
0	0	0	12	144	0
1	1	1	13	169	1
2	4	4	14	196	4
3	9	9	15	225	9
4	16	16	16	256	16
5	25	1	17	289	1
6	36	12	18	324	12
7	49	1	19	361	1
8	64	16	20	400	16
9	81	9	21	441	9
10	100	4	22	484	4
11	121	1	23	529	1

Table 1: Some perfect square numbers and their modulo 24.

2 The generator functions

The above four series can be generated using two generator functions $f_+(g)$ and $f_-(g)$ for any integer g .

$$f_+(g) = 6 \cdot g + 1 \quad (9)$$

$$f_-(g) = 6 \cdot g - 1 \quad (10)$$

We will call g the generator number.

3 Program usage

Once the program had been launched, it waits to receive on the standard input a generator string. The generator string consists of a generator number g followed optionally by a $+$ (default) or $-$ character. By hitting enter, the program would start calculating the magic squares for the given generator string and at the end it would print a dash $-$ to the standard output. As soon as this happened, the program is ready for the next generator string. If any valid or almost¹ valid magic square had been detected, the program would write a file containing the most important information to the disk and prints the filename to the standard output.

4 Program flow

The last character of the generator string defines the generator function to be applied. Either it ends by $-$, then the function (10) would be used or it ends by $+$ (default), then the function (9) would be applied.

Let n_5 be this calculated number. Therefore the center cell of our magic square would become

$$s_5 = n_5^2 \quad (11)$$

In the program we would store n_5 in the variable `number` and s_5 in `numberSquared`.

Example

Let the generator string be "141-". That means our program would calculate

$$n_5 = f_-(141) = 6 \cdot 141 - 1 = 845$$

and

$$s_5 = 845^2 = 714\,025.$$

Next we calculate all factor pairs for n_5 , that is we build the set of factor pairs

$$\mathbb{P} = \{(p_1, q_1), (p_2, q_2), \dots, (p_K, q_K)\} \quad (12)$$

where $K = |\mathbb{P}|$ and where

$$p_k \cdot q_k = n_5 \quad \forall k \in [1, K] \subset \mathbb{Z}. \quad (13)$$

Example

The set of factor pairs for our example would be

$$\mathbb{P} = \{(13, 65), (5, 169), (1, 845)\} \quad (14)$$

To be continued...

¹More than six perfect square numbers

Example

```
> 141-
Input: 141, -1
Number: 845, Number^2: 714025
-- Factor Pairs --
13 x 65
5 x 169
1 x 845
-- Arithmetic Progressions --
519841, 714025, 908209 | 194184
508369, 714025, 919681 | 205656
373321, 714025, 1054729 | 340704
207025, 714025, 1221025 | 507000
89401, 714025, 1338649 | 624624
28561, 714025, 1399489 | 685464
25, 714025, 1428025 | 714000
-- Valid Combinations --
(194184, 205656)...
508369 1113865 519841
725497 714025 702553
908209 314185 919681
(194184, 340704)...
373321 1248913 519841
860545 714025 567505
908209 179137 1054729
(194184, 507000)...
207025 1415209 519841
1026841 714025 401209
908209 12841 1221025
(194184, 624624) Skip as a + b >= c
(194184, 685464) Skip as a + b >= c
(194184, 714000) Skip as a + b >= c
(205656, 340704)...
373321 1260385 508369
849073 714025 578977
919681 167665 1054729
(205656, 507000)...
207025 1426681 508369
1015369 714025 412681
919681 1369 1221025
ps,6,141M,1.result
(205656, 624624) Skip as a + b >= c
(205656, 685464) Skip as a + b >= c
(205656, 714000) Skip as a + b >= c
(340704, 507000) Skip as a + b >= c
(340704, 624624) Skip as a + b >= c
(340704, 685464) Skip as a + b >= c
(340704, 714000) Skip as a + b >= c
(507000, 624624) Skip as a + b >= c
(507000, 685464) Skip as a + b >= c
(507000, 714000) Skip as a + b >= c
(624624, 685464) Skip as a + b >= c
(624624, 714000) Skip as a + b >= c
(685464, 714000) Skip as a + b >= c
```

Content of the file ps,6,141M,1.result

845

714025

1 0 1 | 0 1 0 | 1 1 1

207025 1426681 508369 | 1015369 714025 412681 | 919681 1369 1221025

References

- [1] P. Zimmermann, P. Pierrat, and F. Thiriet, “Magic squares of squares.” <http://www.loria.fr/~zimmerma/papers/squares.pdf>.