

Necesarry libraries

```
library(readr)
library(dplyr)

##
## Attaching package: 'dplyr'

## The following objects are masked from 'package:stats':
##
##   filter, lag

## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union
```

Loading dataset


```
dataset <- read_csv("preprocessed_data.csv")

## Rows: 1460 Columns: 81
## -- Column specification -----
## Delimiter: ","
## chr (43): MSZoning, Street, Alley, LotShape, LandContour, Utilities, LotConf...
## dbl (38): Id, MSSubClass, LotFrontage, LotArea, OverallQual, OverallCond, Ye...
##
## i Use 'spec()' to retrieve the full column specification for this data.
## i Specify the column types or set 'show_col_types = FALSE' to quiet this message.

dataset1 = subset(dataset, BedroomAbvGr != 8)

dataset1$ppsqm = dataset1$SalePrice / dataset1$GrLivArea

plot(dataset1$BedroomAbvGr, dataset1$ppsqm)
```



```


quartiles = quantile(dataset1$ppsqm, probs = c(.25, .75), na.rm=FALSE)
IQRppsqm = IQR(dataset1$ppsqm)

lower <- quartiles[1] - 1.5*IQRppsqm
upper <- quartiles[2] + 1.5*IQRppsqm

no_outliers = subset(dataset1, dataset1$ppsqm > lower & dataset1$ppsqm < upper)

plot(no_outliers$BedroomAbvGr, no_outliers$ppsqm)

```



```
dataset2 <- dataset1[names(dataset1) %in% c('ppsqm', 'BedroomAbvGr')]
```


```
no_outliers %>%
  group_by(BedroomAbvGr) %>%
  count() -> dataset3
```

```
dataset3
```


```
## # A tibble: 7 x 2
## # Groups:   BedroomAbvGr [7]
##   BedroomAbvGr     n
##         <dbl> <int>
## 1             0      5
## 2             1     46
## 3             2    347
## 4             3    796
## 5             4    212
## 6             5     20
## 7             6      7
```

```
data1to4 = subset(no_outliers, no_outliers$BedroomAbvGr > 0 & no_outliers$BedroomAbvGr < 5)
```

```
plot(data1to4$ppsqm ~ data1to4$BedroomAbvGr)
```



```
boxplot(ppsqm ~ BedroomAbvGr, data=data1to4)
```



```
require(nortest)
```

```
## Loading required package: nortest
```

```
bartlett.test(data1to4$ppsqm ~ data1to4$BedroomAbvGr)
```

```
##
## Bartlett test of homogeneity of variances
##
## data: data1to4$ppsqm by data1to4$BedroomAbvGr
## Bartlett's K-squared = 17.446, df = 3, p-value = 0.0005722
```

```
lillie.test(data1to4$ppsqm)
```

```
##
## Lilliefors (Kolmogorov-Smirnov) normality test
##
## data: data1to4$ppsqm
## D = 0.017688, p-value = 0.3579
```

```
lillie.test(data1to4$ppsqm[data1to4$BedroomAbvGr == 1])
```

```
##
```

```
## Lilliefors (Kolmogorov-Smirnov) normality test
##
## data: data1to4$ppsqm[data1to4$BedroomAbvGr == 1]
## D = 0.10279, p-value = 0.2586
```

```
lillie.test(data1to4$ppsqm[data1to4$BedroomAbvGr == 2])
```

```
##
## Lilliefors (Kolmogorov-Smirnov) normality test
##
## data: data1to4$ppsqm[data1to4$BedroomAbvGr == 2]
## D = 0.041909, p-value = 0.1456
```

```
lillie.test(data1to4$ppsqm[data1to4$BedroomAbvGr == 3])
```

```
##
## Lilliefors (Kolmogorov-Smirnov) normality test
##
## data: data1to4$ppsqm[data1to4$BedroomAbvGr == 3]
## D = 0.02798, p-value = 0.1365
```

```
lillie.test(data1to4$ppsqm[data1to4$BedroomAbvGr == 4])
```

```
##
## Lilliefors (Kolmogorov-Smirnov) normality test
##
## data: data1to4$ppsqm[data1to4$BedroomAbvGr == 4]
## D = 0.069636, p-value = 0.01426
```

```
aov = aov(data1to4$ppsqm ~ data1to4$BedroomAbvGr)
```

```
summary(aov)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## data1to4$BedroomAbvGr    1   89864    89864   121.7 <2e-16 ***
## Residuals              1399 1033254     739
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```