



LAG-3: impairs T-cell function and can mark exhausted T cells



Role in normal cell

Lymphocyte-activation gene 3 (LAG-3) is an immune checkpoint receptor expressed on the surface of both activated cytotoxic T cells and regulatory T cells (Tregs).¹⁻³ MHC, which presents antigens to T cells, is one of the ligands for LAG-3.¹⁻⁴




Role in cancer cell

Similar to the expression and function of PD-1, repeated exposure to tumor antigen causes an increase in the presence and activity of LAG-3, leading to T-cell exhaustion.^{5,6} T cells co-expressing both LAG-3 and PD-1 may show an even greater degree of exhaustion compared with those expressing LAG-3 alone.⁷


Preclinical evidence

Research is ongoing to understand how dual inhibition of LAG-3 and other checkpoint pathways may synergistically increase T-cell antitumor activity compared with the inhibition of either pathway alone.

To learn more about specific pathways and download additional educational resources, please visit [IOHCP.com](https://www.iohcp.com)



LAG-3: impairs T-cell function and can mark exhausted T cells



Acronyms

APC=antigen-presenting cell; MHC=major histocompatibility complex; PD-1=programmed death receptor-1; TCR=T-cell receptor.

References

1. Huang CT et al. *Immunity*. 2004;21(4):503-513.
2. Baixeras E et al. *J Exp Med*. 1992;176(2):327-337.
3. Deng W-W et al. *Oncoimmunology*. 2016;5(11):e1239005.
4. Huard B et al. *Eur J Immunol*. 1995;25(9):2718-2721.
5. Blackburn SD et al. *Nat Immunol*. 2009;10(1):29-37.
6. Goding SR et al. *J Immunol*. 2013;190(9):4899-4909.
7. Matsuzaki J et al. *Proc Natl Acad Sci U S A*. 2010;107(17):7875-7880.

To learn more about specific pathways and download additional educational resources, please visit [IOHCP.com](https://www.iohcp.com)