



# CTLA-4: negatively regulates long-term immune responses



## Role in normal cell

Cytotoxic T-lymphocyte antigen 4 (CTLA-4) is an immune checkpoint receptor expressed on the surface of activated T cells and Tregs.<sup>1-3</sup>

- Binding of CTLA-4 on cytotoxic T cells to CD80/86 on APCs inhibits T-cell activation<sup>4</sup>
- Continuous expression of CTLA-4 on Tregs is critical for their suppressive activity<sup>2,5</sup>



## Role in cancer cell

Tumor cells utilize the CTLA-4 pathway to suppress initiation of an immune response, decreasing T-cell activation and ability to proliferate into memory T cells.<sup>6,7</sup>

## Preclinical evidence


Preclinical data demonstrate that treatment with antibodies specific for CTLA-4 can restore an immune response through increased activation, accumulation, function, and survival of T cells and memory T cells as well as the depletion of Tregs.<sup>6,8,9</sup>

## Optimizing CTLA-4 blockade


Novel approaches to enhance either the degree or specificity of immune activation with CTLA-4 blockade are under investigation.<sup>10,11</sup>

- One recent approach aims to improve the specificity of CTLA-4 blockade by reducing antibody binding outside of the tumor microenvironment –To localize anti-CTLA-4 activity to the tumor, probody technology has been developed where the antibodies are masked with a peptide that is removed by enzymes that are active primarily in the TME<sup>10-12</sup>
- An alternative approach developed is the generation of a nonfucosylated (NF) form of anti-CTLA-4 antibodies that increases their affinity for FcγR, leading to enhanced Treg depletion by immune-mediated ADCC and increased effector T-cell activation<sup>13,14</sup>

To learn more about specific pathways and download additional educational resources, please visit [IOHCP.com](https://www.iohcp.com)



# CTLA-4: negatively regulates long-term immune responses



## Acronyms

---

ADCC=antibody-dependent cellular cytotoxicity; APC=antigen-presenting cell; FcγR=fragment crystallizable receptor gamma; MHC=major histocompatibility complex; TCR=T-cell receptor; TME=tumor microenvironment; Treg=regulatory T cell.

## References

---

1. Perkins D et al. *J Immunol*. 1996;156(11):4154-4159.
2. Wing K et al. *Science*. 2008;322(5899):271-275.
3. Le Mercier I et al. *Front Immunol*. 2015. doi:10.3389/fimmu.2015.00418
4. Chen L, Flies DB. *Nat Rev Immunol*. 2013;13(4):227-242.
5. Takahashi T et al. *J Exp Med*. 2000;192(2):303-309.
6. Buchbinder EI, Desai A. *Am J Clin Oncol*. 2016;39(1):98-106.
7. Chambers CA et al. *Eur J Immunol*. 1998;28(10):3137-3143.
8. Pedicord VA et al. *Proc Natl Acad Sci U S A*. 2011;108(1):266-271.
9. Simpson TR et al. *J Exp Med*. 2013;210(9):1695-1710.
10. Chen I-J et al. *Sci Rep*. 2017;7(1):11587.
11. Tuve S et al. *Cancer Res*. 2007;67(12):5929-5939.
12. Franssen MF et al. *Clin Cancer Res*. 2013;19(19):5381-5389.
13. Simpson TR et al. *J Exp Med*. 2013;210(9):1695-1710.
14. Satoh M et al. *Expert Opin Biol Ther*. 2006;6(11):1161-1173.

To learn more about specific pathways and download additional educational resources, please visit [IOHCP.com](http://IOHCP.com)