
An Exploration of Multigrid Methods

Avi Schwarzschild and Andres Soto

Numerical Methods for PDEs 4301

Department of Applied Physics and Applied Math
Columbia University

UNIs: aks2203 and ads2206

Abstract

We used both one and two dimensional poisson problems to study multigrid methods for solving partial differential equations. Using iterative solvers for linear systems we show how coarsening the discretization can lead to approximations which converge to the true solution of the PDE with fewer iterations of the solver.

1 Motivation

The motivation for using coarser grids, while convergence analysis shows that finer grids should lead to more accurate approximations, comes from an observation about waves, discretizations, and aliasing. The iterative methods, often referred to as smoothers, smooth out error such that in the earlier iterations it is the high frequency components of the error that disappear. As the algorithms sweep more times the error gets smoother, containing lower frequencies and tending toward zero. The trouble with the classical iterative methods like this is that low frequencies in the error can take many iterations to smooth. In general these iterative methods are $\mathcal{O}(n^2)$. Thus, some improvement in speed is desired.

From the Shannon Sampling Theorem we know that to retain all of the wave information, we need the discretization to have just over two points per wavelength.¹ The implication of this on our work is that the highest frequency in the error is determined by the mesh grid. Knowing this, we can coarsen the grid so that we have fewer sample points of the error function and then the low frequencies will be among the higher ones still contained in the coarse-grid error. By smoothing the error on this coarser grid we can eliminate more components of the error than without the coarsening. We leave the explanation and implementation of this for later sections.

2 One Dimensional Problem

For the one dimensional problem we chose to demonstrate the Poisson problem with Dirichlet boundary conditions as follows:

$$\begin{aligned}\frac{d^2u}{dx^2} &= f(x), \quad x \in (a, b), \\ u(a) &= \alpha, \\ u(b) &= \beta.\end{aligned}\tag{1}$$

The values of $f(x)$, a , b , α , and β are specified below for two different examples.

¹C. E. Shannon, "Communications in the presence of noise", Proc. IRE, vol. 37, pp. 10-21, Jan. 1949.

2.1 Implementation

2.2 Results

3 Two Dimensional Problem

3.1 Implementation

3.2 Results

4 Error Analysis

4.1 Multigrid

4.2 Gauss-Seidel

4.3 Successive Over Relaxation

5 Conclusion