## A strategy for sample retrieval and possible onboard biosafety controls: Perspectives

Y. Takano<sup>1</sup>, H. Yano<sup>2</sup>, Y. Sekine<sup>1,3</sup>, R. Funase<sup>3</sup>, K. Takai<sup>1,2</sup>

Issues of planetary protection are generally handled by the space community, although these issues often come to the attention of various academic communities and the public (e.g., [1, 2]). Since the establishment of the Panel on Plantary Protection (PPP) and the Committee on Space Rsearch (COSPAR), an international consensus has emerged regarding the development and promulgation of planetary protection knowledge and policies, and regarding plans for mitigating the harmful effects of biological contamination on Earth. The investigation of biological quarantine for planetary protection against both forward and back-contamination has been discussed from the viewpoint of risk management and public consensus, in the context of further planetary exploration. However, selection of a candidate location for initial quarantine, especially for materials with high biosafety levels, is problematic due to the potential risk of biological back-contamination and the difficulty of obtaining public consensus in the host countries of the sample recovery site.

To resolve key issues related to extraterrestrial sample-return projects, we suggest that international waters (i.e., areas of oceans, seas, and waters outside of national jurisdiction; Figure 1) are a meaningful option for the location of sample retrieval, likewise the pioneering Apollo missions in 1960's. To conduct an initial investigation of onboard biological control, we propose application of a BSL laboratory on a developed research vessel operating in international waters. According to the United Nations Convention on the Law of the Sea (UNCLOS), international waters are defined as all waters beyond national boundaries with freedom of navigation and also freedom of scientific research (see, Article 87: Freedom of the High Seas). We think that the international waters are the most likely place for the future public consensus of onboard quarantine because of the potentially minimum risk of back-contamination in the ocean environments and the most rapid and convincing processing of the subsequent scientific research. On this basis, we propose potential onboard protocols for the initial biological control of future sample-return missions (e.g., [3], [4]).

## **Acknowledgements:**

This research was supported by the JAMSTEC and JAXA cooperative agreement (FY2012). The authors

would like to thank Dr. P. Tsou for discussions and scientific advice. The authors also thank the reviewing panels on the Decadal Survey proposal (Y. Sekine [4]) of the Japanese Society of Planetary Science (JSPS).

## **References:**

- [1] Rummel, J.D. (2001) Planetary exploration in the time of astrobiology: protecting against biological contamination. *PNAS*, 98, 2128-2131.
- [2] Nicholson, W.L., Schuerger, A.C., Race, M.S., (2009) Migrating microbes and planetary protection. *TRENDS in Microbiology*, 17, 389-392.
- [3] Tsou, P., Brownlee, D.E., McKay, C.P., Anbar, A.D., Yano, H., Altwegg, K., Beegle, L.W., Dissly, R., Strange, N.J., Kanik, I. (2012) LIFE: Life Investigation For Enceladus A sample return mission concept in search for evidence of life. *Astrobiology*, 12, 730-742.
- [4] Sekine, Y., Takano, Y., Yano, H., Funase, R., Takai, K., Ishihara, M., Shibuya, T., Tachibana, S., Kuramoto, K., Yabuta, H., Kimura, J., Furukawa, Y., (2013) Exploration of Enceladus' water-rich plumes toward understanding of chemistry and biology of the interior ocean. *Transactions of the Japan Society for Aeronautical and Space Sciences, Aerospace Technology Japan*, submitted.




Figure 1. Global distribution of international waters.

<sup>&</sup>lt;sup>1</sup>Japan Agency for Marine-Earth Science & Technology (JAMSTEC),

<sup>&</sup>lt;sup>2</sup>Japan Aerospace Exploration Agency (JAXA), <sup>3</sup>University of Tokyo,