



This is not about
pathologizing faith.

A Note on Framing

This work focuses on harmful sexual messaging, not religion itself.

- Faith can be a source of meaning, identity, ethics, and belonging.
- Clinical concern arises when sexual teachings become fear-based, shame-based, coercive, or rigid.
- The goal is not to remove values.
- The goal is to help clients distinguish chosen values from inherited fear.



What did you learn
about sex before you
had language for it?

Early Sexual Messages

Childhood sexual messages are often absorbed before they are questioned. Explore:

What was said directly?

What was implied?

Who taught you what was "pure," "dangerous," or "wrong"?

What messages still live in your body or relationship?

Clinical note: The earliest sexual scripts often become automatic adult reactions.

THE SHAME WE CARRY

2 / 8

Guilt says,
"I did something wrong."

Shame says,
"I am wrong."

Guilt vs. Shame

Guilt and shame may look similar but function differently.

- Guilt can be connected to behavior or values.
- Shame attaches to identity and worth.
- Sexual shame may create avoidance, shutdown, secrecy, or distress.
- Therapy can help clients move from self-condemnation toward integration.



The body may remember
what the mind
has outgrown.

Embodied Responses

Clients may cognitively reject rigid sexual teachings but still experience embodied responses.

- Tension or pain
- Dissociation
- Low desire
- Anxiety during intimacy
- Difficulty receiving pleasure

Clinical note: Do not rush insight into behavior change.

Safety often has to be rebuilt through the body.



The pathway from
teaching to distress
is rarely linear.

A Clinical Map


Childhood sexual teachings



Internalized sexual scripts



Shame, guilt, fear, or confusion



Clinical repair, agency, and relational safety



Avoidance, low desire, pain, dissociation, or relational conflict



How it shows up
in the bedroom.

COMMON PRESENTATIONS

Rigid sexual scripts may appear as:

- Low desire or desire discrepancy
- Sexual avoidance
- Pain or tension
- Guilt after pleasure
- Difficulty initiating
- Dissociation or "checking out"
- Conflict around roles, expectations, or morality

CLINICAL NOTE:

The presenting issue may be sexual, but the wound may be relational, spiritual, embodied, or systemic.



The couple is not
the problem.
The script may be.

Externalizing the Script

MFTs can help couples externalize inherited sexual scripts.

An MFT can support:

- Naming the script
- Reducing blame
- Increasing emotional safety
- Improving sexual communication
- Clarifying shared and individual values
- Renegotiating intimacy with consent and care



Clinical repair
begins with choice.

Pathways to Repair

Meaningful care may include:

- Family-of-origin exploration
- Values clarification
- Shame reduction
- Emotion regulation
- Attachment and safety work
- Consent-centered intimacy conversations
- Collaboration with sex therapy, pelvic health, trauma, or medical providers

Reflection question:

What would intimacy look like if it were guided by safety, consent, values, and connection rather than fear?