

Introduction to extreme event analysis

Stéphane Girard

Inria Grenoble Rhône-Alpes

<http://mistis.inrialpes.fr/~girard>

June 2024



Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data
- 5 Bibliography

Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data
- 5 Bibliography


Extreme rainfall in Montpellier, France, 2014. Three hours of rainfall = 50% of mean annual rainfall (source <http://www.dailymail.co.uk>).



© NICOLAS GAILLARD/APERCU/SIPA/REX

Nidd river (England)


The flows of the Nidd river have been measured every three months during 38.5 years. This results in $n = 154$ measures, the maximum observed flow being about $300m^3/s$.



Nidd river (England)

From these data, it is possible to compute some exceedance probabilities such as:


$$\mathbb{P}(X \geq 160) \simeq nb(X_i \geq 160)/n = 11/154 = 1/14$$



Nidd river (England)

From these data, it is possible to compute some exceedance probabilities such as:

$$\mathbb{P}(X \geq 255) \simeq nb(X_i \geq 255)/n = 3/154$$




Nidd river (England)

From these data, it is possible to compute some exceedance probabilities

... but not extreme ones!

$$\mathbb{P}(X \geq 500) \simeq nb(X_i \geq 500)/n = 0$$



Statistical formulation

- Let X_1, \dots, X_n be n independent and identically distributed (iid) random variables with cumulative distribution function (cdf) F and survival function $\bar{F} = 1 - F$.
- The maximum is denoted by $X_{n,n} = \max(X_1, \dots, X_n)$. The cdf of $X_{n,n}$ is given by $\mathbb{P}(X_{n,n} \leq x) = F^n(x)$.
- As illustrated in the previous slides, the estimation of $\bar{F}(s) = \mathbb{P}(X > s)$ for $s > X_{n,n}$ is a non-trivial problem which is referred to as the **estimation of small tail probabilities**. As shown in the next slides, the two classical statistical approaches are ill-adapted.

Nonparametric approach

- $\bar{F}(s) = \mathbb{P}(X \geq s)$ is estimated by the proportion of observations above the threshold s :

$$\hat{\bar{F}}_n(s) = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{X_i \geq s\},$$


which is referred to as the **empirical survival function**.

- However, as illustrated on the Nidd river: $\hat{\bar{F}}_n(s) = 0$ if $s > X_{n,n}$.

Parametric approach

- Suppose *a priori* a parametric model for the survival function $\bar{F} \in \{\bar{F}_\theta, \theta \in \Theta\}$,
- Estimate θ by $\hat{\theta}_n$ and estimate $\bar{F}(s)$ by $\bar{F}_{\hat{\theta}_n}(s)$.
- However, a good fit on the sample does not necessarily lead to a good modeling above the maximum.
- This phenomena is illustrated on the next figure: Even though the Student t_ν distribution converges to a $\mathcal{N}(0, 1)$ distribution as $\nu \rightarrow \infty$, their large quantiles remain different.

Illustration



Horizontally: p . Vertically: relative error between the quantile of order p estimated from $\mathcal{N}(0, 1)$ and Student t_4 , Student t_8 , Student t_{12} distributions.

Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data
- 5 Bibliography

Goal

- The **Central Limit Theorem** (CLT) provides the asymptotic distribution of the **mean**

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

of a sample $\{X_1, \dots, X_n\}$ of iid random variables:

$$\sqrt{n} \left(\frac{\bar{X}_n - \mathbb{E}(X)}{\sigma(X)} \right) \xrightarrow{d} \mathcal{N}(0, 1).$$

Equivalently, in terms of cdf,

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\sqrt{n} \left(\frac{\bar{X}_n - \mathbb{E}(X)}{\sigma(X)} \right) \leq x \right) = \Phi(x),$$

where Φ is the cdf of the $\mathcal{N}(0, 1)$ distribution.

- The **Extreme-Value Theorem** is a similar result for the **maximum**.

Extreme-Value Theorem

Under mild conditions on F , there exist three parameters a_n , b_n and γ such that:

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\frac{X_{n,n} - a_n}{b_n} \leq x \right) = H_\gamma(x),$$

with


$$H_\gamma(x) = \exp \left(-(1 + \gamma x)_+^{-1/\gamma} \right)$$

and where $y_+ = \max(0, y)$.

- H_γ is the **Extreme-Value Distribution (EVD)**,
- γ is the **extreme-value parameter**,
- a_n are b_n normalization parameters.

We note $F \in \text{MDA}(H_\gamma)$.

Illustration on a Gaussian distribution



Comparison between $H_\gamma(x)$, $\mathbb{P}\left(\frac{X_{n,n}-a_n}{b_n} \leq x\right)$ with $n = 10$ and $\mathbb{P}\left(\frac{X_{n,n}-a_n}{b_n} \leq x\right)$ with $n = 100$.

Extreme-Value Distribution

In practice,

$$\mathbb{P}(X_{n,n} \leq x) \simeq H_\gamma \left(\frac{x - a_n}{b_n} \right),$$


on has a three-parameter distribution

- a_n is a **location parameter**, playing the role of $\mathbb{E}(X)$ in the TCL,
- b_n is a **scale parameter**, playing the role of $\sigma(X)/\sqrt{n}$ in the TCL,
- γ is a **shape parameter**, without any equivalent in the TCL.

Three cases appear (and thus three kind of distributions):

- If $\gamma > 0$, F is said to belong to the maximum domain of attraction of **Fréchet**.
- If $\gamma = 0$, F is said to belong to the maximum domain of attraction of **Gumbel**.
- If $\gamma < 0$, F is said to belong to the maximum domain of attraction of **Weibull**.

Extreme-Value Distribution



Examples of densities associated with the Extreme-Value Distribution ($\gamma = 0$, $\gamma = 1$ and $\gamma = -1$).

Maximum Domain of Attractions (MDA)

MDA	Gumbel $\gamma = 0$	Fréchet $\gamma > 0$	Weibull $\gamma < 0$
Tail	Light	Heavy	Short
Distribution	Gaussian Exponential Lognormal Gamma Weibull	Cauchy Pareto Student	Uniform Beta

Application to extrapolation

Since $\mathbb{P}(X_{n,n} \leq x) = F^n(x)$, the Extreme-Value Theorem yields an approximation of $F(x)$ for large values of x ,

$$F(x) = 1 - \bar{F}(x) \simeq H_\gamma^{1/n} \left(\frac{x - a_n}{b_n} \right).$$


Taking the logarithm, it follows

$$\log(1 - \bar{F}(x)) \simeq \frac{1}{n} \log H_\gamma \left(\frac{x - a_n}{b_n} \right).$$

Since x is large, $\bar{F}(x)$ is small, a first order expansion of $\log(1 + u)$ entails

$$\bar{F}(x) \simeq -\frac{1}{n} \log H_\gamma \left(\frac{x - a_n}{b_n} \right) = \frac{1}{n} \left[1 + \gamma \left(\frac{x - a_n}{b_n} \right) \right]^{-1/\gamma}.$$

Illustration on a Gaussian distribution



Comparison between $\bar{F}(x)$, $-\frac{1}{n} \log H_\gamma \left(\frac{x-a_n}{b_n} \right)$ with $n = 10$ and $-\frac{1}{n} \log H_\gamma \left(\frac{x-a_n}{b_n} \right)$ with $n = 100$.

Here, the theoretical values of a_n , b_n and γ have been used (they are known for the standard Gaussian distribution). **In practice, they are unknown (since F is unknown) and have to be estimated.**

Estimation of EVD parameters

Let $\{Y_1, \dots, Y_k\}$ be a sample of k iid random variables from the extreme-value cdf $H_{\gamma,a,b}$. This starting point may be a problem since, usually, one does not observe directly a sample of maxima. They have to be extracted from an initial sample $\{X_1, \dots, X_n\}$ with arbitrary cdf F . The common practice is to divide the data into k blocks and to extract the maxima of each block (*block-maxima method*).

Two drawbacks:

- Loss of information (n data $\rightarrow k$ data),
- The maxima are not exactly distributed from an EVD distribution.

Estimation of EVD parameters

The cdf of the EVD distribution is given by:

$$H_{\gamma,a,b}(x) \stackrel{\text{def}}{=} H_{\gamma} \left(\frac{x-a}{b} \right) = \exp \left\{ - \left[1 + \gamma \left(\frac{x-a}{b} \right) \right]_+^{-1/\gamma} \right\}.$$

Maximum-likelihood estimators:

- not closed-form (optimization procedure).
- Consistency if $\gamma > -1$,
- Asymptotic normality if $\gamma > -1/2$ (Fisher information matrix available).

Selection of a sample of maxima




Illustration on Nidd data. Three block sizes are considered, the block maxima are depicted in red.

Application to Nidd data

```
> # Fit GEV to the data in nidd.annual, the annual maximum water
> # levels of the River Nidd, using the "BFGS" optimization method
> gev_nidd <- gev(nidd.annual, method = "BFGS", control = list(maxit = 500))
```

\$data

```
[1] 65.08 65.60 75.06 76.22 78.55 81.27 86.93 87.76 88.89 90.28
[11] 91.80 91.80 92.82 95.47 100.40 111.54 111.74 115.52 131.82 138.72
[21] 148.63 149.30 151.79 153.04 158.01 162.99 172.92 179.12 181.59 189.04
[31] 213.70 226.48 251.96 261.82 305.75
```

\$par.ests

	xi	sigma	mu
	0.321221	36.154177	103.118249

\$varcov

	[,1]	[,2]	[,3]
[1,]	0.04758274	-0.4142656	-0.7770124
[2,]	-0.41426557	43.6098796	35.7316149
[3,]	-0.77701236	35.7316149	58.0116406


MLE estimation of the GEV parameters with R using the *block-maxima method* on Nidd data. Annual blocks are considered so that $k = 35$ block-maxima are obtained. The estimated parameters are provided as well as the asymptotic covariance matrix.

Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data
- 5 Bibliography

Excess

Let X be random variable with cdf F . For all $u \in \mathbb{R}$ and $x \geq 0$, the cdf F_u defined by $F_u(x) = \mathbb{P}(X - u \leq x | X > u)$ is called the cdf of Y , the excess of X above u .



Generalized Pareto Distribution

The Generalized Pareto Distribution (GPD) with parameters $\gamma \in \mathbb{R}$ and $\sigma > 0$ is defined by the cdf

$$G_{\gamma,\sigma}(y) = 1 - \left(1 + \gamma \frac{y}{\sigma}\right)_+^{-1/\gamma}.$$

Particular cases:

- If $\gamma = 0$, then $G_{0,\sigma}$ is the cdf of $\text{Exp}(\sigma)$,
- If $\gamma = -1$, then $G_{-1,\sigma}$ is the cdf of $\text{Unif}[0, \sigma]$,
- If $\gamma > 0$, then $G_{\gamma,\sigma}$ is the cdf of a translated Pareto distribution.

Pickands theorem

$F \in \text{MDA}(H_\gamma)$ if and only if there exists a positive function $a(\cdot)$ such that

$$\lim_{u \rightarrow \infty} \sup_{y > 0} |F_u(y) - G_{\gamma, a(u)}(y)| = 0.$$

Interpretation:

- The excess is asymptotically distributed from a GPD if and only if the maximum is asymptotically distributed from an EVD.
- In both cases, the shape parameter γ is the extreme-value index.

Application to extrapolation

From Pickands theorem, we have, for all $y \geq 0$,

$$\bar{F}_u(y) = \frac{\bar{F}(u+y)}{\bar{F}(u)} \simeq \bar{G}_{\gamma,\sigma}(y),$$

where $\bar{G}_{\gamma,\sigma}$ is the survival function of the GPD. The change of variable $x = u + y$ yields for all $x \geq u$:

$$\bar{F}(x) \simeq \bar{F}(u) \bar{G}_{\gamma,\sigma}(x - u).$$

Finally, introducing $\alpha = \mathbb{P}(X > u) = \bar{F}(u)$, it follows

$$\bar{F}(x) \simeq \alpha \bar{G}_{\gamma,\sigma}(x - \bar{F}^{-1}(\alpha)).$$

We thus obtain an approximation of the small tail probabilities:

$$\bar{F}(x) \simeq \alpha \left[1 + \gamma \left(\frac{x - \bar{F}^{-1}(\alpha)}{\sigma} \right) \right]^{-1/\gamma}.$$

Comparison between EVD and GPD approaches

For both approaches, the approximations are the same:

$$\bar{F}(x) \simeq \alpha \left[1 + \gamma \left(\frac{x - \bar{F}^{-1}(\alpha)}{\sigma} \right) \right]^{-1/\gamma} \quad (\text{GPD})$$

$$\bar{F}(x) \simeq \frac{1}{n} \left[1 + \gamma \left(\frac{x - a_n}{b_n} \right) \right]^{-1/\gamma} \quad (\text{EVD})$$

There are three parameters to be estimated:

- the extreme value index γ (shape parameter),
- a scale parameter σ (GPD) or b_n (EVD),
- a position parameter $\bar{F}^{-1}(\alpha)$ (GPD) or a_n (EVD).

Estimation of GPD parameters

Let $\{Y_1, \dots, Y_k\}$ be a sample of k iid random variables from the Generalized Pareto cdf $G_{\gamma, \sigma}$. Similarly to EVD case, this starting point may be a problem since, usually, one does not observe directly a sample of excesses. They have to be extracted from an initial sample $\{X_1, \dots, X_n\}$ with arbitrary cdf F . The common practice is to choose a number of k excesses and select

$$\{Y_1, \dots, Y_k\} := \{X_{n-k+1,n} - X_{n-k,n}, \dots, X_{n,n} - X_{n-k,n}\}$$

(*peaks over threshold method*).

Advantages / drawbacks:

- The excesses are not exactly distributed from a GPD distribution,
- Dependence issues,
- $\alpha = k/n$, the position parameter $\bar{F}^{-1}(k/n)$ is estimated by the empirical quantile $X_{n-k,n}$. It only remains to estimate γ and σ (by Maximum likelihood for instance).

Selection of a sample of excesses




Illustration on Nidd data. Three numbers of excesses k are considered. The associated thresholds are depicted in red.

Illustration on Nidd data

```
> # Fit GPD to the data in nidd.thresh
> gpdnidd <- gpd(nidd.thresh, 100)

$data
 [1] 189.02 115.52 119.28 162.99 102.92 143.06 153.04 149.30 116.77 131.82
[11] 107.97 104.19 261.82 110.48 181.59 104.19 158.01 172.92 179.12 213.70
[21] 111.74 100.40 104.19 151.79 111.54 148.63 251.96 121.73 107.58 108.14
[31] 131.92 138.72 133.06 257.62 123.71 157.12 305.75 226.48 110.98

$threshold
[1] 100

$p.less.thresh
[1] 0.7467532

$n.exceed
[1] 39


$par.ests
      xi      beta
0.003508321 50.608623759

$varcov
      [,1]      [,2]
[1,] 0.04562003 -2.303872
[2,] -2.30387212 182.476944
```

MLE estimation of the GPD parameters with R. The threshold $u = 100$ is considered so that $k = 39$ excesses are obtained. The estimated parameters are provided as well as the asymptotic covariance matrix.

In practice, the choice of k is difficult ...

- If k is small, $\hat{\gamma}(k)$ is based on few observations, it has therefore a **large variance**.
- If k is large, then $X_{n-k,n}$ is no longer in the distribution tail and the approximation of the excesses distribution by a GPD is no longer true, $\hat{\gamma}(k)$ has a **large bias**.



Estimation of γ on three samples of size $n = 500$ from a t_2 distribution ($\gamma = 1/2$). Horizontally: k . Vertically: $\hat{\gamma}(k)$ for $k = 1, \dots, 200$.

Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data**
- 5 Bibliography

Time dependent extremes

Data $X_1(\cdot), \dots, X_n(\cdot)$ are stochastic processes.

Probabilistic problem: Find two sequences of continuous functions $a_n(\cdot)$ and $b_n(\cdot)$ such that

$$\left\{ \frac{\max_{i=1, \dots, n} X_i(t) - a_n(t)}{b_n(t)} \right\}_{t \in [0,1]}$$

converges in distribution to a stochastic process $\{Y(t)\}_{t \in [0,1]}$. From the univariate theory, the margin at point $t \in [0,1]$ is an EVD with shape parameter $\gamma(t)$. In practice, for each time $t \in [0,1]$:

$$\mathbb{P} \left(\max_{i=1, \dots, n} X_i(t) \leq x \right) \simeq H_{\gamma(t)} \left(\frac{x - a_n(t)}{b_n(t)} \right).$$

Two statistical problems:

- Estimations of functions $\gamma(\cdot)$, $a_n(\cdot)$ and $b_n(\cdot)$.
- Estimation of the dependence structure

$$\mathbb{P} \left(\max_{i=1, \dots, n} X_i(t_1) \leq x_1 \cap \max_{i=1, \dots, n} X_i(t_2) \leq x_2 \right).$$

Not discussed here, cf Chapter 9 of de Haan & Ferreira (2006).

Estimation of extreme-value parameters as functions of covariates

Parametric approach

- Chavez-Demoulin & Davison (2005): GEV + linear model of the trend (expansion on a splines basis), penalized maximum likelihood.

Semi-parametric approach

- Davison & Ramesh (2000) and Hall & Tajvidi (2000): GEV/GPD + nonlinear model of the trend (local polynomial smoothing).

Nonparametric approach

- Daouia *et al* (2013): Domain of attraction condition + kernel estimators of extreme-value parameters.
- Einmahl *et al* (2016) and Ahmad *et al* (2020): restriction to heavy-tails and $\gamma(\cdot)$ constant.

Outline

- 1 The problem
- 2 Introduction to extreme-value analysis: Maxima
- 3 Introduction to extreme-value analysis: Excesses
- 4 Non iid data
- 5 Bibliography

S. Coles (2001), An introduction to statistical modeling of extreme values, Springer.

P. Embrechts, C. Klüppelberg and T. Mikosch (1997), Modelling extremal events, Springer.

L. de Haan and A. Ferreira (2006), Extreme Value Theory, Springer.

Time dependent extremes

- Ahmad, A., Deme, E., Diop, A., Girard, S. and Usseglio-Carleve, A. (2020). Estimation of extreme quantiles from heavy-tailed distributions in a location-dispersion regression model, *Electronic Journal of Statistics*, 14, 4421–4456, 2020.
- Chavez-Demoulin, V. and Davison, A. (2005). Generalized additive modelling of sample extremes. *Journal of the Royal Statistical Society C*, 54, 207–222.
- Daouia, A., Gardes, L. and Girard, S. (2013). On kernel smoothing for extremal quantile regression. *Bernoulli*, 19, 2557–2589.
- Davison, A. and Ramesh, N. (2000). Local likelihood smoothing of sample extremes. *Journal of the Royal Statistical Society B*, 62, 191–208.
- Einmahl, J., de Haan, L. and Zhou, C. (2016). Statistics of heteroscedastic extremes. *Journal of the Royal Statistical Society B*, 78, 31–51.
- Hall, P. and Tajvidi, N. (2000). Nonparametric analysis of temporal trend when fitting parametric models to extreme-value data. *Statistical Science*, 15, 153–167.