

PHY101: Introduction to Physics I

Monsoon Semester 2023

Lecture 31

Department of Physics, School of Natural Sciences,
Shiv Nadar Institution of Eminence, Delhi NCR

Previous Lecture

Monoatomic gas

Involvement of temperature

This Lecture


Diatomeric gas system

**Energy associated with degrees
of freedom**

Equipartition theorem

Diatomeric gas and its dynamics

Example



Molecules of neon, a monatomic gas, have only one atom.

Molecules of diatomic gases, such as hydrogen, have two atoms.

Rotational and vibrational motion



Total kinetic energy of a diatomic gas molecule

DIATOMIC MOLECULE

- Consider now a gas containing a diatomic molecule. The diatomic molecule consists of atoms A and B. The molecule has a center of mass motion as well as the internal motion (rotation, vibration).
- If \vec{v}_A and \vec{v}_B are the corresponding velocities and the gas is at temperature T , then

$$\left\langle \frac{1}{2}m_A v_A^2 \right\rangle = \frac{3}{2}kT = \left\langle \frac{1}{2}m_B v_B^2 \right\rangle.$$

- Thus the total kinetic energy of A+B (the molecule)= $3kT$
(Here $k \equiv k_B$, the Boltzmann constant)

Kinetic energy of the center of mass

- Now consider the Center of Mass (COM) of the molecule,
- The velocity of the COM is $\vec{V}_{CM} = \frac{m_A \vec{v}_A + m_B \vec{v}_B}{m_A + m_B}$.
- Therefore,
$$\begin{aligned} V_{CM}^2 &= \vec{V}_{CM} \cdot \vec{V}_{CM} \\ &= \left(\frac{m_A \vec{v}_A + m_B \vec{v}_B}{m_A + m_B} \right) \cdot \left(\frac{m_A \vec{v}_A + m_B \vec{v}_B}{m_A + m_B} \right) \\ &= \frac{1}{M^2} (m_A^2 v_A^2 + m_B^2 v_B^2 + 2m_A m_B \vec{v}_A \cdot \vec{v}_B) \end{aligned}$$
- where $M = m_A + m_B$
$$\Rightarrow \frac{1}{2} M V_{CM}^2 = \frac{1}{M} \left(\frac{1}{2} m_A^2 v_A^2 + \frac{1}{2} m_B^2 v_B^2 + m_A m_B \vec{v}_A \cdot \vec{v}_B \right)$$

Average kinetic energy of the center of mass

Take average,

$$\left\langle \frac{1}{2} M V_{CM}^2 \right\rangle = \frac{1}{M} \left(m_A \left\langle \frac{1}{2} m_A v_A^2 \right\rangle + m_B \left\langle \frac{1}{2} m_B v_B^2 \right\rangle + m_A m_B \langle \vec{v}_A \cdot \vec{v}_B \rangle \right)$$

Now, as already seen, for individual motions of A and B,

$$\left\langle \frac{1}{2} m_A v_A^2 \right\rangle = \frac{3}{2} kT = \left\langle \frac{1}{2} m_B v_B^2 \right\rangle.$$

Thus,

$$\begin{aligned} \left\langle \frac{1}{2} M V_{CM}^2 \right\rangle &= \frac{1}{M} \left((m_A + m_B) \frac{3}{2} kT + m_A m_B \langle \vec{v}_A \cdot \vec{v}_B \rangle \right) \\ \Rightarrow \left\langle \frac{1}{2} M V_{CM}^2 \right\rangle &= \frac{3}{2} k_B T + \frac{m_A m_B}{m_A + m_B} \langle \vec{v}_A \cdot \vec{v}_B \rangle \end{aligned}$$

Assumptions valid for monoatomic gases and diatomic gases

- If A and B were part of independent gases we could have intuitively set $\langle \vec{v}_A \cdot \vec{v}_B \rangle = 0$, as we did earlier. However, in this case we must analyze this term as A and B are bound as a molecule.
- Even in this case, for the relative velocity $\vec{w} = \vec{v}_A - \vec{v}_B$, we can safely assume that

$$\langle \vec{w} \cdot \vec{V}_{CM} \rangle = 0$$

$$\Rightarrow \left\langle (\vec{v}_A - \vec{v}_B) \cdot \left(\frac{m_A \vec{v}_A + m_B \vec{v}_B}{m_A + m_B} \right) \right\rangle = 0$$

$$\Rightarrow \langle (\vec{v}_A - \vec{v}_B) \cdot (m_A \vec{v}_A + m_B \vec{v}_B) \rangle = 0$$

Final form of the average kinetic energy of the center of mass

$$\Rightarrow \langle m_A \vec{v}_A \cdot \vec{v}_A + m_B \vec{v}_A \cdot \vec{v}_B - m_A \vec{v}_B \cdot \vec{v}_A - m_B \vec{v}_B \cdot \vec{v}_B \rangle = 0$$

$$\Rightarrow \langle m_1 v_1^2 - m_2 v_2^2 + (m_2 - m_1) \vec{v}_1 \cdot \vec{v}_2 \rangle = 0$$

$$\Rightarrow \langle m_1 v_1^2 \rangle - \langle m_2 v_2^2 \rangle + (m_2 - m_1) \langle \vec{v}_1 \cdot \vec{v}_2 \rangle = 0$$

Now, $\left\langle \frac{1}{2} m_1 v_1^2 \right\rangle = \left\langle \frac{1}{2} m_2 v_2^2 \right\rangle$, thus

[A, B and 1, 2 in the subscripts represents the same quantity]

$$(m_2 - m_1) \langle \vec{v}_1 \cdot \vec{v}_2 \rangle = 0$$

$$\Rightarrow \langle \vec{v}_1 \cdot \vec{v}_2 \rangle = 0$$

So this relation holds in this case also. Therefore,


$$\left\langle \frac{1}{2} M V_{CM}^2 \right\rangle = \frac{3}{2} kT$$

Thus kinetic energy associated with the center of mass motion is $\frac{3}{2} kT$.


Average kinetic energy of the internal motion of the molecule

- Total average kinetic energy of the molecule = $3kT$.
- Average kinetic energy of center of mass = $\frac{3}{2}kT$
(The bodily motion of the entire molecule, regarded as a single particle of mass $M = m_1 + m_2$)
- Therefore, the average kinetic energy associated with the internal motion of the molecule
(rotation+vibration)= $3kT - \frac{3}{2}kT = \frac{3}{2}kT$

Degrees of freedom: motion of the diatomic molecule



Translational Motion



Rotational Motion



Vibrational Motion

Energy associated with each type of motion

Consider for example a diatomic molecule.

The total average kinetic energy has to be $2 \times \frac{3kT}{2} = 3kT$.

This has contribution:

- $\frac{3kT}{2}$ from the center of mass motion.
- $2 \times \frac{kT}{2}$ from rotational degrees of freedom.
(There are two axes of rotation)
- $\frac{kT}{2}$ from vibrational degree of freedom.

Energies associated with polyatomic molecule

Recall the relation

$$PV = (\gamma - 1)U = NkT$$

For a gas with monatomic molecules, $\gamma = 5/3$. In this case the internal energy $U = 3NkT/2$ is entire due to kinetic energy.

However, for polyatomic molecules the contribution to U comes from the total average kinetic energy and potential energy associated with the bound atoms in the molecule.

Total energy: Kinetic + Potential

At low energies the vibration of the molecule can be very well be modeled by a harmonic potential. In this case we know that the average kinetic and potential energies are same.

Thus potential energy associated with vibration = $\frac{kT}{2}$.

Therefore total internal energy of the gas,

$$U = N \left(3kT + \frac{kT}{2} \right) = \frac{7NkT}{2}.$$

Thus $(\gamma - 1)U = (\gamma - 1) \frac{7NkT}{2} = NkT$, giving $\gamma = 9/7$.

In absence of energy contribution from vibrational motion

- However, usually the vibrational degree of freedom is not excited (for example at the room temperature: *Quantum mechanical reason*), and then the total contribution to the total internal energy is $U = \frac{5NkT}{2}$, which gives $\gamma = \frac{7}{5} = 1.4$
- *The terrestrial air is primarily made up of diatomic gases (~78% nitrogen (N_2) and ~21% oxygen (O_2)) and at standard conditions it can be considered to be an ideal gas. The above value of 1.4 is highly consistent with the measured values 1.403, 1.400, 1.401 at 0°C, 20°C, 100°C respectively.*

EQUIPARTITION THEOREM

If there are f degrees of freedom, then each of them has an associated contribution $\frac{kT}{2}$ to the internal energy.

Thus,

$$U = N \frac{f kT}{2}, \text{ and thus } (\gamma - 1)U = (\gamma - 1) \frac{f N kT}{2} = N kT$$

gives

$$\gamma = 1 + 2/f.$$

Problem set for practice

- ❖ Reference book: Fundamentals of Physics, 5th Edition, (Chapter 20, problem 20-4)
- ❖ Reference book: Physics for Scientists and Engineers, 9th Edition (Chapter 21, problem 21-1, 21-2, 21-3, 21-4)

What is average translational kinetic energy of the oxygen and nitrogen molecule at room temperature?

Hints: Average translational kinetic energy depends only on the temperature and not on the nature of molecule.

Thus, using the following relationship the answer is 0.039 eV

$$E = \frac{3kT}{2}$$