

Series Solution of O.D.E

Dr Tracey Berry



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON



Apply Separation of Variables to Schrodinger Equ, Laplace,
in cylindrical, spherical coordinates...

-> get O.D.E in corresponding coordinates

Usually non-constant co-efficients (like Euler)

Usually no simple variable sub to make to make constant coeff

Need new approaches...

1) Series Solution

2) Frobenius

Old method...



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON

To solve..

$$y'' + y = 0 .$$

General solution is: (only because we know already..!)

$$y(x) = A \cos x + B \sin x .$$

For a more general solution: use a power series..

$$y(x) = \sum_{n=0}^{\infty} a_n x^n .$$

Solve with a power series..



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON

Substitute

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

Into the D.E equation.. $y'' + y = 0$.

$$y'(x) = \sum_{n=0}^{\infty} a_n n x^{n-1},$$

$$y''(x) = \sum_{n=0}^{\infty} a_n n(n-1) x^{n-2}.$$

Gives:

$$\sum_{n=0}^{\infty} a_n n(n-1) x^{n-2} + \sum_{n=0}^{\infty} a_n x^n = 0.$$

Solve with a power series..



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON

$$\sum_{n=0}^{\infty} a_n n(n-1)x^{n-2} + \sum_{n=0}^{\infty} a_n x^n = 0 .$$

Use the trick.. let $m=n-2$


Terms with $m=-2, -1$ are 0 anyway.

$$\sum_{m=-2}^{\infty} a_{m+2}(m+2)(m+1)x^m + \sum_{n=0}^{\infty} a_n x^n = 0 .$$

So now let $m=n$ and rewrite

$$\sum_{n=0}^{\infty} [a_{n+2}(n+2)(n+1) + a_n] x^n = 0 .$$

All coefficients must = 0, for all terms.


$$0 = \sum_{n=0}^{\infty} 0 \times x^n$$



Gives the recurrence relation for the coefficients:

$$a_{n+2} = -\frac{a_n}{(n+1)(n+2)}.$$

Relates odd and even series....

$$a_2 = -\frac{a_0}{2 \times 1},$$

$$a_3 = -\frac{a_1}{3 \times 2},$$

$$a_4 = -\frac{a_2}{4 \times 3} = \frac{a_0}{4 \times 3 \times 2 \times 1},$$

$$a_5 = -\frac{a_3}{5 \times 4} = -\frac{a_1}{5 \times 4 \times 3 \times 2}.$$



Pattern is:

$$a_n = \begin{cases} \frac{(-1)^{n/2}}{n!} a_0, & n \text{ even,} \\ \frac{(-1)^{(n-1)/2}}{n!} a_1, & n \text{ odd.} \end{cases}$$

Relates odd and even series....

$$y(x) = a_0 \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \right) + a_1 \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right) .$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots ,$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots .$$

Same solution as before!! 😊

Summary



ROYAL
HOLLOWAY
UNIVERSITY
OF LONDON

This lecture:

Series Solutions to solve D.E

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

Next lecture:

Frobenius Method