



Evaluating scoring streaks and game excitement using in-match trend estimation

Claus Thorn Ekstrøm and Andreas Kryger Jensen

March 24th 2021

Biostatistics, Department of Public Health
University of Copenhagen


1. Introduction (CE)
2. Latent dynamics GP Regression (AKJ)
3. The 2019–2020 NBA season (CE)
4. Conclusion

Introduction (CE)

Introduction

Motivating example

Game development in the final match of the NBA 2019–2020 season between Los Angeles Lakers and Miami Heat at October 11, 2020. Positive values \Rightarrow LA Lakers are leading.



How exciting was this match?

Latent dynamics GP Regression (AKJ)

Gaussian Process regression with latent dynamics

$$\otimes_m \mid \Psi_m, \mathbf{t}_m \sim H(\otimes_m \mid \Psi_m)$$

$$d_m(t) \mid \otimes_m \sim \mathcal{GP}(\mu_{\beta_m}(t), C_{\theta_m}(s, t))$$

$$D_m(t_{mi}) \mid d_m(t_{mi}), t_{mi}, \otimes_m \stackrel{iid}{\sim} N(d_m(t_{mi}), \sigma_m^2)$$

$$\begin{bmatrix} d_m(s) \\ d'_m(t) \\ d''_m(u) \end{bmatrix} \mid \otimes_m \sim \mathcal{GP} \left(\begin{bmatrix} \mu_{\beta_m}(s) \\ \mu'_{\beta_m}(t) \\ \mu''_{\beta_m}(u) \end{bmatrix}, \begin{bmatrix} C_{\theta_m}(s, s') & \partial_2 C_{\theta_m}(s, t) & \partial_2^2 C_{\theta_m}(s, u) \\ \partial_1 C_{\theta_m}(t, s) & \partial_1 \partial_2 C_{\theta_m}(t, t') & \partial_1 \partial_2^2 C_{\theta_m}(t, u) \\ \partial_1^2 C_{\theta_m}(u, s) & \partial_1^2 \partial_2 C_{\theta_m}(u, t) & \partial_1^2 \partial_2^2 C_{\theta_m}(u, u') \end{bmatrix} \right)$$

Gaussian Process regression with latent dynamics

The posterior distribution of the latent processes conditional on the hyper-parameters is the multivariate Gaussian process

$$\begin{bmatrix} d_m(s) \\ d'_m(t) \\ d''_m(u) \end{bmatrix} \mid \mathcal{D}_m, \otimes_m \sim \mathcal{GP} \left(\begin{bmatrix} \mu_{d_m}(s) \\ \mu_{d'_m}(t) \\ \mu_{d''_m}(u) \end{bmatrix}, \begin{bmatrix} \Sigma_{d_m}(s, s') & \Sigma_{d_m d'_m}(s, t) & \Sigma_{d_m d''_m}(s, u) \\ \Sigma_{d'_m d_m}(t, s) & \Sigma_{d'_m}(t, t') & \Sigma_{d'_m d''_m}(t, u) \\ \Sigma_{d''_m d_m}(u, s) & \Sigma_{d''_m d'_m}(u, t) & \Sigma_{d''_m}(u, u') \end{bmatrix} \right)$$

Gaussian Process regression with latent dynamics

The posterior distribution of the latent processes conditional on the hyper-parameters is the multivariate Gaussian process

$$\begin{bmatrix} d_m(s) \\ d'_m(t) \\ d''_m(u) \end{bmatrix} \mid \mathcal{D}_m, \otimes_m \sim \mathcal{GP} \left(\begin{bmatrix} \mu_{d_m}(s) \\ \mu_{d'_m}(t) \\ \mu_{d''_m}(u) \end{bmatrix}, \begin{bmatrix} \Sigma_{d_m}(s, s') & \Sigma_{d_m d'_m}(s, t) & \Sigma_{d_m d''_m}(s, u) \\ \Sigma_{d'_m d_m}(t, s) & \Sigma_{d'_m}(t, t') & \Sigma_{d'_m d''_m}(t, u) \\ \Sigma_{d''_m d_m}(u, s) & \Sigma_{d''_m d'_m}(u, t) & \Sigma_{d''_m}(u, u') \end{bmatrix} \right)$$

Where the components evaluated at any \mathbf{t}^* are given by

$$\mu_{d_m}(\mathbf{t}^*) = \mu_{\beta_m}(\mathbf{t}^*) + C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} (\mathbf{D}_m - \mu_{\beta_m}(\mathbf{t}_m))$$

$$\mu_{d'_m}(\mathbf{t}^*) = \mu'_{\beta_m}(\mathbf{t}^*) + \partial_1 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} (\mathbf{D}_m - \mu_{\beta_m}(\mathbf{t}_m))$$

$$\mu_{d''_m}(\mathbf{t}^*) = \mu''_{\beta_m}(\mathbf{t}^*) + \partial_1^2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} (\mathbf{D}_m - \mu_{\beta_m}(\mathbf{t}_m))$$

$$\Sigma_{d_m}(\mathbf{t}^*, \mathbf{t}^*) = C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

$$\Sigma_{d'_m}(\mathbf{t}^*, \mathbf{t}^*) = \partial_1 \partial_2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - \partial_1 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} \partial_2 C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

$$\Sigma_{d''_m}(\mathbf{t}^*, \mathbf{t}^*) = \partial_1^2 \partial_2^2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - \partial_1^2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} \partial_2^2 C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

$$\Sigma_{d_m d'_m}(\mathbf{t}^*, \mathbf{t}^*) = \partial_2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} \partial_2 C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

$$\Sigma_{d_m d''_m}(\mathbf{t}^*, \mathbf{t}^*) = \partial_2^2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} \partial_2^2 C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

$$\Sigma_{d'_m d''_m}(\mathbf{t}^*, \mathbf{t}^*) = \partial_1 \partial_2^2 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}^*) - \partial_1 C_{\theta_m}(\mathbf{t}^*, \mathbf{t}_m) \left(C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I \right)^{-1} \partial_2^2 C_{\theta_m}(\mathbf{t}_m, \mathbf{t}^*)$$

Trend Direction Index (TDI)

$$\begin{aligned}\text{TDI}_m(t \mid \otimes_m) &= P(d'_m(t) > 0 \mid \mathcal{D}_m, \otimes_m) \\ &= \frac{1}{2} + \frac{1}{2} \text{Erf} \left(\frac{\mu_{d'_m}(t)}{2^{1/2} \Sigma_{d'_m}(t, t)^{1/2}} \right)\end{aligned}$$

Excitement Trend Index

$$\begin{aligned}\text{ETI}_m \mid \otimes_m &= \mathbb{E} \left[\# \{t \in \mathcal{I}_m : d'_m(t) = 0\} \mid \mathcal{D}_m, \otimes_m \right] \\ &= \int_{\mathcal{I}_m} \int_{-\infty}^{\infty} |v| f_{d'_m(t), d''_m(t)}(0, v \mid \mathcal{D}_m, \otimes_m) dv dt \\ &= \int_{\mathcal{I}_m} d\text{ETI}_m(t \mid \otimes_m) dt\end{aligned}$$

Excitement Trend Index

?

$$d\text{ETI}_m(t \mid \otimes_m) = \lambda_m(t) \phi \left(\frac{\mu_{d'_m}(t)}{\Sigma_{d'_m}(t, t)^{1/2}} \right) \left(2\phi(\zeta_m(t)) + \zeta_m(t) \text{Erf} \left(\frac{\zeta_m(t)}{2^{1/2}} \right) \right)$$

$$\lambda_m(t) = \frac{\Sigma_{d''_m}(t, t)^{1/2}}{\Sigma_{d'_m}(t, t)^{1/2}} (1 - \omega_m(t)^2)^{1/2}, \quad \omega_m(t) = \frac{\Sigma_{d'_m d''_m}(t, t)}{\Sigma_{d'_m}(t, t)^{1/2} \Sigma_{d''_m}(t, t)^{1/2}}$$
$$\zeta_m(t) = \frac{\mu_{d'_m}(t) \Sigma_{d''_m}(t, t)^{1/2} \omega_m(t) \Sigma_{d'_m}(t, t)^{-1/2} - \mu_{d''_m}(t)}{\Sigma_{d''_m}(t, t)^{1/2} (1 - \omega_m(t)^2)^{1/2}}$$

$$\mu_{\beta_m}(t) = \beta_m, \quad C_{\theta_m}(s, t) = \alpha_m^2 \exp\left(-\frac{(s-t)^2}{2\rho_m^2}\right)$$

The hyper-parameters are given the following prior distribution

$$H(\otimes_m \mid \Psi_m) = H(\beta_m \mid \Psi_{\beta_m})H(\alpha_m \mid \Psi_{\alpha_m})H(\rho_m \mid \Psi_{\rho_m})H(\sigma_m \mid \Psi_{\sigma_m})$$

with

$$\beta_m \sim T_4\left(\widehat{\beta_m^{\text{ML}}}, 5\right), \alpha_m \sim T_4^+\left(\widehat{\alpha_m^{\text{ML}}}, 5\right), \rho_m \sim T_4^+\left(\widehat{\rho_m^{\text{ML}}}, 5\right), \sigma_m \sim T_4^+\left(\widehat{\sigma_m^{\text{ML}}}, 5\right)$$

Super-script ML indicates the marginal maximum-likelihood estimate (empirical Bayes).

Empirical bayes

To find the empirical Bayes estimate of the hyper-parameters we maximize the marginal log-likelihood function. This can be written as

$$\begin{aligned}\log L(\otimes_m \mid \mathcal{D}_m) &\propto -\frac{1}{2} \log |C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I| \\ &\quad - \frac{1}{2} (\mathbf{D}_m - \mu_{\beta_m}(\mathbf{t}_m))^T [C_{\theta_m}(\mathbf{t}_m, \mathbf{t}_m) + \sigma_m^2 I]^{-1} (\mathbf{D}_m - \mu_{\beta_m}(\mathbf{t}_m))\end{aligned}$$

The solution $\widehat{\otimes}_m^{\text{ML}} = \arg \sup_{\otimes} \log L(\otimes \mid \mathcal{D}_m)$ can be obtained by numerical optimization.

Our model is implemented in Stan and for each match we ran 4 chains for 50,000 iterations with half of them for warm-up.





The 2019–2020 NBA season (CE)

The 2019–2020 NBA season



We applied our method for all 1143 matches from the 2019–2020 NBA basketball season.

We disregarded overtime.

Final match between LA Lakers and Miami Heat




Distribution of the 1143 median posterior ETIs




Season minimum ETI

Dallas Mavericks vs. Golden State Warriors, ETI = 0.19 (season minimum)




Season 25% percentile ETI

Oklahoma City Thunder vs. Washington Wizards, ETI = 7.5 (season 25% percentile)




Season median ETI

Los Angeles Clippers vs. Charlotte Hornets, ETI = 10.12 (season median)




Season 75% percentile ETI

Chicago Bulls vs. Detroit Pistons, ETI = 13.56 (season 75% percentile)



Season maximum ETI

New Orleans Pelicans vs. Utah Jazz, ETI = 26.23 (season maximum)




An algorithm for collapsing a factor into sub-groups

Clustering results

	Average	SD	2.5%	50%	97.5%	Group
New Orleans Pelicans	11.67	4.91	3.10	11.59	23.54	A
Washington Wizards	11.36	4.46	2.98	11.63	18.28	A
San Antonio Spurs	11.31	5.31	3.52	9.88	23.44	A
Oklahoma City Thunder	11.26	5.32	1.68	11.55	23.74	A
Portland Trail Blazers	11.21	4.46	3.31	10.82	18.48	A
Los Angeles Lakers	11.16	4.43	3.61	10.52	19.79	A
Philadelphia 76ers	11.09	4.70	2.73	10.28	21.53	A
Minnesota Timberwolves	11.09	4.28	4.01	10.69	19.46	A
Utah Jazz	10.96	5.44	3.09	9.76	23.54	A
Memphis Grizzlies	10.94	4.59	3.04	10.36	20.84	A
Cleveland Cavaliers	10.90	4.50	4.93	10.39	22.36	A
Toronto Raptors	10.69	4.44	3.48	10.33	21.79	B
Houston Rockets	10.67	4.12	3.52	10.74	19.17	B
Chicago Bulls	10.66	3.98	2.95	10.65	17.49	B
Boston Celtics	10.61	4.66	2.81	10.19	20.92	B
Orlando Magic	10.56	4.89	2.72	10.41	21.01	B
Milwaukee Bucks	10.55	4.36	2.77	9.59	18.13	B
Los Angeles Clippers	10.54	4.14	4.05	10.72	20.56	B
Miami Heat	10.44	4.29	3.28	9.91	18.54	B
Golden State Warriors	10.40	4.58	3.37	10.24	18.72	B
Indiana Pacers	10.37	3.99	3.31	10.22	18.54	B
Brooklyn Nets	10.33	4.57	3.07	10.06	18.89	C
Dallas Mavericks	10.27	4.99	3.00	9.04	22.09	C
Atlanta Hawks	10.26	4.35	2.50	9.75	18.31	C
Phoenix Suns	10.25	4.32	3.78	9.29	19.81	C
Denver Nuggets	10.16	4.39	3.42	9.59	19.61	C
New York Knicks	10.11	4.05	2.84	9.87	18.47	C
Sacramento Kings	9.94	4.19	3.66	9.30	18.93	C
Detroit Pistons	9.86	4.00	3.73	9.44	17.08	C
Charlotte Hornets	9.26	3.74	1.96	9.22	15.95	D

Team specific seasonal SD vs. average ETIs



Conclusion

Conclusion