

THE COMBINATORICS OF FAREY WORDS AND THEIR TRACES

ALEX ELZENAAR, GAVEN MARTIN, AND JEROEN SCHILLEWAERT

ABSTRACT. We introduce a family of 3-variable “Farey polynomials” that are closely connected with the geometry and topology of 3-manifolds and orbifolds as they can be used to produce concrete realisations of the boundaries and local coordinates for one complex dimensional deformation spaces of Kleinian groups. As such, this family of polynomials has a number of quite remarkable properties. We study these polynomials from an abstract combinatorial viewpoint, including a recursive definition extending that which is known in the literature for the special case of manifolds, even beyond what the geometry predicts. We also present some intriguing examples and conjectures which we would like to bring to the attention of researchers interested in algebraic combinatorics and hypergeometric functions.

The results in this paper additionally provide a practical approach to various classification problems for rank-two subgroups of $\mathrm{PSL}(2, \mathbb{C})$ since they, together with other recent work of the authors, make it possible to provide certificates that certain groups are discrete and free, and effective ways to identify relators.

1. INTRODUCTION

We introduce a fascinating family of 3-variable polynomials $\Phi_{p/q}^{a,b}(z)$ defined for all $a, b \in \mathbb{N} \cup \{\infty\}$ (though the geometric meaning of these polynomials only makes sense for a and b that are integers that are at least 2 or ∞ and excluding the case $a = b = 2$) and all rational numbers $p/q \in [0, 1]$ assumed to be in reduced form $\gcd(r, s) = 1$. The goal of this paper is to study the abstract combinatorial properties of these polynomials, which we call **Farey polynomials**. These polynomials have been studied before in the case $a = b = \infty$, since they appear naturally in studies of the Riley slice of Schottky space, a much studied model for one complex dimensional deformation spaces of Kleinian groups; even in that case many of our results or their proofs seem to be new.

We will now explain the mathematical background behind this paper in more detail. A **Kleinian group** is a discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$; these groups have been intensively studied for a long time in association with hyperbolic geometry and conformal geometry [Bea83; Mar16; Mas87]. The **closed Riley slice** is the deformation space of Kleinian groups which are free on two parabolic generators. After normalisation, every group on two parabolic generators may be written as

$$\Gamma = \left\langle X = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, Y = \begin{bmatrix} 1 & 0 \\ z & 1 \end{bmatrix} \right\rangle;$$

(AE) MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, LEIPZIG, GERMANY

(GM) INSTITUTE FOR ADVANCED STUDY, MASSEY UNIVERSITY, AUCKLAND, NEW ZEALAND

(JS) DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF AUCKLAND, NEW ZEALAND

E-mail addresses: aelz176@aucklanduni.ac.nz, g.j.martin@massey.ac.nz,

j.schillewaert@auckland.ac.nz.

2020 *Mathematics Subject Classification.* 11B37, 11B39, 11B57, 20F36, 57K20 (primary), 32G15, 30F40, 33C45, 41A10 (secondary).

Key words and phrases. Kleinian groups, Schottky groups, two-bridge links, mapping class groups, trace identities, Stern-Brocot tree, Farey sequences, recurrence relations.

the closed Riley slice, which we denote by $\overline{\mathcal{R}}$, is therefore naturally identified with the set of $z \in \mathbb{C}$ such that Γ is discrete and free; in this guise, it forms the region of Section 1.3 lying outside the diamond-shaped unshaded region (along with the boundary of that set). The interior of $\overline{\mathcal{R}}$ is the **Riley slice**, which can also be characterised as the set of $z \in \mathbb{C}$ such that Γ is free, discrete, and the Riemann surface $\Omega(\Gamma)/\Gamma$ ($\Omega(\Gamma)$ being the domain of discontinuity of Γ , as defined in [Mas87, E.2]) has homeomorphism type a 4-times punctured sphere.

A detailed historical account of the Riley slice, together with background information for the non-expert and motivating applications (including a brief outline of earlier work on the classification of two-generated arithmetic groups), can be found in [EMS23b]. Though it was first defined in the mid-20th century, recent work on the Riley slice follows on from a 1994 paper of Linda Keen and Caroline Series [KS94] (with some corrections by Yohei Komori and Caroline Series [KS98]). In this paper, they constructed a foliation of the Riley slice via a two-step process:

- (1) Firstly, a lamination of \mathcal{R} is defined, with leaves indexed by $\mathbb{Q}/2\mathbb{Z}$. These leaves consist of certain branches of preimages of $(-\infty, -2)$ under a family of polynomials $\Phi_{p/q}$ ($p/q \in \mathbb{Q}$) and are called the **rational pleating rays**.
- (2) Then, by a completion construction similar to the completion of \mathbb{Q} via Dedekind cuts, the lamination is extended to a foliation whose leaves are indexed by $\mathbb{R}/2\mathbb{Z}$. The leaves adjoined at this step are called the **irrational pleating rays**.

The polynomials $\Phi_{p/q}$ are constructed as trace polynomials of certain words in the group, which we call **Farey words**; these words enumerate all but one of the non-boundary-parallel closed curves on the sphere, and the distinguished branches on the preimages of $(-\infty, -2)$ correspond to curves in the Riley slice along which the lengths of these curves change in a particularly natural way; in particular, the geometric limit of $\Omega(\Gamma)/\Gamma$ as z travels down the (p/q) -pleating ray towards the preimage of -2 is precisely a pair of three-times punctured spheres, where the additional pair of punctures appears as the loxodromic word $W_{p/q}$ is pinched to a parabolic word with trace -2 . The group corresponding to the geometric limit is known as a **cusp group**, and the set of points corresponding to cusp groups are dense in the boundary of the slice by a result of Curtis McMullen [McM91]. The cusp points were studied in detail by David Wright [Wri05].

Punctured spheres can be viewed as a limiting case of cone-pointed spheres as the cone angles tend to zero. It is therefore natural to study the case of spheres with four cone points, which have corresponding groups (after normalisation)

$$(1.1) \quad \Gamma_z^{a,b} = \left\langle X = \begin{bmatrix} \alpha & 1 \\ 0 & \alpha^{-1} \end{bmatrix}, Y_z = \begin{bmatrix} \beta & 0 \\ z & \beta^{-1} \end{bmatrix} \right\rangle;$$

where $\alpha = \exp(\pi i/a)$ and $\beta = \exp(\pi i/b)$ (so X and Y_z have respective orders a and b) and z is some element of \mathbb{C} such that Γ_z is discrete and isomorphic to $\mathbb{Z}/a\mathbb{Z} * \mathbb{Z}/b\mathbb{Z}$. The deformation space of these groups is called the (a,b) -**Riley slice**, and we denote it by $\mathcal{R}^{a,b}$; in our paper [EMS22], we show that these spaces admit foliations which are defined in a completely analogous way to the Keen-Series foliation of the parabolic Riley slice. In fact, the polynomials which appear in the construction of this foliation are the trace polynomials of the same Farey words as the parabolic case (i.e. the only changes are in the coefficients of the matrices, not in the construction of the word whose trace is taken).

The abstract setting in which we work is a formalised version of this setup. Throughout:

- $z \in \mathbb{C} \setminus \{0\}$ is a complex parameter (which we do not assume lies in a Riley slice unless otherwise stated);

- a and b are real numbers or infinity (we write $a, b \in \hat{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ and use ‘hats’ in this way for all subsets of \mathbb{C} , so for instance $\hat{\mathbb{C}}$ is the Riemann sphere);
- $\alpha = \exp(\pi i/a)$ and $\beta = \exp(\pi i/b)$;
- and Γ_z to be the group generated by the elements X and Y_z defined symbolically in Equation (1.1).

To simplify notation when z is fixed we often drop the subscripts, so $Y = Y_z$ and $\Gamma = \Gamma_z$. We also use the convention $x := X^{-1}$, $y := Y^{-1}$ (and we use similar conventions throughout without comment when writing group words).

1.1. Structure of the paper. The remainder of this introduction describes some of our results and places them in a geometric context; it is not necessary for the reader who is only interested in the abstract combinatorics of the polynomials to read this, and they may skip to the foundational section Section 2 which gives a combinatorial definition of the Farey words using cutting sequences; we take care to avoid requiring any knowledge of geodesic coding. In Section 3 we list some definitions and results about Farey sequences from classical number theory. Our main result (Theorem 4.7) is proved in Section 4; in the following section, Section 5, we use our results to compute the commutators of the Farey words in a couple of different ways. The final part of the paper consists of two sections; in Section 6, we apply the work of Chesebro and his collaborators [Che+20; Che20] to show a formal analogy between the Farey polynomials and the Chebyshev polynomials. Then in Section 7 we discuss some applications to the approximation of irrational pleating rays and cusps, and give some computational results which show that there are interesting connections with dynamical systems and number theory for future work to explore.

Remark. A MathRepo page for this paper may be found at <https://mathrepo.mis.mpg.de/farey/index.html>.

1.2. The recursion formula. Our main result is Theorem 4.7, which gives a straightforward recursion formula for these Farey polynomials that is independent of the geometry and topology from which they naturally arise. These formulas make it possible to effectively compute deep into rank-two Kleinian groups to look at special words (Farey words) conjecturally describing the relators in a non-free group. Similar recursions for the commutators $[X^n, Y] - 2$ and the traces $\text{tr}^2(X^n) - 4$ were found by Alaqaq, Gong, and Martin in [AGM21], and for the analogues of the Farey polynomials for the Maskit slice in [MSW02, pp. 283–285].

As an example, we describe the interesting special case of our formula where $\alpha = \beta = \exp(i\theta)$ (so $a = b = \pi/\theta$). Write $f_{p/q}$ for $\Phi_{p/q}^{\pi/\theta, \pi/\theta}$; then our recursion becomes

$$\begin{aligned} f_{\frac{r_1}{s_1} \oplus \frac{r_2}{s_2}}(z) &= 8 \cos^2(\theta) - p_{\frac{r_1}{s_1}}(z) p_{\frac{r_2}{s_2}}(z) - p_{\frac{r_1}{s_1} \ominus \frac{r_2}{s_2}}(z) \\ &\quad f_{0/1}(z) = 2 - z, \\ &\quad f_{1/1}(z) = z + 2 \cos(2\theta). \end{aligned}$$

Here the connection with the name “Farey” becomes clear: in our recursion we require $\frac{r_1}{s_1}$ and $\frac{r_2}{s_2}$ to be **Farey neighbours**, $r_1 s_2 - r_2 s_1 = \pm 1$ and \oplus and \ominus are **Farey addition** and **subtraction**,

$$\frac{r_1}{s_1} \oplus \frac{r_2}{s_2} = \frac{r_1 + r_2}{s_1 + s_2}, \quad \frac{r_1}{s_1} \ominus \frac{r_2}{s_2} = \frac{|r_1 - r_2|}{|s_1 - s_2|}.$$

In the case $\alpha \neq \beta$ the formula is similar but the constant term is one of two values which depends on the parity of $s_1 + s_2$ and the combinatorial pattern follows that of the Stern-Brocot tree.

One expects that the Chebyshev polynomials should appear, and indeed it is not difficult to see that

$$\Phi_{1/n}^{\infty,\infty}(z) = \frac{2}{z-4} \left(T_n \left(\frac{z}{2} - 1 \right) z - 4 \right)$$

where T_n is the n th Chebyshev polynomial of the first kind, see Section 6. More generally, for other values of θ ,

$$(1.2) \quad \Phi_{1/n}^{\pi/\theta, \pi/\theta}(z) = \frac{1}{z-4} \left(2T_n \left(\frac{z}{2} - 1 \right) (z - 4 \sin^2(\theta)) - 8 \cos^2(\theta) \right)$$

1.3. Applications and connections to geometry and topology. It is the following remarkable connection to three dimensional hyperbolic geometry and potential applications that motivates our study; we give here only a very rough idea here, precise definitions can be found below or in our work [EMS23a]. Fix a, b integers which are at least 2 or infinity, not both 2, and let $\mathcal{R}^{a,b}$ denote the closure of the quasiconformal deformation space of faithful representations (up to conjugacy) of the free product group $\mathbb{Z}/a\mathbb{Z} * \mathbb{Z}/b\mathbb{Z}$ into $\mathrm{PSL}(2, \mathbb{C})$. We have already described a natural embedding of $\mathcal{R}^{a,b}$ in the complex plane, namely it is the set of $z \in \mathbb{C}$ such that the group $\Gamma_z^{a,b}$ is discrete and isomorphic to $\mathbb{Z}/a\mathbb{Z} * \mathbb{Z}/b\mathbb{Z}$. The set $\mathcal{R}^{a,b}$ is open and its complement is connected and bounded (it is contained in the ball of radius 4 about 0). Generalising work of Keen, Komori, and Series [KS94; KS98] and using McMullen's theorem on cusp density [McM91] in [Elz22; EMS23a; EMS23b] we established the following:

Theorem 1.3. *The set $\mathcal{R}^{a,b}$ is the interior of the unbounded component of*

$$\mathbb{C} \setminus \overline{\bigcup_{p/q \in [0,1]} \{z : \Phi_{p/q}^{a,b}(z) = -2\}}$$

The set $\mathcal{R}^{\infty,\infty}$, the classical Riley slice studied by Keen and Series, is illustrated in Section 1.3; a detailed historical account, together with background information for the non-expert and motivating applications (including a brief outline of earlier work on the classification of two-generated arithmetic groups), can be found in [EMS23b].

A theorem of Miyachi and Ohshika [OM10] informs us that $\mathbb{C} \setminus \mathcal{F}_{\infty,\infty}$ is a Jordan domain which is not a quasicircle, so one expects a complicated boundary. A point $z \in \partial \mathcal{R}^{a,b}$ where $\Phi_{p/q}^{a,b}(z) = -2$ is called a **cusp**, and in fact Equation (1.2) can be used to precisely identify the shape of the most prominent cusp of the moduli space $\mathcal{R}^{a,b}$. Miyachi [Miy03] states that there is a “universal shape” to the all cusps in $\mathcal{R}^{\infty,\infty}$ of the form $y^2 = tx^3$, with t depending on the particular cusp, and this persists for $\mathcal{R}^{a,b}$ (work in progress). For example, for the Riley slice at the 0/1-cusp at 4 we have $t = \frac{1}{\pi^2}$.

A point to Theorem 1.3, and evidenced by the above example where we allowed θ to range over values which do not correspond to orbifolds arising in generalised Keen–Series theory, is that the Farey polynomials have allowed us to abstract away the very challenging questions concerning the discreteness of Kleinian groups and replace them with the study of root sets of recursively defined polynomials to obtain information about the geometry of deformation spaces. These formulae may be used to generate high-resolution pictures of the deformation spaces far more quickly than working directly from the definition of cut sequences as it bypasses slow matrix multiplications; a working software implementation of this in `Python` may be found online [Elz21]. Similar recursions for the commutators $[X^n, Y] - 2$ and the traces




FIGURE 1. Cusp groups on the boundary of the Riley slice (coloured and shown with pleating rays. Courtesy of Y. Yamashita). As $n \rightarrow \infty$ the $\frac{1}{n}$ -cusp groups accumulate at the limit group $\frac{0}{1} = 4$ which is the (∞, ∞, ∞) -triangle group. Each of the $\frac{1}{n}$ -cusp groups has limit set a different Apollonian type circle packing. The other solutions to $p_{p/q}(z) = -2$ are shown in red.

$\text{tr}^2(X^n) - 4$ were found by Alaquad, Gong, and Martin in [AGM21], and for the analogues of the Farey polynomials for the Maskit slice in [MSW02, pp. 283–285]. Links between these Farey words and other combinatorial words which occur in Teichmüller theory have been surveyed by Gilman [Gil06].

We wish to make a few more observations about these Farey polynomials and their geometric interpretations. The first is that the Farey polynomials are “trace polynomials”. The simple closed curves on the four times punctured sphere which separate one pair of points from another are enumerated by a rational slope $p/q \in [0, 1]$. Each such slope generates a so-called **Farey word** $W_{p/q}$ in the fundamental group of the four times punctured sphere which represents the p/q simple closed curve. As noted above, this was used by Keen and Series [KS94] in this context and is carefully explained in Section 2 below via cut sequences. Relatedly, these rational slopes and Farey words also enumerate the two-bridge knots and links. Riley [Ril72] proved that the fundamental group of any two-bridge link has a faithful presentation on the two parabolic generators

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \text{ and } \begin{bmatrix} 1 & 0 \\ z & 1 \end{bmatrix},$$

for some $z \in \mathbb{C}$; that is, these fundamental groups are of the form $\Gamma_z^{\infty, \infty}$ for some z . Farey words can be viewed as relators in the fundamental groups of the 3-manifold




FIGURE 2. Two rapidly computed deformation space complements. Left: the quasiconformal deformation space of Kleinian groups generated by two elements of order 3. Right: the quasiconformal deformation space of Kleinian groups generated by two elements of order 4.

knot or link complement $\mathbb{S}^3 \setminus K$ when the generators are chosen as meridians, and in fact this can be taken as another definition of a Farey word [Ril72]. In particular when z is such that $\Gamma_z^{\infty, \infty}$ is a knot group the word $W_{p/q}$ is trivial, hence for the knot or link of slope p/q we have $\Phi_{p/q}^{\infty, \infty}(z) = 2$. One might ask what geometric interpretations exist when the parameters a, b are non-integral.

Far more is true due to the classification theorem [Aki+21] of all discrete groups generated by two parabolics. A corollary to that work is the following.

Theorem 1.4. *If $\Gamma = \Gamma_z^{\infty, \infty}$ is discrete then either Γ is free or there is a Farey word $W_{p/q}$ and integers r and s so that*


$$\Phi_{p/q}^{\infty, \infty}(z_0) = 2 \cos \frac{r\pi}{s} \in [-2, 2].$$

1.4. Conclusion. Taken altogether, these observations concerning the Farey polynomials yield a practical and effective approach to computing in these quasiconformal deformation spaces, see as illustrated in Figure 2. We hope to ultimately describe effective algorithms (perhaps partial algorithms) to solve the following decision problem for rank-two subgroups of $\mathrm{SL}(2, \mathbb{C})$:

Problem 1.5. Given two matrices in $\mathrm{SL}(2, \mathbb{C})$, decide whether the group G they generate is discrete. If G is discrete decide whether it is free or not, and if G is not free effectively produce a relator.

The combination of the current paper with our paper [EMS23a] goes some way towards solving this problem.

1.5. Acknowledgements. We profoundly thank the referee for the numerous good suggestions which have improved this paper. Much of this paper appeared as part of the first author's MSc thesis [Elz22]. We would like to thank the organisers of the Groups and Geometry workshop on Waiheke Island, New Zealand, in December, 2021 for the opportunity to speak, and the attendees of that conference for their interesting questions and helpful comments. The first author also thanks Thu Hien Nguyen for hosting him at the Mathematisches Forschungsinstitut Oberwolfach during June 2022.

FIGURE 3. The cutting sequence of the $1/2$ Farey word.

2. CUTTING SEQUENCES AND FAREY WORDS

We define the Farey word¹ of slope p/q via cutting sequences: this is essentially an interpretation of Dehn's classical algorithm for curves on surfaces [Sti83, Chapter 6] in the language of symbolic dynamics and mapping class groups by Birman and Series [BS87], and for the concrete case of interest to us we follow [Ser85], [Ser91], and [KS94]. Since we want to give a purely abstract version of the theory, we start with an abstract marked tiling of the plane in what follows. Nevertheless in the classical situations where $\Gamma = \langle X, Y \rangle$ is a Schottky-type group uniformising a surface, this covering is topologically equivalent to the lift to the universal cover of the tiling of $\hat{\mathbb{C}}$ induced by the action of Γ . The interested reader should look at the references just listed for details on this interpretation.

Consider the marked tiling of \mathbb{R}^2 shown in Figure 3, and let $L_{p/q}$ be the line through $(0,0)$ of slope p/q ; now define $S_{p/q} = L_{p/q} \cap 2\mathbb{Z}^2$. The **Farey word of slope p/q** , $W_{p/q}$, is the word of length $2q$ such that the k th letter is the label on the right-hand side of the k th vertical line segment crossed by $L_{p/q}$ (i.e. the label to the right of the point $(p/q)i$); if $(p/q)k$ is a lattice point then this definition is ambiguous and by convention we take the label on the north-east side. In other words, the k th letter of $W_{p/q}$ is determined by the parity of $\text{ceil}(p/q)k$ with the convention that $\text{ceil } n = n + 1$ for integral n .

Example 2.1. As an example of the construction process, from Figure 3 we can read off that $W_{1/2} = yxYX$; we include a list of the Farey words with $q \leq 12$ as Table 1.

Remark. Note that the number of words of length at most n in $\langle X, Y \rangle$ grows like $4 \cdot 3^{n-1}$. The word length of $W_{p/q}$ is $2q$. Hardy and Wright prove [HW60,

¹We name these words and the related polynomials after John Farey Sr. as they are closely related to the so-called Farey sequences of rational numbers which we will discuss briefly later in this paper; with regard to this attribution, we quote from the historical notes to Chapter III of Hardy and Wright [HW60, pp. 36–37]: “The history of ‘Farey series’ is very curious... [their properties] seem to have been stated and proved first by Haros in 1802... Farey did not publish anything on the subject until 1816. [...] Mathematicians generally have followed Cauchy’s example in attributing the results to Farey, and the series will no doubt continue to bear his name. Farey has a notice of twenty lines in the *Dictionary of national biography* where he is described as a geologist. As a geologist he is forgotten, and his biographer does not mention the one thing in his life which survives.”

TABLE 1. Farey words $W_{p/q}$ for small q .

p/q	$W_{p/q}$
0/1	yX
1/1	YX
1/2	$yxYX$
1/3	$yXYxYX$
2/3	$yxyXYX$
1/4	$yXyxYxYX$
3/4	$yxyxYXYX$
1/5	$yXyXYxYxYX$
2/5	$yXYxyXyxYX$
3/5	$yxYXYxyXYX$
4/5	$yxyxyXYXYX$
1/6	$yXyXyxYxYxYX$
5/6	$yxyxyxYXYXYX$
1/7	$yXyXyXYxYxYxYX$
2/7	$yXyxYxyXyXYxYX$
3/7	$yXYxyXYxYXYxYX$
4/7	$yXyXYxyXYxyXYX$
5/7	$yxyXYXYxyxYXYX$
6/7	$yxyxyxyXYXYXYX$
1/8	$yXyXyXyxYxYxYxYX$
3/8	$yXYxYXyxYxyXyxYX$
5/8	$yxYXYxyxYXyxyXYX$
7/8	$yxyxyxyxYXYXYXYX$
1/9	$yXyXyXyXYxYxYxYxYX$
2/9	$yXyXYxYxyXyxYxYxYX$
4/9	$yXYxyXYxyXyxYxYxYX$
5/9	$yxYXyxYXYxyXYxyXYX$
7/9	$yxyxYXYXYxyxyXYXYX$
8/9	$yxyxyxyxyXYXYXYXYX$
1/10	$yXyXyXyXyxYxYxYxYxYX$
3/10	$yXyxYxyXyxYxYxYxYxYX$
7/10	$yxyXYXyxyxYXYxyxYXYX$
9/10	$yxyxyxyxyxYXYXYXYXYX$
1/11	$yXyXyXyXyXYxYxYxYxYxYX$
2/11	$yXyXyxYxYxyXyXYxYxYxYX$
3/11	$yXyxYxYxyXYxYxyXyXYxYX$
4/11	$yXYxYXyXYxyXyxYxyXyxYX$
5/11	$yXYxyXYxyXYxYXyxYXyxYX$
6/11	$yxYXyxYXyxyXYxyXYxyXYX$
7/11	$yXYYxyxyXYxyXYxyXYXYX$
8/11	$yxyXYXYxyxyXYXyxYxyXYX$
9/11	$yxyxyXYXYXYxyxyXYXYXYX$
10/11	$yxyxyxyxyxyXYXYXYXYXYX$
1/12	$yXyXyXyXyXyxYxYxYxYxYX$
5/12	$yXYxyXyxYXyxYxyXYxYxYxYX$
7/12	$yXYYxyxyXYxyxYXyxYXYxyXYX$
11/12	$yxyxyxyxyxyxYXYXYXYXYXYX$

Theorem 330 p.268] that the length of the Farey sequence of order $2q$, that is the sequence of all reduced fractions between 0 and 1 with denominator at most $2q$, is asymptotic to $\frac{3}{\pi^2}(2q)^2$. Thus, assuming the conjectures and given $\langle X, Y \rangle$ is discrete but not free, we have a very powerful tool for finding relators of length q or less. We need only examine the approximately q^2 values $\Phi_{p/q}^{\infty,\infty}(z_0)$ of the (roughly) $4 \cdot 3^{2q-1}$ possibilities for the relator. Note that it is not claimed this is a complete presentation (and sometimes it is not) but once we know $W_{p/q}$ is of finite order then this is often enough to identify the group as a Dehn surgery on a knot or link complement. This has proved a useful tool in the enumeration of arithmetic groups of $\mathrm{PSL}(2, \mathbb{C})$ generated by two elements. Conjecturally there are only finitely many such groups, and it is known that there are only finitely many such groups of $\mathrm{PSL}(2, \mathbb{C})$ generated by two parabolic or elliptic generators [MM99]. For instance the groups generated by two parabolic elements are known, and there are only four of them [GMM98]. All of these groups are knot or link groups and lie in the exterior of the Riley slice.

There are various symmetries visible in the Farey words; for instance, they are alternating products of $X^{\pm 1}$ and $Y^{\pm 1}$ which always end in X (this is obvious from the definition). The following slightly less trivial symmetry will be useful in the sequel:

Lemma 2.2. *Let $W_{p/q}$ be a Farey word; then the word consisting of the first $2q-1$ letters of $W_{p/q}$ is conjugate to X or Y according to whether the q th letter of $W_{p/q}$ is $X^{\pm 1}$ or $Y^{\pm 1}$ (i.e. according to whether q is even or odd respectively).*

Proof. This identity comes from considering the rotational symmetry of the line of slope p/q about the point (q, p) ; it is clear from the symmetry of the picture that the first $p-1$ letters of $W_{p/q}$ are obtained from the $(p+1)$ th to $(2p-1)$ th letters by reversing the order and swapping the case (imagine rotating the line by 180 degrees onto itself and observe the motion of the labelling). \square

Remark. For a more precise statement see e.g. the first lemma of Section 2 of our sister paper [EMS23a].

We do not use the following lemma in this paper, but we state it for interest's sake while we are studying the symmetries of the Farey words.

Lemma 2.3. *Let p/q be a fraction in least terms; then $W_{-p/q}$ is obtained from $W_{p/q}$ by the following process: swap the sign of the exponent of every letter except for the q th and $2q$ th letters.*

Proof. For all i such that p/qi is non-integral, we have $\text{ceil} -\frac{p}{q}i = -\text{floor} \frac{p}{q}i = 1 - \text{ceil} \frac{p}{q}i$. In particular, the parity of the ceiling is swapped and so the sign of the exponent is swapped. For i such that p/q is integral (i.e. i is q or $2q$), due to our convention on the ceiling function, we have $\text{ceil} -\frac{p}{q}i = 1 - \frac{p}{q}i$ and $\text{ceil} \frac{p}{q}i = \frac{p}{q}i + 1$, so the parities of $\text{ceil} -\frac{p}{q}i$ and $\text{ceil} \frac{p}{q}i$ are equal and the exponent of the corresponding letter is unchanged. \square

The **Farey polynomial of slope p/q** is defined by $\Phi_{p/q} := \text{tr } W_{p/q}$; this is a polynomial in z of degree q , with coefficients rational functions of α and β . If we wish to emphasise the dependence on a and b , we write $\Phi_{p/q}^{a,b}$.

Remark. The Farey polynomials are not to be confused with the so-called **Riley polynomials** $\Lambda_{p/q}$ defined by Riley [Ril72] and studied e.g. by Chesebro [Che20]; these $\Lambda_{p/q}$ is the polynomial in z which is the top-left entry of the word consisting of the first $q-1$ letters of $W_{p/q}$ (i.e. the conjugating word in Lemma 2.2). See also Remark 5.3.13 of [Aki+07].

TABLE 2. Farey polynomials $\Phi_{p/q}^{a,b}(z)$ for small q .

p/q	
0/1	$\frac{\alpha}{\beta} + \frac{\beta}{\alpha} - z$
1/1	$\alpha\beta + \frac{1}{\alpha\beta} + z$
1/2	$2 + (\alpha\beta - \frac{\alpha}{\beta} - \frac{\beta}{\alpha} + \frac{1}{\alpha\beta})z + z^2$
1/3	$\frac{1}{\alpha\beta} + \alpha\beta + \left(3 - \frac{1}{\alpha^2} - \alpha^2 - \frac{1}{\beta^2} - \beta^2 + \frac{\alpha^2}{\beta^2} + \frac{\beta^2}{\alpha^2}\right)z$ $+ \left(\alpha\beta - 2\frac{\alpha}{\beta} - 2\frac{\beta}{\alpha} + \frac{1}{\alpha\beta}\right)z^2 + z^3$
2/3	$\frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \left(-3 + \alpha^2 + \frac{1}{\alpha^2} - \frac{1}{\alpha^2\beta^2} - \alpha^2\beta^2 + \beta^2 + \frac{1}{\beta^2}\right)z$ $+ \left(-2\alpha\beta - \frac{2}{\alpha\beta} + \frac{\alpha}{\beta} + \frac{\beta}{\alpha}\right)z^2 - z^3$
1/4	$2 + \left(\frac{\alpha}{\beta^3} - \frac{\alpha^3}{\beta^3} + \frac{2}{\alpha\beta} - 3\frac{\alpha}{\beta} + \frac{\alpha^3}{\beta} + \frac{\beta}{\alpha^3} - 3\frac{\beta}{\alpha} + 2\alpha\beta - \frac{\beta^3}{\alpha^3} + \frac{\beta^3}{\alpha}\right)z$ $+ \left(6 - \frac{2}{\alpha^2} - 2\alpha^2 - \frac{2}{\beta^2} + 3\frac{\alpha^2}{\beta^2} - 2\beta^2 + 3\frac{\beta^2}{\alpha^2}\right)z^2$ $+ \left(\frac{1}{\alpha\beta} - 3\frac{\alpha}{\beta} - 3\frac{\beta}{\alpha} + \alpha\beta\right)z^3 + z^4$
3/4	$2 + \left(\frac{1}{\alpha^3\beta^3} - \frac{1}{\alpha\beta^3} - \frac{1}{\alpha^3\beta} + \frac{3}{\alpha\beta} - 2\frac{\alpha}{\beta} - 2\frac{\beta}{\alpha} + 3\alpha\beta - \alpha^3\beta - \alpha\beta^3 + \alpha^3\beta^3\right)z$ $+ \left(6 - \frac{2}{\alpha^2} - 2\alpha^2 - \frac{2}{\beta^2} + \frac{3}{\alpha^2\beta^2} - 2\beta^2 + 3\alpha^2\beta^2\right)z^2$ $+ \left(\frac{3}{\alpha\beta} - \frac{\alpha}{\beta} - \frac{\beta}{\alpha} + 3\alpha\beta\right)z^3 + z^4$

Example 2.4. We list the Farey polynomials $\Phi_{p/q}^{a,b}$ with $q \leq 4$ in Table 2. This illustrates some of the difficulty in studying these polynomials: they are clearly very symmetric, but quickly become too unwieldy to write explicitly and so actually guessing what the symmetries *are* in general is hard. We also list the first few ‘Fibonacci’ Farey polynomials $\Phi_{\text{fib}(q-1)/\text{fib}(q)}^{\infty,\infty}$ (as usual, $\text{fib}(1) = 1$, $\text{fib}(2) = 1$, and $\text{fib}(n) := \text{fib}(n-1) + \text{fib}(n-2)$) in Table 3.

3. FAREY THEORY

Our recursion formula, like that for the Maskit slice in [MSW02, pp. 283–285], is a recursion down the **Farey diagram**. We recall some of the notation and ideas, which can be found for instance in [GKP94, §4.5] or [HW60, Chapter III].

There is a subgroup Γ_1 of $\text{PSL}(2, \mathbb{Z})$ generated by the two matrices

$$L = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \quad \text{and} \quad R = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

which acts via componentwise fractional linear transformations on the set of ordered pairs $(p/q, r/s)$ with $p/q < r/s$ and $|ps - rq| = 1$ (the action being essentially the faithful action of Γ_1 on itself by left multiplication). These matrices appear naturally in relation to the geometry and topology as they generate the orientation-preserving half of the mapping class group of the four-times punctured sphere and hence are the natural group which acts on the complex of geodesics (for more information on this perspective see the discussion in [Aki+07] and [BZ03, Chapter 12]).

TABLE 3. Farey polynomials of the form $\Phi_{\text{fib}(q-1)/\text{fib}(q)}^{\infty, \infty}(z)$ for small q .

$\frac{\text{fib}(q-1)}{\text{fib}(q)}$	
0/1	$2 - z$
1/1	$2 + z$
1/2	$2 + z^2$
2/3	$2 - z - 2z^2 - z^3$
3/5	$2 + z + 2z^2 + 3z^3 + 2z^4 + z^5$
5/8	$2 + 4z^4 + 8z^5 + 8z^6 + 4z^7 + z^8$
8/13	$2 - z - 2z^2 - 5z^3 - 12z^4 - 22z^5 - 32z^6 - 44z^7 - 54z^8$ $- 53z^9 - 38z^{10} - 19z^{11} - 6z^{12} - z^{13}$
13/21	$2 + z + 2z^2 + 7z^3 + 14z^4 + 31z^5 + 64z^6 + 124z^7 + 214z^8$ $+ 339z^9 + 498z^{10} + 699z^{11} + 936z^{12} + 1148z^{13} + 1216z^{14}$ $+ 1064z^{15} + 746z^{16} + 409z^{17} + 170z^{18} + 51z^{19} + 10z^{20} + z^{21}$
21/34	$2 + z^2 + 8z^4 + 24z^5 + 68z^6 + 192z^7 + 516z^8 + 1256z^9 + 2834z^{10}$ $+ 5912z^{11} + 11460z^{12} + 20816z^{13} + 35598z^{14} + 57248z^{15}$ $+ 86446z^{16} + 122560z^{17} + 163199z^{18} + 203952z^{19} + 238564z^{20}$ $+ 259704z^{21} + 260686z^{22} + 238320z^{23} + 195694z^{24} + 142328z^{25}$ $+ 90451z^{26} + 49552z^{27} + 23058z^{28} + 8952z^{29} + 2831z^{30} + 704z^{31}$ $+ 130z^{32} + 16z^{33} + z^{34}$
34/55	$2 - z - 4z^2 - 10z^3 - 34z^4 - 103z^5 - 286z^6 - 791z^7$ $- 2078z^8 - 5221z^9 - 12680z^{10} - 29824z^{11} - 67872z^{12}$ $- 149896z^{13} - 321800z^{14} - 671896z^{15} - 1364228z^{16}$ $- 2692102z^{17} - 5158232z^{18} - 9587668z^{19} - 17273444z^{20}$ $- 30141702z^{21} - 50903644z^{22} - 83138942z^{23} - 131230688z^{24}$ $- 200056876z^{25} - 294348624z^{26} - 417663240z^{27} - 571010576z^{28}$ $- 751328456z^{29} - 950188464z^{30} - 1153232920z^{31} - 1340813030z^{32}$ $- 1490107333z^{33} - 1578696308z^{34} - 1589182962z^{35} - 1513960786z^{36}$ $- 1358696535z^{37} - 1142850158z^{38} - 896137319z^{39} - 651440922z^{40}$ $- 436582355z^{41} - 268228504z^{42} - 150207744z^{43} - 76207672z^{44}$ $- 34797892z^{45} - 14193584z^{46} - 5125756z^{47} - 1621110z^{48}$ $- 442809z^{49} - 102556z^{50} - 19630z^{51} - 2990z^{52}$ $- 341z^{53} - 26z^{54} - z^{55}$

Definition 3.1. If $p/q, r/s \in \hat{\mathbb{Q}}$ satisfy the determinant condition $|ps - rq| = 1$ then we say that p/q and r/s are **Farey neighbours**.

The primary elementary lemma on Farey neighbourliness is the following, which appears in [HW60, §3.3]:

Lemma 3.2. If p/q and r/s are Farey neighbours with $p/q < r/s$, then $p/q < (p+r)/(q+s) < r/s$, and $(p+r)/(q+s)$ is the unique fraction of minimal denominator between p/q and r/s . More precisely, let u/v be any fraction in $(p/q, r/s)$; then there exist two positive integers λ, μ such that

$$u = \lambda p + \mu r \text{ and } v = \lambda r + \mu s$$

(so of course the minimal denominator is obtained when $\lambda = \mu = 1$). \square

The operation

$$(p/q, r/s) \mapsto (p+r)/(q+s)$$




FIGURE 4. The subset of the Farey triangulation of \mathbb{H}^2 with vertices $p/q \in [0, 1] \cap \mathbb{Q}$ such that $q \leq 10$.

will be fundamental to our later study; it is called the **mediant** or **Farey addition** operation. We write $(p/q) \oplus (r/s)$ for the Farey addition of p/q to r/s . We will be careful to only combine p/q and r/s in this way if they are Farey neighbours.

We now check that the usual action of Γ_1 on $\hat{\mathbb{Q}}$ as a group of fractional linear transformations agrees with the action of Γ_1 on the space of homology bases, in the following way. Suppose that r_1/s_1 and r_2/s_2 are Farey neighbours with $p/q = r_1/s_1 \oplus r_2/s_2$, and let $A \in \Gamma_1$ be represented by the matrix

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix};$$

A acts on the pair $\{r_1/s_1, r_2/s_2\}$ by matrix multiplication,

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \left\{ \frac{r_1}{s_1}, \frac{r_2}{s_2} \right\} = \left\{ \frac{ar_1 + bs_1}{cr_1 + ds_1}, \frac{ar_2 + bs_2}{cr_2 + ds_2} \right\},$$

a simple computation shows that the Farey sum of the images is exactly

$$\frac{a(r_1 + r_2) + b(s_1 + s_2)}{c(r_1 + r_2) + d(s_1 + s_2)} = \frac{ap + bq}{cp + dq},$$

i.e. the image of p/q under A with respect to the usual action of $\text{PSL}(2, \mathbb{R})$ on the upper half plane as a group of fractional linear transformations.

The ideal triangle Δ in the upper half plane \mathbb{H}^2 spanned by

$$(1/0, 1/1, 0/1) = (\infty, 1, 0)$$

does not bound a fundamental domain for this action, since $X^{-1}(D) = Y^{-1}(D)$; however, the images of Δ under $\text{PSL}(2, \mathbb{Z})$ do tessellate \mathbb{H}^2 in the sense that if $W\Delta \cap W'\Delta \neq \emptyset$ then either the intersection is a shared edge or the two images are equal. The simplicial complex thus formed is called the **Farey triangulation** of \mathbb{H}^2 ; we denote it by \mathcal{D} . The portion of \mathcal{D} ‘above’ the segment $[0, 1]$ is depicted in Figure 4. The vertices of \mathcal{D} are exactly the points of $\hat{\mathbb{Q}}$, and the 2-faces are triangles with vertex triples of the form $\{p/q, (p+r)/(q+s), r/s\}$ where $ps - qr = \pm 1$. We denote the set of 2-faces by $\mathcal{D}(2)$. This second assertion follows immediately from the definition of \mathcal{D} as a tessellation: if $\{a/b, c/d, e/f\} \in \mathcal{D}(2)$, then there exists some matrix

$$\begin{bmatrix} p & r \\ q & s \end{bmatrix} \in \text{PSL}(2, \mathbb{Z})$$




FIGURE 5. The Farey addition graph.

which acts on the original triangle as


$$\begin{bmatrix} p & r \\ q & s \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} a & c & e \\ b & d & f \end{bmatrix},$$

giving immediately that $(a/b, c/d, e/f) = (p/q, (p+r)/(q+s), r/s)$; i.e. the triangles of \mathcal{D} are exactly the triples $(\alpha, \alpha \oplus \beta, \beta)$ for Farey neighbours $\alpha, \beta \in \hat{\mathbb{Q}}$. The **Farey graph** is the digraph with vertices $\hat{\mathbb{Q}}$ and directed edges from p/q and r/s to $p/q \oplus r/s$ (Figure 5). This is just the 1-skeleton of \mathcal{D} with an added orientation.

It will be convenient, finally, to have the notation $p/q \ominus r/s$ for the fraction $(p-r)/(q-s)$; we shall only use this when it is known that $(p-r)/(q-s)$ and r/s are Farey neighbours (which is implied by neighbourliness of p/q and r/s). If $(p/q, r/s)$ is an edge in the Farey triangulation, then it forms the boundary between the two triangles $(p/q, p/q \oplus r/s, r/s)$ and $(p/q, p/q \ominus r/s, r/s)$.

4. A RECURSION FORMULA TO GENERATE FAREY POLYNOMIALS

In this section, we will give a recursion formula for the Farey polynomials. This recursion will be a recursion ‘down the Farey graph’, in the sense that its input will be the Farey polynomials at the vertices of a triangle $(p/q, p/q \ominus r/s, r/s)$ and its output will be the Farey polynomial at $p/q \oplus r/s$:



We begin by finding a similar recurrence for the Farey words; we will then produce a recurrence for the polynomials using standard trace identities and elbow grease. One might guess, for instance by analogy with the Maskit slice [MSW02, p. 277], that $W_{p/q} W_{r/s} = W_{p/q \oplus r/s}$. It is easy to check whether or not this is true:

Example 4.1.

- $W_{1/2} = yxYX, W_{1/1} = YX,$

$$W_{1/2}W_{1/1} = yx\textcolor{red}{Y}XYX, \text{ and}$$

$$W_{1/2\oplus 1/1} = W_{2/3} = yx\textcolor{red}{y}XYX.$$

- $W_{1/3} = yXYxYX, W_{2/5} = yXYxyXyxYX,$

$$W_{1/3}W_{2/5} = yXYxYXy\textcolor{red}{X}YxyXyxYX, \text{ and}$$

$$W_{1/3\oplus 2/5} = W_{3/8} = yXYxYXy\textcolor{red}{x}YxyXyxYX.$$

Example 4.1 shows that our guess is almost correct; the corrected statement is:

Lemma 4.2. *Let p/q and r/s be Farey neighbours with $p/q < r/s$. Then $W_{p/q\oplus r/s}$ is equal to:*

- (1) $W_{p/q}W_{r/s}$ with the sign of the $(q+s)$ th exponent swapped;
- (2) $W_{r/s}W_{p/q}$ with the sign of the $(2s)$ th exponent swapped.

Proof. We prove (1); the proof of (2) is done in the same way, and we briefly indicate the difference at the end. The situation is diagrammed in Figure 6 for convenience. To simplify notation, in this proof we write $h(i)$ for the height $(\frac{p}{q} \oplus \frac{r}{s})i$. Observe that $h(i)$ is integral only at $i = 0$ and $i = 2q + 2s$ (at both positions trivially the letters in $W_{(p/q)\oplus(r/s)}$ and $W_{p/q}W_{r/s}$ are identical) and at $i = q + s$. The lemma will follow once we check that at the positions $i \notin \{0, 2q + 2s\}$,

$$(4.3) \quad \text{if } 0 < i \leq 2q \text{ then } (p/q)i < h(i) < \text{ceil}(p/q)i$$

and

$$(4.4) \quad \text{if } 0 < i < 2s \text{ then } (r/s)i + 2q < h(i + 2p) < \text{ceil}[(r/s)i + 2p] :$$

indeed, these inequalities show that at every integral horizontal distance the height of the line corresponding to $W_{p/q\oplus r/s}$ is meeting the same vertical line segment as the line corresponding to $W_{p/q}$ or $W_{r/s}$, and so the letter chosen is the same except at $i = q + s$ since at this position the height of the line of slope $(p/q) \oplus (r/s)$, being integral, is rounded up to $h(i) + 1$ while the height of the line of slope r/s is non-integral so is rounded up to the integer $h(i)$.

Observe now that the inequalities Equations (4.3) and (4.4) are equivalent to the following: there is no integer between $(p/q)i$ and $(p/q \oplus r/s)i$ (exclusive) if $0 < i \leq 2q$, and there is no integer between $(r/s)i + 2q$ and $h(i + 2p)$ if $0 < i < 2s$. But these follow from Lemma 3.2. Indeed, the lemma shows that no integer lies between p/q and $(p/q) \oplus (r/s)$; suppose a/b is a rational between $(p/q)i$ and $h(i)$, then $a = i(\lambda p + \mu(p+r))$ and $b = (\lambda q + \mu(q+s))$ for some positive λ, μ ; suppose $a/b \in \mathbb{Z}$, so $\lambda q + \mu(q+s)$ divides $i(\lambda p + \mu(p+r))$. By the case $i = 1$, $\lambda p + \mu(p+r)$ and $\lambda q + \mu(q+s)$ are coprime, so $\lambda q + \mu(q+s)$ divides i ; but $\lambda q + \mu(q+s) \geq 2q$. The case of $(r/s)i + 2q$ and $h(i + 2p)$ is proved in a similar way. (The proof of part (2) differs only in the relative position of the ‘kink’ between the concatenated p/q and r/s segments and the $(p+r)/(q+s)$ segment: the kink is at position $2s$ horizontally and lies above the $(p+r)/(q+s)$ line rather than below, so it is at the $(2s)$ th position that the ‘modified ceiling’ makes a nontrivial appearance.) \square

Lemma 4.2 is not new; for instance, part (1) appears as Proposition 4.2.4 of [Zha10], and part (2) was known to Keen and Series at least (though as far as we know they did not publish it, see [KS94, Remark at end of §4]). However, the following consequences for the Farey polynomials are comparatively nontrivial to prove and have, to the best of our knowledge, not appeared before in the literature.




FIGURE 6. Farey addition versus word multiplication for $W_{p/q}$ (red) and $W_{r/s}$ (green).

Lemma 4.5 (Product lemma). *Let p/q and r/s be Farey neighbours with $p/q < r/s$. Then the following trace identity holds:*

$$\mathrm{tr} W_{p/q} W_{r/s} + \mathrm{tr} W_{p/q \oplus r/s} = \begin{cases} 2 + \alpha^2 + \frac{1}{\alpha^2} & \text{if } q+s \text{ is even,} \\ \alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} & \text{if } q+s \text{ is odd.} \end{cases}$$

Proof. Trace is invariant under cyclic permutations, thus (applying Lemma 4.2) we can write

$$\mathrm{tr} W_{p/q} W_{r/s} = \mathrm{tr} AB \text{ and } \mathrm{tr} W_{p/q \oplus r/s} = \mathrm{tr} AB^{-1},$$

where B is the $(q+s)$ th letter of $W_{p/q} W_{r/s}$ and A is the remainder of the word but with the final letters cycled to the front. Now we know that $\mathrm{tr} AB = \mathrm{tr} A \mathrm{tr} B - \mathrm{tr} AB^{-1}$ (see the useful list of trace identities found in Section 3.4 of [MR03]), so it suffices to check that $\mathrm{tr} A \mathrm{tr} B = 2 + \alpha^2 + \frac{1}{\alpha^2}$ if $q+s$ is even and $\alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta}$ otherwise.

Case I. $q+s$ is even. Observe now that B is $X^{\pm 1}$ if $q+s$ is even; then $\mathrm{tr} B = 2\Re\alpha$. The identity to show is therefore $\mathrm{tr} A = (2 + \alpha^2 + \frac{1}{\alpha^2})/(2\Re\alpha)$; recalling that $|\alpha| = 1$ and using the double angle formulae we have

$$\frac{2}{2\Re\alpha} + \frac{\alpha^2}{2\Re\alpha} + \frac{\alpha^{-2}}{2\Re\alpha} = \frac{1}{\cos\theta} + \left(\alpha - \frac{1}{2\cos\theta}\right) + \left(\bar{\alpha} - \frac{1}{2\cos\theta}\right) = 2\cos\theta$$

where $\theta = \mathrm{Arg}\alpha$. Thus we actually just need to show $\mathrm{tr} A = 2\cos\theta$. As an aside, this shows that A is parabolic if X is, and is elliptic if X is.

Case II. $q+s$ is odd. In this case, B is $Y^{\pm 1}$ and so $\mathrm{tr} B = 2\Re\beta$; we therefore wish to show that $\mathrm{tr} A = (\alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta})/(2\Re\beta)$; again using trigonometry we may simplify the right side,

$$\frac{\alpha\beta}{2\Re\alpha} + \frac{\alpha/\beta}{2\Re\alpha} + \frac{\beta/\alpha}{2\Re\alpha} + \frac{1/(\alpha\beta)}{2\Re\beta} = 2\cos\theta$$

and so again we need only show that $\mathrm{tr} A = 2\cos\theta$ where $\theta = \mathrm{Arg}\alpha$.

Both cases then reduce to the identity $\mathrm{tr} A = \mathrm{tr} X$. It will be enough to show that A is conjugate to X ; by construction of A , this is equivalent to showing that in $W_{p/q \oplus r/s}$ the $(q+s+1)$ th to $(2q+2s-1)$ th letters are obtained from the first $q+s-1$ letters by reversing the order and swapping the case. But this is just Lemma 2.2. \square

In the case that X and Y are parabolics and $\alpha = \beta = 1$, the two formulae unify to become:

$$\mathrm{tr} W_{p/q} W_{r/s} = 4 - \mathrm{tr} W_{p/q \oplus r/s}.$$

We may similarly prove the following ‘quotient lemma’:

Lemma 4.6 (Quotient lemma). *Let p/q and r/s be Farey neighbours with $p/q < r/s$. Then the following trace identity holds:*

$$\mathrm{tr} W_{p/q} W_{r/s}^{-1} + \mathrm{tr} W_{p/q \ominus r/s} = \begin{cases} 2 + \beta^2 + \frac{1}{\beta^2} & \text{if } q - s \text{ is even,} \\ \alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} & \text{if } q - s \text{ is odd.} \end{cases}$$

Proof. We begin by setting up notation. By Lemma 2.2 we may write $W_{p/q} = UAuX$ with $A = X^{\pm 1}$ if q is even and $A = Y^{\pm 1}$ if q is odd; similarly, write $W_{r/s} = VBvX$ with B one of $X^{\pm 1}$ or $Y^{\pm 1}$. Then

$$W_{p/q} W_{r/s}^{-1} = UAuXvBv = UAuVbv;$$

by Lemma 4.2, we have also that $W_{r/s} W_{p/q \ominus r/s}$ is $W_{p/q}$ with the sign of the exponent of the q th letter swapped; explicitly,

$$W_{p/q \ominus r/s} VBvX = UauX \implies W_{p/q \ominus r/s} = UauXxVbv = UauVbv.$$

Our goal is therefore to compute $\mathrm{tr} UAuVbv + \mathrm{tr} UauVbv$; performing a cyclic permutation again, this is equivalent to $\mathrm{tr} A(uVbvU) + \mathrm{tr} a(uVbvU)$. In this form, this becomes

$$\mathrm{tr} A(uVbvU) + \mathrm{tr} a(uVbvU) = \mathrm{tr} A \mathrm{tr} uVbvU = \mathrm{tr} A \mathrm{tr} b.$$

Consider now the cases for the product $\mathrm{tr} A \mathrm{tr} b$:

	q odd	q even
s odd	$\mathrm{tr}^2 Y$	$\mathrm{tr} X \mathrm{tr} Y$
s even	$\mathrm{tr} X \mathrm{tr} Y$	$\mathrm{tr}^2 X$

If $p/q, r/s$ are Farey neighbours then it is not possible for both q and s to be even since $ps - rq \equiv 1 \pmod{2}$. Further, $q - s$ is odd iff exactly one of p and q is odd, otherwise $q - s$ is even. Thus we see that if $q - s$ is even then

$$\mathrm{tr} W_{p/q} W_{r/s}^{-1} + \mathrm{tr} W_{p/q \ominus r/s} = \mathrm{tr}^2 Y = (\beta + 1/\beta)^2$$

and if $q - s$ is odd then

$$\mathrm{tr} W_{p/q} W_{r/s}^{-1} + \mathrm{tr} W_{p/q \ominus r/s} = \mathrm{tr} X \mathrm{tr} Y = (\alpha + 1/\alpha)(\beta + 1/\beta)$$

which are the claimed formulae. \square

Using the product and quotient lemmata, we can prove the desired recursion formula for the trace polynomials.

Theorem 4.7 (Recursion formulae). *Let p/q and r/s be Farey neighbours. If $q + s$ is even, then*

$$(4.8) \quad \Phi_{p/q} \Phi_{r/s} + \Phi_{p/q \oplus r/s} + \Phi_{p/q \ominus r/s} = 4 + \frac{1}{\alpha^2} + \alpha^2 + \frac{1}{\beta^2} + \beta^2.$$

Otherwise if $q + s$ is odd, then

$$(4.9) \quad \Phi_{p/q} \Phi_{r/s} + \Phi_{p/q \oplus r/s} + \Phi_{p/q \ominus r/s} = 2 \left(\alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} \right).$$

Proof. Suppose $q + s$ is even; then $q - s$ is also even, so

$$\begin{aligned} \Phi_{p/q} \Phi_{r/s} + \Phi_{(p+r)/(q+s)} + \Phi_{(p-r)/(q-s)} \\ = \mathrm{tr} W_{p/q} \mathrm{tr} W_{r/s} + \mathrm{tr} W_{p/q \oplus r/s} + \mathrm{tr} W_{p/q \ominus r/s} \\ = \mathrm{tr} W_{p/q} W_{r/s} + \mathrm{tr} W_{p/q} W_{r/s}^{-1} + \mathrm{tr} W_{p/q \oplus r/s} + \mathrm{tr} W_{p/q \ominus r/s} \\ = 2 + \alpha^2 + \frac{1}{\alpha^2} + 2 + \beta^2 + \frac{1}{\beta^2} \end{aligned}$$




FIGURE 7. The induced colouring of the Stern-Brocot tree.

where in the final step we used Lemma 4.5 and Lemma 4.6. Similarly, when $q - s$ is odd then $q + s$ is also odd and

$$\begin{aligned}
 & \Phi_{p/q} \Phi_{r/s} + \Phi_{(p+r)/(q+s)} + \Phi_{(p-r)/(q-s)} \\
 &= \text{tr } W_{p/q} \text{tr } W_{r/s} + \text{tr } W_{p/q \oplus r/s} + \text{tr } W_{p/q \ominus r/s} \\
 &= \text{tr } W_{p/q} W_{r/s} + \text{tr } W_{p/q} W_{r/s}^{-1} + \text{tr } W_{p/q \oplus r/s} + \text{tr } W_{p/q \ominus r/s} \\
 &= \alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} + \alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta}
 \end{aligned}$$

as desired. \square

Again in the parabolic case the two formulae unify and the recursion identity becomes


$$(4.10) \quad \Phi_{p/q} \Phi_{r/s} + \Phi_{(p+r)/(q+s)} + \Phi_{(p-r)/(q-s)} = 8;$$

in fact, such a unification occurs whenever $\alpha = \beta$. In the parabolic case this formula has been known to experts for a while but proofs of it go via either passing to a common cover of the four-punctured sphere and the punctured torus and using the analogous identity for the latter [Aki+07; Bow98], or by using directly some version of Bowditch's theory on Markoff triples [MPT15], which we discuss further in the next section. A combinatorial group theory approach as described above which allows the definition of the family of polynomials in a non-geometric abstract context does not appear in the literature.

As an aside, Figure 7 shows the picture we get when we just draw the edges of the Farey graph corresponding to Farey neighbours which appear as products in some iterate of the recursion: it is a nice colouring of the Stern-Brocot tree.²

Observe that $0/1$ and $1/0$ are Farey neighbours in $\hat{\mathbb{Q}}$. Thus, applying Equation (4.8) *formally* to the diamond

²There is a standard bijection in computer science between finite binary strings and vertices of the infinite binary tree; the Stern-Brocot tree is the tree structure on $I = (0, 1] \cap \mathbb{Q}$ obtained by writing every $x \in I$ as the image of $1/1$ under a sequence of R 's and L 's (here R and L are the generators of Γ_1 , the orientation-preserving part of $\text{PSL}(2, \mathbb{Z})$). For a more explicit discussion, see [GKP94, Section 6.7].



we obtain

$$\Phi_{0/1}\Phi_{1/1} + \Phi_{1/0} + \Phi_{1/2} = 4 + \frac{1}{\alpha^2} + \alpha^2 + \frac{1}{\beta^2} + \beta^2;$$

substituting for $\Phi_{1/1}$, $\Phi_{1/2}$, and $\Phi_{0/1}$ from Table 2 we get the following formal expression for $\Phi_{1/0}$:

$$\begin{aligned} \Phi_{1/0} &= 4 + \frac{1}{\alpha^2} + \alpha^2 + \frac{1}{\beta^2} + \beta^2 - \left(\frac{\alpha}{\beta} + \frac{\beta}{\alpha} - z \right) \left(\alpha\beta + \frac{1}{\alpha\beta} + z \right) - 2 \\ &\quad - \left(\alpha\beta - \frac{\alpha}{\beta} - \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} \right) z - z^2 \\ &= 2. \end{aligned}$$

Observe that $\Phi_{1/0}^{-1}((-\infty, -2]) = \emptyset$, so this is compatible with the Keen–Series theory; it is also a polynomial of degree q (here, $q = 0$) with constant term 2, which all agrees with the properties of the higher-degree polynomials. On the other hand, it is not monic!

5. COMMUTATORS OF FAREY WORDS

Recall **Fricke’s identity** for $A, B \in \mathrm{PSL}(2, \mathbb{C})$ [MR03, (3.15)]:

$$\mathrm{tr}[A, B] = \mathrm{tr}^2 A + \mathrm{tr}^2 B + \mathrm{tr}^2 AB - \mathrm{tr} A \mathrm{tr} B \mathrm{tr} AB - 2.$$

As a direct consequence of Lemma 4.5, we obtain the following special case for Farey words:

Lemma 5.1. *If p/q and r/s are Farey neighbours with $p/q < r/s$, then*

$$\begin{aligned} \mathrm{tr}[W_{p/q}, W_{r/s}] &= \Phi_{p/q}^2 + \Phi_{r/s}^2 + \Phi_{(p/q)\oplus(r/s)}^2 + \Phi_{p/q}\Phi_{r/s}\Phi_{(p/q)\oplus(r/s)} \\ &\quad + \kappa(p/q, r/s)^2 - \kappa(p/q, r/s) (\Phi_{p/q}\Phi_{r/s} + 2\Phi_{(p/q)\oplus(r/s)}) - 2; \end{aligned}$$

where

$$\kappa(p/q, r/s) := \begin{cases} 2 + \alpha^2 + \frac{1}{\alpha^2} & \text{if } q+s \text{ is even,} \\ \alpha\beta + \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + \frac{1}{\alpha\beta} & \text{if } q+s \text{ is odd.} \end{cases}$$

In particular, if $\alpha = \beta = 1$, then

$$\begin{aligned} (5.2) \quad \mathrm{tr}[W_{p/q}, W_{r/s}] &= \Phi_{p/q}^2 + \Phi_{r/s}^2 + \Phi_{(p/q)\oplus(r/s)}^2 + \Phi_{p/q}\Phi_{r/s}\Phi_{(p/q)\oplus(r/s)} \\ &\quad - 4(\Phi_{p/q}\Phi_{r/s} + 2\Phi_{(p/q)\oplus(r/s)}) + 14. \end{aligned}$$

In the case of the Maskit slice, the commutator of the equivalent of neighbouring Farey words ('slope words') is always -2 , since in that case the commutator represents a closed curve about the puncture on the torus. Combining the Fricke identity with the equivalent to Lemma 4.5 in this case gives that Farey triplets (f, g, h) of the trace polynomials of the slope words on the punctured torus satisfy the **Markoff equation**

$$f^2 + g^2 + h^2 = 3fgh.$$

A little experimentation shows that no nice identity holds for the commutators of Farey words in a Riley group—nor should we expect any, since the product of two Farey words is usually strictly loxodromic for any fixed ρ ; there are Markoff-like theories for the 4-times punctured sphere, and one is the theory of Beardon which

depends on picking hyperbolic representatives for the simple closed curves [BLS86], but the more relevant to us is the theory of Maloni, Palesi, and Tan [MPT15] which (as we briefly mentioned above) recovers our recursion: in §2.8 *op. cit.*, take $\mu = (p, q, r, s) = (8, 8, 8, -28)$ to recover the recursion (their Equations (5)–(7)) and so obtain the μ -Markoff equation

$$f^2 + g^2 + h^2 + fgh = 8f + 8g + 8h - 28.$$

In any case we can generalise in a combinatorial manner the process used to compute the commutator in the Maskit case to obtain a second expression for the commutator (and bypass the Markoff theory) in the Riley slice case by using both parts of Lemma 4.2 together. By that lemma,

$$[W_{p/q}, W_{r/s}] = W_{p/q} W_{r/s} (W_{r/s} W_{p/q})^{-1} = \widehat{W}_{(p/q)\oplus(r/s)} \cdot \tilde{W}_{(p/q)\oplus(r/s)}^{-1}$$

where the hat here indicates that the $(q+s)$ th exponent is flipped in sign and the tilde indicates a flip in the $(2s)$ th exponent. We therefore have two cases: either $q+s < 2s$, or $2s < q+s$. Without loss of generality, assume that $q+s < 2s$ and write $W_{p/q\oplus r/s} = u_1 t u_2 t' u_3$ where u_1 is the word made up of the first $(q+s-1)$ letters, t is the $(q+s)$ th letter, u_2 is the $(q+s+1)$ th to $(2s-1)$ th letters, t' is the $(2s)$ th letter, and u_3 is the remainder of the word; then

$$\begin{aligned} \text{tr}[W_{p/q}, W_{r/s}] &= \text{tr } u_1 T u_2 t' u_3 (u_1 t u_2 T' u_3)^{-1} \\ &= \text{tr } u_1 T u_2 t' u_3 U_3 t' U_2 T U_1 = \text{tr } u_1 T u_2 (t')^2 U_2 T U_1. \end{aligned}$$

By rotational symmetry, this last trace is equal to $\text{tr } u_2 (t')^2 U_2 T^2$. By assumption, t and t' are parabolic so there exists some transformation $M \in \text{PSL}(2, \mathbb{C})$ (perhaps the identity) such that $mt'M = t$. Writing $u = u_2 m$, the trace becomes $\text{tr } ut^2 UT^2 = \text{tr}[u, t^2]$. We now use the Fricke identity again, to see that

$$\begin{aligned} \text{tr}[u, t^2] &= \text{tr}^2 u + \text{tr}^2 t^2 + \text{tr}^2 ut^2 - \text{tr } u \text{tr } t^2 \text{tr } ut^2 - 2 \\ &= \text{tr}^2 u + \text{tr}^2 ut^2 - 2 \text{tr } u \text{tr } ut^2 + 2 \\ &= (\text{tr } u - \text{tr } ut^2)^2 + 2. \end{aligned}$$

This proves the following result, which may be known to experts:

Lemma 5.3. *If p/q and r/s are Farey neighbours, then $\text{tr}[W_{p/q}(z), W_{r/s}(z)] - 2$ is a square in $\mathbb{Z}[z]$.* \square

As an amusing corollary, we see that the complicated expression to the right of Equation (5.2) is a positive function.

6. CLOSED-FORM SOLUTIONS FOR SOME CASES

Recall that the **Chebyshev polynomials** (of the first kind) are the family of polynomials T_n defined via the recurrence relation


$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x). \end{aligned}$$

It is well-known that these polynomials satisfy the product relation

$$2T_m(x)T_n(x) = T_{m+n}(x) + T_{|m-n|}(x)$$

for $m, n \in \mathbb{Z}_{\geq 0}$. Compare this relation with the relation Equation (4.10) developed above for the parabolic Farey polynomials (but note that the Chebyshev product rule holds for all m, n and the identities for the Farey polynomials hold only for Farey neighbours).

We may apply the theory of ‘Farey recursive functions’ [Che+20; Che20] in order to explain this analogy. The following diagram may be useful for translating the notation of that paper (right) into the notation we use here (left):



Definition 6.1 (Definition 3.1 of [Che+20]). Let R be a (commutative) ring, and suppose $d_1, d_2 : \hat{\mathbb{Q}} \rightarrow R$. A function $\mathcal{F} : \hat{\mathbb{Q}} \rightarrow R$ is a **(d_1, d_2)-Farey recursive function** if, whenever $\alpha, \beta \in \hat{\mathbb{Q}}$ are Farey neighbours,

$$(6.2) \quad \mathcal{F}(\beta \oplus \alpha) = -d_1(\alpha)\mathcal{F}(\beta \ominus \alpha) + d_2(\alpha)\mathcal{F}(\beta).$$

Observe that the relation Equation (4.10) looks essentially of this form; to make this clearer, we rewrite it slightly as

$$(6.3) \quad \Phi(\beta \oplus \alpha) = 8 - \Phi(\beta \ominus \alpha) - \Phi(\alpha)\Phi(\beta)$$

(where we set $\beta = p/q$, $\alpha = r/s$, and changed from subscript notation to functional notation). In our case, then, $d_1(\alpha)$ is constantly 1 and $d_2 = -\Phi$. Note also that our relation is not homogeneous. We therefore adapt the definition of [Che+20] to the following:

Definition 6.4. Let R be a (commutative) ring, and suppose $d_1, d_2, d_3 : \hat{\mathbb{Q}} \rightarrow R$. A function $\mathcal{F} : \hat{\mathbb{Q}} \rightarrow R$ is a **(d_1, d_2)-Farey recursive function** if, whenever $\alpha, \beta \in \hat{\mathbb{Q}}$ are Farey neighbours,

$$(6.5) \quad \mathcal{F}(\beta \oplus \alpha) = -d_1(\alpha)\mathcal{F}(\beta \ominus \alpha) + d_2(\alpha)\mathcal{F}(\beta) + d_3(\alpha).$$

If d_3 is the zero function, we say that the relation of Equation (6.5) is **homogeneous**, otherwise it is **non-homogeneous**.

The relevant generalisations of the existence-uniqueness results of [Che+20, Section 4] follow easily (the same proofs work, with the usual property that the space of non-homogenous solutions is the sum of a particular solution and the space of homogeneous solutions).

In our case, there is an obvious explicit solution to the non-homogeneous Farey polynomial recursion, Equation (6.3): namely, the map Φ which sends every $\alpha \in \mathbb{Q}$ to the constant polynomial $2 \in \mathbb{Z}[z]$. It therefore remains to solve the corresponding homogeneous equation.

6.1. A Fibonacci-like subsequence of the homogeneous Farey polynomials.

Notation. In this section, we work exclusively with the parabolic Farey polynomials, $\Phi_{p/q}^{\infty, \infty}$. We will reuse the symbols α and β for rational numbers (since the roots of unity used to define X and Y become 1 in this situation and so there is no need for special notation for them).

In Table 4, we list the first few polynomials Φ^h which solve the homogeneous recursion relation

$$(6.6) \quad \Phi^h(\beta \oplus \alpha) = -\Phi^h(\beta \ominus \alpha) - \Phi^h(\alpha)\Phi^h(\beta)$$

with the initial values $\Phi^h(0/1) = 2 - z$, $\Phi^h(1/0) = 2$, and $\Phi^h(1/1) = 2 + z$.

The polynomials with numerator 1 listed in the table have very nice properties: immediately one sees that the constant terms alternate in sign and increase in

TABLE 4. Selected polynomials Φ^h satisfying the homogeneous recursion relation for small q , with the initial values as given.

p	q	$\Phi_{p/q}^h$
1	0	2
0	1	$-z + 2$
1	1	$z + 2$
1	2	$z^2 - 6$
1	3	$z^3 - 2z^2 - 7z + 10$
2	3	$-z^3 - 2z^2 + 7z + 10$
1	4	$z^4 - 4z^3 - 4z^2 + 24z - 14$
3	4	$z^4 + 4z^3 - 4z^2 - 24z - 14$
1	5	$z^5 - 6z^4 + 3z^3 + 34z^2 - 55z + 18$
2	5	$-z^5 + 2z^4 + 13z^3 - 22z^2 - 41z + 58$
3	5	$z^5 + 2z^4 - 13z^3 - 22z^2 + 41z + 58$
4	5	$-z^5 - 6z^4 - 3z^3 + 34z^2 + 55z + 18$
1	6	$z^6 - 8z^5 + 14z^4 + 32z^3 - 119z^2 + 104z - 22$
1	7	$z^7 - 10z^6 + 29z^5 + 10z^4 - 186z^3 + 308z^2 - 175z + 26$
1	8	$z^8 - 12z^7 + 48z^6 - 40z^5 - 220z^4 + 648z^3 - 672z^2 + 272z - 30$
1	9	$z^9 - 14z^8 + 71z^7 - 126z^6 - 169z^5 + 1078z^4 - 1782z^3 + 1308z^2 - 399z + 34$
1	10	$z^{10} - 16z^9 + 98z^8 - 256z^7 + 35z^6 + 1456z^5 - 3718z^4 + 4224z^3 - 2343z^2 + 560z - 38$

magnitude by 4 each time; also, we have that $\Phi_{1/q}^h(1)$ cycles through the values 3, -5 , 2, $\Phi_{1/q}^h(2)$ cycles through the values 4, -2 , -4 , 2; and when we evaluate at 3 and 4 we get a 6-cycle and an arithmetic sequence of step 4 respectively. When we consider $\Phi^h(1/q)(5)$, though, we obtain more interesting behaviour: this is OEIS sequence A100545³ and satisfies the Fibonacci-type relation

$$\Phi_{1/q}^h(5) = 3\Phi_{1/(q-1)}^h(5) - \Phi_{1/(q-2)}^h(5) \quad \text{with } \Phi_{1/1}^h(5) = 7, \Phi_{1/2}^h(5) = 19.$$

Of course, from the way that we defined the Φ^h such types of relations ought to be expected. In this section, we use the standard diagonalisation technique to explain the behaviour of the sequence $a_q := \Phi^h(1/q)(z)$ for fixed $z \in \mathbb{C}$. From Equation (6.6), we have that

$$(6.7) \quad a_q = -(2-z)a_{q-1} - a_{q-2}.$$

We may rewrite this equation in matrix form as the following:

$$(6.8) \quad \begin{bmatrix} 0 & 1 \\ -1 & z-2 \end{bmatrix} \begin{bmatrix} a_{q-2} \\ a_{q-1} \end{bmatrix} = \begin{bmatrix} a_{q-1} \\ a_q \end{bmatrix}.$$

One easily computes that the eigenvalues of the transition matrix are

$$\lambda^\pm = \frac{z-2 \pm \alpha}{2}$$

(where $\alpha = \sqrt{z^2 - 4z}$) with respective eigenvectors

$$v^\pm = \begin{bmatrix} z-2 \mp \alpha \\ 2 \end{bmatrix}$$

(note the alternated sign). Thus the transition matrix may be diagonalised as

$$(6.9) \quad \frac{-1}{2\alpha} \begin{bmatrix} z-2-\alpha & z-2+\alpha \\ 2 & 2 \end{bmatrix} \begin{bmatrix} \frac{z-2+\alpha}{2} & 0 \\ 0 & \frac{z-2-\alpha}{2} \end{bmatrix} \begin{bmatrix} 2 & 2-z-\alpha \\ -2 & z-2-\alpha \end{bmatrix}$$

³<http://oeis.org/A100545>

and so a_q is the first coordinate of

$$\frac{-1}{2\alpha} \begin{bmatrix} z-2-\alpha & z-2+\alpha \\ 2 & 2 \end{bmatrix} \begin{bmatrix} \left(\frac{z-2+\alpha}{2}\right)^q & 0 \\ 0 & \left(\frac{z-2-\alpha}{2}\right)^q \end{bmatrix} \begin{bmatrix} 2 & 2-z-\alpha \\ -2 & z-2-\alpha \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \end{bmatrix};$$

expanding out, we get

$$(6.10) \quad a_q = 2^{-1-q}\alpha^{-1} \left((a_0(z-2+\alpha)-2a_1)(z-2-\alpha)^q + (a_0(2-z+\alpha)+2a_1)(z-2+\alpha)^q \right).$$

We may also characterise the z for which $\Phi_{1/q}^h(z)$ is cyclic: this occurs precisely when the diagonal matrix of Equation (6.9) is of finite order, i.e. whenever both $(z-2\pm\alpha)/2$ are roots of unity.

As an application of the theory above, we have seen that the Chebyshev polynomials also satisfy a second-order recurrence relation with transition matrix

$$\begin{bmatrix} 0 & 1 \\ -1 & 2x \end{bmatrix}.$$

If we set $z = 2x+2$, then we get back our transition matrix from Equation (6.8). Thus our sequence $\Phi_{1/q}^h(z)$ for fixed z is of the form $W_q(\frac{z-2}{2})$ where W_q is the q th Chebyshev polynomial in the sequence beginning with $W_0 = 2x$ and $W_1 = 2x+4$.

Finally, we consider the solution of the non-homogeneous equation for $\Phi_{1/q}$. Above, we observed that there is a constant solution to the global recursion relation on the entire Stern-Brocot tree; we therefore guess that there is a similar solution to this recursion. Such a solution f will satisfy

$$8 = f(z) + (2-z)f(z) + f(z)$$

and arithmetic gives $f(z) = 8/(4-z)$. Combining this with Equation (6.10) above gives us the following general solution to the non-homogeneous relation:

$$a_q = \frac{8}{4-z} + 2^{-1-q}\alpha^{-1} \left((\lambda(z-2+\alpha)-2\mu)(z-2-\alpha)^q + (\lambda(2-z+\alpha)+2\mu)(z-2+\alpha)^q \right).$$

In our case, we have $a_0 = \Phi_{1/0}(z) = 2$ and $a_1 = \Phi_{1/1}(z) = 2+z$. Solving the resulting system of equations gives

$$(\lambda, \mu) = \left(\frac{2z}{z-4}, \frac{2z-z^2}{z-4} \right)$$

and hence

$$a_q = \frac{8}{4-z} + \frac{2^{-q}z}{z-4} \left((-2+z-\sqrt{z^2-4z})^q + (-2+z+\sqrt{z^2-4z})^q \right)$$

6.2. Solving the homogeneous recursion relation in general. In the previous section, we computed a closed form formula for $\Phi_{1/q}^h(n)$ using standard techniques from the theory of second-order linear recurrences. We now tackle the general problem of finding a closed-form formula for $\Phi_{p/q}^h(n)$; in order to do this, we use the theory of Section 6 of [Che+20] but with a slight modification: in that paper, the authors define a special case of Farey recursive function, a **Farey recursive function of determinant d** (where $d : \hat{\mathbb{Q}} \rightarrow R$), to be a Farey recursive function \mathcal{F} with $d_1 = d$ and $d_2 = \mathcal{F}$. That is, they replace the recurrence of Equation (6.2) with

$$\mathcal{F}(\beta \oplus \alpha) = -d(\alpha)\mathcal{F}(\beta \ominus \alpha) + \mathcal{F}(\alpha)\mathcal{F}(\beta).$$

This is very similar to our situation, except that instead of $d_2 = \mathcal{F}$ we have $d_2 = -\mathcal{F}$. To reflect this, we shall call a Farey recursive function satisfying a relation of the form

$$(6.11) \quad \mathcal{F}(\beta \oplus \alpha) = -d(\alpha)\mathcal{F}(\beta \ominus \alpha) - \mathcal{F}(\alpha)\mathcal{F}(\beta).$$

a Farey recursive function of **anti-determinant** d . We shall work for the time being in this setting (i.e. we shall work with the general function \mathcal{F} rather than the particular example Φ) in order to restate in sufficient generality the theorem which we need (Theorem 6.1 of [Che+20]).

Let $\alpha \in \mathbb{Q}$. The **boundary sequence** $\partial(\alpha)$ is defined inductively by the process of ‘continuing to expand down the Farey graph by constant steps’. More precisely, let $\beta \oplus^k \gamma$ denote $((\beta \oplus \underbrace{\gamma \oplus \dots}_{k \text{ iterates}}) \oplus \gamma)$ for $\beta, \gamma \in \hat{\mathbb{Q}}$ and let γ_L, γ_R be the unique Farey neighbours such that $\alpha = \gamma_L \oplus \gamma_R$; then we set

$$\partial(\alpha) := \{\gamma_L \oplus^k \alpha : k \in \mathbb{Z}_{\geq 0}\} \cup \{\gamma_R \oplus^k \alpha : k \in \mathbb{Z}_{\geq 0}\}.$$

If we allow the Farey graph to embed in the Euclidean upper halfplane by sending $\mathbb{Q} \ni p/q \mapsto (p/q, 1/q) \in \mathbb{H}^2$, then except for the exceptional cases $\alpha = 1/0$ and $\alpha = n/1$ for $n \in \mathbb{Z}$ the subgraph spanned by $\partial(\alpha)$ corresponds to a Euclidean triangle containing α in its interior, see Figure 3 of [Che+20]; for example, the triangle spanned by $\partial(1/n)$ is the triangle with vertices $0, (1/2, 1/2), 1$.

It will be useful to have specific names for the terms in each of the two subsequences and so we set, for $k \in \mathbb{Z}$,

$$(6.12) \quad \beta_k := \begin{cases} \gamma_L \oplus^{-k-1} \alpha & \text{if } k < -1 \\ \gamma_L & \text{if } k = -1 \\ \gamma_R & \text{if } k = 0 \\ \gamma_R \oplus^k \alpha & \text{if } k > 0. \end{cases}$$

For every $\alpha \in \mathbb{Q}$, define

$$(6.13) \quad M_\alpha = \begin{bmatrix} 0 & 1 \\ -d(\alpha) & \mathcal{F}(\alpha) \end{bmatrix}.$$

Given any Farey neighbour β of α , we have

$$M_\alpha^n \begin{bmatrix} \mathcal{F}(\gamma \oplus^0 \alpha) \\ \mathcal{F}(\gamma \oplus^1 \alpha) \end{bmatrix} = \begin{bmatrix} \mathcal{F}(\gamma \oplus^n \alpha) \\ \mathcal{F}(\gamma \oplus^{n+1} \alpha) \end{bmatrix}$$

and so the recursion Equation (6.11) is equivalent to a family of second-order linear recurrences, one down $\partial(\alpha)$ for each α .

We may now state the following theorem:

Theorem 6.14 (Adaptation of Theorem 6.1 of [Che+20]). *Let $d : \hat{\mathbb{Q}} \rightarrow R$ be a multiplicative function (in the sense that $d(\gamma \oplus \beta) = d(\gamma)d(\beta)$ for all pairs of Farey neighbours $\beta, \gamma \in \mathbb{Q}$) to a commutative ring R , such that $d(\hat{\mathbb{Q}})$ contains no zero divisors. Suppose that \mathcal{F} is a Farey recursive function with anti-determinant d . Given $\alpha \in \mathbb{Q}$, define M_α as in Equation (6.13) and $(\beta_k)_{k \in \mathbb{Z}}$ as in Equation (6.12).*

Then, for all $n \in \mathbb{Z}$,

$$M_\alpha^n \begin{bmatrix} \mathcal{F}(\beta_0) \\ \mathcal{F}(\beta_1) \end{bmatrix} = \begin{cases} \begin{bmatrix} \mathcal{F}(\beta_n) \\ \mathcal{F}(\beta_{n+1}) \end{bmatrix} & n \geq 0, \\ \begin{bmatrix} \frac{1}{d(\beta_{-1})} \mathcal{F}(\beta_{-1}) \\ \mathcal{F}(\beta_0) \end{bmatrix} & n = -1, \text{ and} \\ \begin{bmatrix} \frac{1}{d(\beta_{-1})d_\alpha^{-n-1}} \mathcal{F}_{\beta_n} \\ \frac{1}{d(\beta_{-1})d_\alpha^{-n-2}} \mathcal{F}_{\beta_{n+1}} \end{bmatrix} & n < -1. \end{cases}$$

We proceed to prove Theorem 6.14 by exactly the same argument as given in [Che+20]. The key point is the following lemma, which is the analogue of the discussion directly preceding the statement of Theorem 6.1 in that paper.

Lemma 6.15. *With the setup of Theorem 6.14, we have*

$$\begin{aligned} M_\alpha^{-1} \begin{bmatrix} \mathcal{F}(\beta_0) \\ \mathcal{F}(\beta_1) \end{bmatrix} &= \begin{bmatrix} \frac{d(\beta_0)}{d(\alpha)} \mathcal{F}(\beta_{-1}) \\ \mathcal{F}(\beta_0) \end{bmatrix} \\ M_\alpha^{-2} \begin{bmatrix} \mathcal{F}(\beta_0) \\ \mathcal{F}(\beta_1) \end{bmatrix} &= \begin{bmatrix} \frac{1}{d(\alpha)d(\beta_{-1})} \mathcal{F}(\beta_{-2}) \\ \frac{1}{d(\beta_{-1})} \mathcal{F}(\beta_{-1}) \end{bmatrix} \end{aligned}$$

Proof. The formula involving M_α^{-1} comes directly from computing the product on the left via the definition and simplifying with the formula

$$\mathcal{F}(\beta_1) = -d(\beta_0)\mathcal{F}(\beta_{-1}) - \mathcal{F}(\alpha)\mathcal{F}(\beta_0)$$

which is almost exactly the same as Equation (8) of [Che+20]—the single sign change cancels exactly with the sign change between the ‘determinant’ and ‘anti-determinant’ recurrences so we get the same overall formula for the M_α^{-1} product as they do in Equation (11) of their paper.

The formula for M_α^{-2} comes from applying the analogues of Equations (9) and (10) of their paper,

$$\begin{aligned} \mathcal{F}(\beta_{-2}) &= -d(\beta_{-1})\mathcal{F}(\beta_0) - \mathcal{F}(\alpha)\mathcal{F}(\beta_{-1}) \\ d(\alpha) &= d(\beta_{-1})d(\beta_0) \end{aligned}$$

and simplifying; again the minus signs cancel and we get the same formula. \square

Proof of Theorem 6.14. The formula for $n \geq 0$ holds for all Farey recursive formulae as noted above; the formulae for $n = -1$ and $n = -2$ are just the formulae of Lemma 6.15; and we proceed to prove the formula for $n < -2$ by induction. Assume that the formula holds for some fixed $n \leq -2$; then from the definitions we have

$$\mathcal{F}(\beta_{n-1}) = -\mathcal{F}(\alpha)\mathcal{F}(\beta_n) - d(\alpha)\mathcal{F}(\beta_{n+1})$$

and so we can compute

$$\begin{aligned} M_\alpha^{n-1} \begin{bmatrix} F(\beta_0) \\ F(\beta_1) \end{bmatrix} &= M_\alpha^{-1} M_\alpha^n \begin{bmatrix} F(\beta_0) \\ F(\beta_1) \end{bmatrix} \\ &= \frac{1}{d(\alpha)} \begin{bmatrix} -\mathcal{F}(\alpha) & -1 \\ d(\alpha) & -0 \end{bmatrix} \begin{bmatrix} \frac{1}{d(\beta_{-1})d(\alpha)^{-n-1}} F(\beta_n) \\ \frac{1}{d(\beta_{-1})d(\alpha)^{-n-2}} F(\beta_{n+1}) \end{bmatrix} \\ &= \frac{1}{d(\alpha)} \begin{bmatrix} -\frac{1}{d(\beta_{-1})d(\alpha)^{-n-1}} (\mathcal{F}(\alpha)\mathcal{F}(\beta_n) + d(\alpha)\mathcal{F}(\beta_{n+1})) \\ \frac{1}{d(\beta_{-1})d(\alpha)^{-n-2}} \mathcal{F}(\beta_n) \end{bmatrix} \\ &= \begin{bmatrix} -\frac{1}{d(\beta_{-1})d(\alpha)^{-n}} \mathcal{F}(\beta_{n-1}) \\ \frac{1}{d(\beta_{-1})d(\alpha)^{-n-1}} \mathcal{F}(\beta_n) \end{bmatrix} \end{aligned}$$

which is the desired result. \square

Corollary 6.16 (Adaptation of Corollary 6.2 of [Che+20]). *Let Φ^h be a family of homogeneous Farey polynomials (i.e. a family solving Equation (6.6) for some starting values). Then, for some $\alpha \in \mathbb{Z}$, if M_α is the matrix*

$$\begin{bmatrix} 0 & 1 \\ -1 & \Phi^h(\alpha) \end{bmatrix}$$

and if $(\beta_k)_{k \in \mathbb{Z}}$ are the boundary values about α as in Equation (6.12), then for all $n \in \mathbb{Z}$ we have

$$M_\alpha^n \begin{bmatrix} \Phi^h(\beta_0) \\ \Phi^h(\beta_1) \end{bmatrix} = \begin{bmatrix} \Phi^h(\beta_n) \\ \Phi^h(\beta_{n+1}) \end{bmatrix}.$$

Proof. This follows directly from Theorem 6.14 with the observation that the anti-determinant of Φ^h is the constant function $d(\gamma) = 1$ for all $\gamma \in \mathbb{Q}$. \square

Thus to determine Φ_α^h for all $\alpha \in \mathbb{Q}$ it suffices to compute and diagonalise the M_α matrices, using the techniques of Section 6.1. (Of course, we need to diagonalise in the ring of rational functions over \mathbb{Q} rather than the ring of polynomials over \mathbb{Z} .) More precisely, we need to compute M_{α_i} for some family (α_i) of rationals with the property that the boundary sets $\partial(\alpha_i)$ cover \mathbb{Q} . (In Section 6.1, we did this computation for $\partial(0/1)$.)

In any case, from Corollary 6.16 we immediately have a qualitative result:

Theorem 6.17. *For any $\gamma \in \mathbb{Q}$, there exists a sequence $\dots, \gamma_{-1}, \gamma_0 = \gamma, \gamma_1, \gamma_2, \dots$ of rational numbers such that $\Phi_{\gamma_n}^h(z)$ is a sequence of Chebyshev polynomials $W_n(\Phi_\gamma^h(z)/2)$. (Namely, let γ_{-1} be a neighbour in the Stern-Brocot tree of γ and take the sequence (γ_k) to be precisely the sequence (β_k) of Equation (6.12) with $\alpha := \gamma \ominus \gamma_{-1}$.)* \square

Remark. Of course, the boundary sequence (γ_k) constructed here is just a geodesic line Λ in the Stern-Brocot tree rooted at γ , defined by choosing one vertical half-ray in the tree starting from γ (where ‘vertical’ refers to the embedding of Figure 7) and then extending that in the Farey graph in the obvious way by repeated Farey arithmetic with the same difference. There are clearly two such natural choices for Λ given a fixed γ (γ has three neighbours, but two correspond to the same geodesic), and a single natural choice is obtained by taking the unique neighbour of γ which lies above.

We easily compute that the eigenvalues of M_α are

$$\lambda^\pm = \frac{1}{2} \left(\Phi_\alpha^h \pm \sqrt{(\Phi_\alpha^h)^2 - 4} \right).$$

Let $x = \Phi_\alpha^h$ and $\kappa = \sqrt{x^2 - 4}$ (this is the analogue of the constant α from Section 6.1); then the respective eigenvectors are

$$v^\pm = \begin{bmatrix} x \mp \kappa \\ 2 \end{bmatrix}.$$

We therefore may diagonalise M_α as

$$M_\alpha = -\frac{1}{4\kappa} \begin{bmatrix} x - \kappa & x + \kappa \\ 2 & 2 \end{bmatrix} \begin{bmatrix} \frac{1}{2}(x + \kappa) & 0 \\ 0 & \frac{1}{2}(x - \kappa) \end{bmatrix} \begin{bmatrix} 2 & -x - \kappa \\ -2 & x - \kappa \end{bmatrix};$$

in particular, $\Phi^h(\beta_n)$ is the first component of

$$\begin{aligned} M_\alpha^n \begin{bmatrix} \Phi^h(\beta_0) \\ \Phi^h(\beta_1) \end{bmatrix} \\ = -\frac{1}{4\kappa} \begin{bmatrix} x-\kappa & x+\kappa \\ 2 & 2 \end{bmatrix} \begin{bmatrix} \frac{1}{2^n}(x+\kappa)^n & 0 \\ 0 & \frac{1}{2^n}(x-\kappa)^n \end{bmatrix} \begin{bmatrix} 2 & -x-\kappa \\ -2 & x-\kappa \end{bmatrix} \begin{bmatrix} \Phi^h(\beta_0) \\ \Phi^h(\beta_1) \end{bmatrix} \end{aligned}$$

computing this, we have

$$\Phi^h(\beta_n) = \frac{(\Phi^h(\beta_0)(x+\kappa) - 2\Phi^h(\beta_1))(x-\kappa)^n + (\Phi^h(\beta_0)(\kappa-x) + 2\Phi^h(\beta_1))(x+\kappa)^n}{2^{1+n}\kappa}.$$

In particular, we have proved the following quantitative improvement of Theorem 6.17:

Theorem 6.18. *Let β_0 and β_1 be Farey neighbours, and let $\alpha = \beta_1 \ominus \beta_0$. Then we have a closed form formula for $\Phi^h(\beta_n)$ ($n \in \mathbb{Z}$), namely*

$$\Phi_{\beta_n}^h = \frac{(\Phi_{\beta_0}^h(\Phi_\alpha^h + \kappa) - 2\Phi_{\beta_1}^h)(\Phi_\alpha^h - \kappa)^n + (\Phi_{\beta_0}^h(\kappa - \Phi_\alpha^h) + 2\Phi_{\beta_1}^h)(\Phi_\alpha^h + \kappa)^n}{2^{1+n}\kappa}.$$

where $\kappa = \sqrt{(\Phi_\alpha^h)^2 - 4}$. \square

This gives a ‘local’ closed form solution for the recursion around any $\alpha \in \mathbb{Q}$; a ‘global’ solution corresponds to a collection of these solutions, each local to a particular geodesic in the graph and which are compatible on intersections. Unfortunately, our original recurrence relied on knowing only three initial values globally in the graph; while this local formula relies on knowing three initial values which are local on the particular geodesic.

7. APPROXIMATING IRRATIONAL PLEATING RAYS

As we mentioned in the introduction to [EMS23a], a version of this theory can be used to give approximations to irrational pleating rays. In order to do this, we must deal with the theory of infinite continued fractions.

Every rational number can be expressed as a finite simple continued fraction: if (a_0, a_1, \dots, a_k) is a finite sequence of integers, we define the simple continued fraction

$$[a_0; a_1, \dots, a_k] := a_0 + \cfrac{1}{a_1 + \cfrac{1}{a_2 + \cfrac{1}{\ddots + \cfrac{1}{a_n + \cfrac{1}{\ddots + \cfrac{1}{a_k}}}}}}$$

and every rational number p/q has exactly two expressions as such a continued fraction [HW60, Theorem 162]. It is also a well-known result in classical number theory that every *irrational* $\lambda \in \mathbb{R} \setminus \mathbb{Q}$ has a unique simple continued fraction approximation of the form

$$\lambda = [a_0, a_1, \dots, a_n, \dots] = a_0 + \cfrac{1}{a_1 + \cfrac{1}{\ddots + \cfrac{1}{a_n + \cfrac{1}{\ddots}}}}$$

which can be computed efficiently by repeated application of the Euclidean algorithm (see, for example, §10.9 of [HW60]). We now show that this exhibits λ as a limit of a sequence of rationals ‘down the Farey tree’.

Proposition 7.1. Suppose that $p/q = [a_1, \dots, a_{N-1}, a_N, 1]$; define

$$\frac{r_1}{s_1} = [a_1, \dots, a_{N-1}, a_N] \text{ and } \frac{r_2}{s_2} = [a_1, \dots, a_{N-1}].$$

Then r_1/s_1 and r_2/s_2 are Farey neighbours and $p/q = (r_1/s_1) \oplus (r_2/s_2)$.

Proof. That the Farey sum is as claimed follows from Theorem 149 of [HW60]: that is, if $p_n/q_n = [a_1, \dots, a_n]$ then

$$p_n = a_n p_{n-1} + p_{n-2} \text{ and } q_n = a_n q_{n-1} + q_{n-2}.$$

Indeed, take $p/q = p_{N+1}/q_{N+1} = [a_1, \dots, a_{N-1}, a_N, 1]$, then $a_{N+1} = 1$ so $p_{N+1} = 1p_N + p_{N-1}$ and $q_{N+1} = q_N + q_{N-1}$.

That the two are Farey neighbours is exactly Theorem 150 of [HW60], which actually gives slightly more information:

$$p_N q_{N-1} - p_{N-1} q_N = (-1)^{N-1}. \quad \square$$

In the previous section, we indicated how to compute in closed form the sequence of Farey polynomials corresponding to the Farey fractions

$$\frac{p_1}{q_1}, \frac{p_2}{q_2} = \frac{p_1}{q_1} \oplus \left(\frac{p_2}{q_2} \ominus \frac{p_1}{q_1} \right), \dots, \frac{p_n}{q_n} = \frac{p_1}{q_1} \oplus^{n-1} \left(\frac{p_2}{q_2} \ominus \frac{p_1}{q_1} \right)$$

where p_1/q_1 and p_2/q_2 are Farey neighbours. That is, we gave a way to compute the Farey polynomials down a branch of the Farey tree with constant difference (for instance, we gave the example of $\Phi_{1/q}$, where $\frac{1}{q} = \frac{1}{0} \oplus^q \frac{0}{1}$). The study of partial fraction decompositions here gives, in general, different sequences: the constant addition sequence rooted at an element ξ in the tree is the sequence which constantly chooses the *left* branch when moving down from ξ (with respect to the embedding of Figure 7), while the sequence corresponding to continually adding the previous two items in the tree (and therefore building a continued fraction decomposition) corresponds to the sequence which is eventually constantly moving *rightwards*. For this reason, we might (rather foolishly, as it turns out) expect there to also exist a nice way to compute closed-form expressions for sequences of Farey polynomials corresponding to finite convergents of infinite continued fraction decompositions, and therefore for there to be a reasonable way to approximate irrational pleating rays and attempt to compute expressions for the analytic functions of which they are subsets of zero-sets.

Remark. The geometric meaning of all this comes from the theory of two-bridge links and is described in [BZ03, Chapter 12]; one is really writing down a sequence of 2-bridge links which converge to the link of interest, increasing by one crossing in each step.


The recursion relation of interest comes, as always, from applying the Farey polynomial operator to the recursion relation $a_n = a_{n-1} \oplus a_{n-2}$; doing this we obtain

$$\Phi_{a_n} = \Phi_{a_{n-1} \oplus a_{n-2}} = 8 - \Phi_{a_{n-1} \ominus a_{n-2}} - \Phi_{a_{n-1}} \Phi_{a_{n-2}}.$$

Observe that $a_{n-1} \ominus a_{n-2}$ is just a_{n-3} and replace the cumbersome notation Φ_{a_k} with x_k to get the relation

$$(7.2) \quad x_n = 8 - x_{n-3} - x_{n-2}x_{n-1}.$$

Unfortunately, this is a non-linear recurrence relation—this means that it is harder to study than the linear left-recursion studied in the previous section, and we are unable to find a closed-form solution. (We believe that if one exists then it will be in terms of combinatorial objects, e.g. Stirling numbers.) However, it does at least give a computationally feasible method for approximating irrational cusp points.

FIGURE 8. The zeros of $\Phi_{\text{fib}(p-1)/\text{fib}(p)} + 2$ for $p \in \{1, \dots, 16\}$.

Example 7.3 (The Fibonacci polynomials). Let us compute an approximation to the pleating ray with asymptotic angle π/ϕ , where ϕ is the golden ratio $(\sqrt{5}+1)/2$. It is well-known that

$$\phi = [1, 1, \dots] = 1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}}$$

and so by Proposition 7.1 we get that $1/\phi$ is approximated by

$$[0, 1] = 1, [0, 1, 1] = 1/2, [1, 1, 1] = 2/3, \dots;$$

in other words, by the Fibonacci fractions $\text{fib}(q-1)/\text{fib}(q)$. Being efficiency-minded, we have already computed the corresponding polynomials, which are listed in Table 3; the inverse images of -2 under the first 16 such polynomials are shown in Figure 8, where the $1/\phi$ -cusp is approximated by the ‘corner’ point, at the top-left of each picture. Quite a good approximation seems to be given after only a few pictures, which is good since after about 15 terms the polynomial coefficients and degrees become too big for both **Mathematica** and **MATLAB** to study without hassle. We also note that eventually the cusp point occurs at quite a ‘sharp’ point in the set; perhaps it would be interesting to use this to distinguish the cusp computationally from the other points in the inverse image.

Example 7.4 ($\sqrt{2}$). For our next trick, we consider another irrational number with a nice continued fraction decomposition:

$$\sqrt{2} = [1, 2, 2, \dots, 2, \dots]$$




FIGURE 9. $(\Phi_{19/27}^{\infty, \infty})^{-1}(-2)$, approximating the $1/\sqrt{2}$ cusp point.

TABLE 5. Eigenproperties of the right-recurrence relation.

fixed point	eigenvalue	eigenvector
-4	$\frac{5+\sqrt{21}}{2}$	$\left(-\frac{5+\sqrt{21}}{5+\sqrt{21}}, \frac{2}{5+\sqrt{21}}, 1 \right)^t$
	-1	$(1, -1, 1)^t$
	$\frac{5-\sqrt{21}}{2}$	$\left(-\frac{5+\sqrt{21}}{5+\sqrt{21}}, -\frac{2}{5+\sqrt{21}}, 1 \right)^t$
	determinant:	-1
2	$\frac{-1+i\sqrt{3}}{2}$	$\left(\frac{-1+i\sqrt{3}}{2}, \frac{-1-i\sqrt{3}}{2}, 1 \right)^t$
	-1	$(1, -1, 1)^t$
	$\frac{-1-i\sqrt{3}}{2}$	$\left(\frac{-1-i\sqrt{3}}{2}, \frac{-1+i\sqrt{3}}{2}, 1 \right)^t$
	determinant:	-1

so we may approximate $1/\sqrt{2}$ by the Farey sequence beginning

$$1, 2/3, 5/7, 7/10, 12/17, \dots$$

The corresponding Farey polynomials are higher degree and so longer to write down than the Fibonacci polynomials, but it is easy enough for the computer to plot the preimages of -2 for the first 10 or so polynomials; we give the preimage for $\Phi_{19/27}$ in Figure 9.

Having seen some examples, we consider the dynamical system a little more closely. Define the transition map $f : k^3 \rightarrow k^3$ by

$$f(x^1, x^2, x^3) = (x^2, x^3, 8 - x^1 - x^2 x^3).$$

By replacing each x_i with $y_i := x_i - 2$ (since 2 is a fixed point), we may replace Equation (7.2) with the homogeneous relation

$$(7.5) \quad y_i = -y_{i-3} - y_{i-2} y_{i-1} - 2(y_{i-2} + y_{i-1});$$

with the initial conditions $(y_1, y_2, y_3) = (-z, z, z^2)$, the result is a sequence of **reduced Farey polynomials** $\phi_{p/q} := \Phi_{p/q}^{\infty, \infty} - 2$ which we studied in [EMS23a] (in that paper, these polynomials were called $Q_{p/q}$). We were surprised to observe, through computing a few examples, that the alternating sum of the coefficients of these polynomials seem to be squares (Table 6)!

More precisely, we make the following conjecture.

Conjecture 7.6. *The reduced Farey polynomial $\phi_\alpha(z)$ is always of the form*

$$(-1)^{\epsilon(\alpha)} z^{k(\alpha)} R(\alpha)^2,$$

TABLE 6. $\sqrt{|\phi_{\text{fib}(q-1)/\text{fib}(q)}(-1)|}$ for small q .

q	$\sqrt{ \phi_{\text{fib}(q-1)/\text{fib}(q)}(-1) }$
1	1
2	1
3	1
4	0
5	1
6	1
7	1
8	2
9	3
10	5
11	17
12	88
13	1491
14	131225
15	195656563
16	25675032478184
17	5023488609594854052817
18	128978233205135262131911855731900891
19	647920685391665774371139137077100918906183250131690881763
20	835676652588773441434735633165055341787601058478245210146 00110051618636610092033658769403650

where $\epsilon : \hat{\mathbb{Q}} \rightarrow \{-1, +1\}$, $k : \hat{\mathbb{Q}} \rightarrow \mathbb{Z}/2\mathbb{Z}$, and $R : \hat{\mathbb{Q}} \rightarrow \mathbb{Z}[z]$ are functions which satisfy Farey recursion properties: if $\alpha, \beta \in \hat{\mathbb{Q}}$ are Farey neighbours, then

- (1) $\epsilon(\alpha \oplus \beta) = \epsilon(\alpha)\epsilon(\beta)$;
- (2) $k(\alpha \oplus \beta) = k(\alpha) + k(\beta)$; and
- (3) either $R(\alpha \oplus \beta) = R(\alpha)R(\beta) + R(\alpha \ominus \beta)$ or $R(\alpha \oplus \beta) = R(\alpha)R(\beta) - R(\alpha \ominus \beta)$ depending on the position of $\alpha \oplus \beta$ in $\hat{\mathbb{Q}}$. In the case of the arboreal geodesic $\text{fib}(p)/\text{fib}(p+1)$, the rule with a minus sign occurs iff $\text{fib}(p)$ is even; the ‘bad points’ in $\hat{\mathbb{Q}}$ where this minus sign rule occurs are shown in red in Figure 10).

The functions ϵ , k , and R satisfy the initial conditions in the following table:

α	$\epsilon(\alpha)$	$k(\alpha)$	$R(\alpha)$
0/1	-1	+1	1
1/1	+1	+1	1
1/2	+1	0	z

This conjecture is a more precise version of Conjecture 1 of [EMS23a], and (if the rule for R could be guessed) can probably be proved by an application of Farey induction applied to Equation (7.5) (compare with Lemma 5.3.12 and Remark 5.3.13 of [Aki+07]), but really the interest to us is the following observation: our initial recurrence relation is uniform on $\hat{\mathbb{Q}}$ (that is, the recurrence rule is the same no matter the position in the Farey graph), but from it there naturally arises a recurrence rule which depends on some *parity* structure in the graph. This is the *second* such recurrence relation which has appeared in this paper (the elliptic Farey polynomial recurrence also had parity), and roughly speaking in the geometric context of the Riley slices this parity is coming from the categorisation of two-bridge knots as either starting with an east-west twist or a north-south twist.




FIGURE 10.]

Points up to height 8 in the Farey graph coloured according to which recursion rule for the function R in Conjecture 7.6 is correct at that fraction: in green are points $\alpha \oplus \beta$ such that $R(\alpha \oplus \beta) = R(\alpha)R(\beta) + R(\alpha \ominus \beta)$, in red are points such that the rule $R(\alpha \oplus \beta) = R(\alpha)R(\beta) - R(\alpha \ominus \beta)$ holds, and in blue are the points such that $0 = R(\alpha \ominus \beta)$ (so both rules hold). The full picture may be viewed at the URL

<https://aelzenaar.github.io/farey/badpoints.ps>.

We end this section with a few open questions.

- (1) Do other parity-dependent recurrence relations on the Farey graph occur in nature?
- (2) Is there an analogue of the theory of Farey recursive functions with constant difference that is applicable to the elliptic Farey polynomial recurrence?
- (3) Is there a geometric proof for Theorem 4.7 (for instance, via some study of the lengths of the represented geodesics)?
- (4) As a stepping stone to (3), is there a more combinatorial reason (for instance, in terms of geodesics on the marked sphere) for parity to matter for the elliptic Farey polynomial recurrence? That is, take two adjacent triangles in the Farey graph; what is it about the arrangements of the corresponding geodesics which causes their lengths to depend on each other by either the odd relation or the even relation?
- (5) Is there a way of determining computationally which roots of $\Phi_{p/q}^{a,b} + 2$ correspond to the cusp points? Computation shows that the shape of the sets $(\Phi_{p/q}^{\infty,\infty})^{-1}(-2)$ as $q \rightarrow \infty$ seems to become quite regularly shaped (possibly with polygonal convex hull), with two extremal points corresponding to the cusps (see Figure 8 and Figure 9). If this can be made precise and shown to hold in general, then it would provide a method for finding cusps.
- (6) Can one give closed forms for the factors of $\phi_{p/q}^{\infty,\infty} = \Phi_{p/q}^{a,b} - 2$?
- (7) How quickly does the sequence of Farey polynomials coming from the continued fraction decomposition approximate the irrational pleating ray?

REFERENCES

- [AGM21] Hala Alaqad, Jianhua Gong, and Gaven J. Martin. *Chebyshev polynomials and inequalities for Kleinian groups*. 2021. arXiv: 2103.05696 [math.CV].
- [Aki+07] Hirotaka Akiyoshi, Makoto Sakuma, Masaaki Wada, and Yasushi Yamashita. *Punctured torus groups and 2-bridge knot groups I*. Lecture Notes in Mathematics 1909. Springer, 2007. ISBN: 9783540718079. DOI: 10.1007/978-3-540-71807-9.
- [Aki+21] Hirotaka Akiyoshi, Ken’ichi Ohshika, John Parker, Makoto Sakuma, and Han Yoshida. “Classification of non-free Kleinian groups generated by two parabolic transformations”. In: *Transactions of the American Mathematical Society* 374 (3 2021), pp. 1765–1814. DOI: 10.1090/tran/8246. arXiv: 2001.09564 [math.GT].
- [Bea83] Alan F. Beardon. *The geometry of discrete groups*. Graduate Texts in Mathematics 91. Springer-Verlag, 1983. ISBN: 0387907882. DOI: 10.1007/978-1-4612-1146-4.
- [BLS86] A. F. Beardon, J. Lehner, and M. Sheingorn. “Closed geodesics on a Riemann surface with application to the Markov spectrum”. In: *Transactions of the American Mathematical Society* 295.2 (1986), pp. 635–647.
- [Bow98] Brian H. Bowditch. “Markoff triples and quasifuchsian groups”. In: *Proceedings of the London Mathematical Society* 77.3 (1998), pp. 697–736. DOI: 10.1112/s0024611598000604.
- [BS87] Joan S. Birman and Caroline Series. “Dehn’s algorithm revisited, with applications to simple curves on surfaces”. In: *Combinatorial group theory and topology*. Ed. by Stephen M. Gersten and John R. Stallings. Annals of Mathematics Studies 111. Princeton University Press, 1987, pp. 451–478. ISBN: 0691084092.

- [BZ03] Gerhard Burde and Heiner Zieschang. *Knots*. 2nd ed. Studies in Mathematics 5. de Gruyter, 2003.
- [Che+20] Eric Chesebro, Cory Emlen, Kenton Ke, Denise Lafontaine, Kelly McKinnie, and Catherine Rigby. “Farey recursive functions”. In: *Involve* 41 (2020), pp. 439–461. DOI: 10.2140/involve.2021.14.439. arXiv: 2008.13303.
- [Che20] Eric Chesebro. “Farey recursion and the character varieties for 2-bridge knots”. In: *Characters in low-dimensional topology*. Ed. by Olivier Collin, Stefan Friedl, Cameron Gordon, Stephan Tillmann, and Liam Watson. Contemporary Mathematics 760. American Mathematical Society, 2020, pp. 9–34. ISBN: 9781470452094. arXiv: 1902.01968 [math.GT].
- [Elz21] Alex Elzenaar. *Riley slice computational package*. GitHub repository. 2021. URL: <https://github.com/aelzenaar/riley>.
- [Elz22] Alex Elzenaar. “Deformation spaces of Kleinian groups”. Master thesis. The University of Auckland, 2022. URL: https://aelzenaar.github.io/msc_thesis.pdf.
- [EMS22] Alex Elzenaar, Gaven J. Martin, and Jeroen Schillewaert. *The elliptic cousins of the Riley slice*. In preparation. 2022.
- [EMS23a] Alex Elzenaar, Gaven J. Martin, and Jeroen Schillewaert. “Approximations of the Riley slice”. In: *Expositiones Mathematicae* 41 (1 2023), pp. 20–54. DOI: 10.1016/j.exmath.2022.12.002. arXiv: 2111.03230 [math.GT].
- [EMS23b] Alex Elzenaar, Gaven J. Martin, and Jeroen Schillewaert. “Concrete one complex dimensional moduli spaces of hyperbolic manifolds and orbifolds”. In: *2021-22 MATRIX annals*. Springer, 2023. arXiv: 2204.11422 [math.GT]. To appear.
- [Gil06] Jane Gilman. “Informative words and discreteness”. In: *Combinatorial group theory, discrete groups, and number theory*. Ed. by Benjamin Fine, Anthony M. Gaglione, and Dennis Spellman. Vol. 421. American Mathematical Society, 2006, pp. 147–155. ISBN: 978-0-8218-3985-0. DOI: 10.1090/conm/421/08034.
- [GKP94] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete mathematics. A foundation for computer science*. 2nd ed. Addison-Wesley, 1994. ISBN: 0201558025.
- [GMM98] Fred W. Gehring, Colin Maclachlan, and Gaven J. Martin. “Two-generator arithmetic Kleinian groups II”. In: *Bulletin of the London Mathematical Society* 30.3 (1998), pp. 258–266. DOI: 10.1112/S0024609397004359.
- [HW60] Godfrey H. Hardy and Edward M. Wright. *An introduction to the theory of numbers*. 4th ed. Oxford University Press, 1960.
- [KS94] Linda Keen and Caroline Series. “The Riley slice of Schottky space”. In: *Proceedings of the London Mathematics Society* 3.1 (69 1994), pp. 72–90. DOI: 10.1112/plms/s3-69.1.72.
- [KS98] Yohei Komori and Caroline Series. “The Riley slice revisited”. In: *The Epstein birthday schrift*. Ed. by Igor Rivin, Colin Rourke, and Caroline Series. Geometry and Topology Monographs 1. Mathematical Sciences Publishers, 1998, pp. 303–316. DOI: 10.2140/gtm.1998.1.303. arXiv: math/9810194.
- [Mar16] Albert Marden. *Hyperbolic manifolds. An introduction in 2 and 3 dimensions*. 2nd ed. First edition was published under the title “Outer circles”. Cambridge University Press, 2016. ISBN: 9781107116740.

- [Mas87] Bernard Maskit. *Kleinian groups*. Grundlehren der mathematischen Wissenschaften 287. Springer-Verlag, 1987. ISBN: 978-3-642-61590-0. DOI: 10.1007/978-3-642-61590-0.
- [McM91] Curt McMullen. “Cusps are dense”. In: *Annals of Mathematics* 133 (1991), pp. 217–247. DOI: 10.2307/2944328.
- [Miy03] Hideki Miyachi. “Cusps in complex boundaries of one-dimensional Teichmüller space”. In: *Conformal Geometry and Dynamics* 7 (2003), pp. 103–151. DOI: 10.1090/S1088-4173-03-00065-1.
- [MM99] Colin Maclachlan and Gaven J. Martin. “2-generator arithmetic Kleinian groups”. In: *Journal für die Reine und Angewandte Mathematik* 511 (1999), pp. 95–117. DOI: 10.1515/crll.1999.511.95/html.
- [MPT15] Sara Maloni, Frédéric Palesi, and Ser Peow Tan. “On the character variety of the four-holed sphere”. In: *Groups, Geometry and Dynamics* 9 (2015), pp. 737–782. DOI: 10.4171/GGD/326.
- [MR03] Colin Maclachlan and Alan W. Reid. *The arithmetic of hyperbolic 3-manifolds*. Graduate Texts in Mathematics 219. Springer-Verlag, 2003. ISBN: 978-1-4757-6720-9. DOI: 10.1007/978-1-4757-6720-9.
- [MSW02] David Mumford, Caroline Series, and David Wright. *Indra’s pearls. The vision of Felix Klein*. Cambridge University Press, 2002. ISBN: 0521352533.
- [OM10] Ken’ichi Ohshika and Hideki Miyachi. “Uniform models for the closure of the Riley slice”. In: *Contemporary Mathematics* 510 (2010), pp. 249–306. DOI: 10.1090/conm/510/10029.
- [Ril72] Robert Riley. “Parabolic representations of knot groups, I”. In: *Proceedings of the London Mathematics Society*. 3rd ser. 24 (1972), pp. 217–242. DOI: 10.1112/plms/s3-24.2.217.
- [Ser85] Caroline Series. “The geometry of Markoff numbers”. In: *The Mathematical Intelligencer* 7.3 (1985), pp. 20–29. DOI: 10.1007/bf03025802.
- [Ser91] Caroline Series. “Geometric methods of symbolic coding”. In: *Ergodic theory, symbolic dynamics and hyperbolic spaces*. Ed. by Tim Bedford, Michael Keane, and Caroline Series. Oxford University Press, 1991, pp. 125–151. ISBN: 0198596855.
- [Sti83] John Stillwell. *Classical topology and combinatorial group theory*. 2nd ed. Graduate Texts in Mathematics 72. Springer, 1983. ISBN: 978-1-4612-4372-4. DOI: 10.1007/978-1-4612-4372-4.
- [Wri05] David Wright. “Searching for the cusp”. In: *Spaces of Kleinian groups*. Ed. by Yair N. Minsky, Makoto Sakuma, and Caroline Series. LMS Lecture Notes 329. Cambridge University Press, 2005, pp. 301–336. DOI: 10.1017/cbo9781139106993.016.
- [Zha10] Qingxiang Zhang. “Two elliptic generator Kleinian groups”. PhD thesis. Massey University, 2010. URL: <http://hdl.handle.net/10179/2044>.