

# MTH 1020 Week 12 tutorial

- ① Three worked integrals
- ② Integration

## Theorem (Archimedes)

The volume of the cone with height  $h$  and radius  $r$  is one-third the volume of the cylinder it lies in.

### Proof.

Slice the cone into discs.

By similar triangles, the disc at distance  $t$  from the tip of the cone has radius  $t/h \cdot r$ . Hence it has area  $\pi \left(\frac{tr}{h}\right)^2$ .

Therefore the volume of the cone is

$$\int_0^h \pi \left(\frac{tr}{h}\right)^2 dt = \frac{\pi r^2}{h^2} \int_0^h t^2 dt = \frac{\pi r^2}{h^2} \cdot \frac{1}{3} h^3 = \frac{\pi}{3} r^2 h.$$



## Integral 1

$$\int \frac{1}{1+x^2} dx$$

## Integral 1

$$\int \frac{1}{1+x^2} dx$$

**Method 1.** Look at it and immediately see, 'it is the derivative of arctan'.

## Integral 1

$$\int \frac{1}{1+x^2} dx$$

**Method 1.** Look at it and immediately see, 'it is the derivative of  $\arctan$ '.

**Method 2.**  $1+x^2$  looks like it might simplify using a trig identity: remember that  $\tan^2 + 1 = \sec^2$ .

## Integral 1

$$\int \frac{1}{1+x^2} dx$$

**Method 1.** Look at it and immediately see, 'it is the derivative of  $\arctan$ '.

**Method 2.**  $1+x^2$  looks like it might simplify using a trig identity:  
remember that  $\tan^2 + 1 = \sec^2$ .

Let  $x = \tan \theta$ , so  $dx = \sec^2 \theta d\theta$ .

## Integral 1

$$\int \frac{1}{1+x^2} dx$$

**Method 1.** Look at it and immediately see, 'it is the derivative of  $\arctan$ '.

**Method 2.**  $1+x^2$  looks like it might simplify using a trig identity: remember that  $\tan^2 + 1 = \sec^2$ .

Let  $x = \tan \theta$ , so  $dx = \sec^2 \theta d\theta$ . Then

$$\begin{aligned}\int \frac{1}{1+x^2} dx &= \int \frac{1}{1+\tan^2 \theta} \sec^2 \theta d\theta \\ &= \int \frac{\sec^2 \theta}{\sec^2 \theta} d\theta \\ &= \int d\theta = \theta = \arctan x.\end{aligned}$$

## Integral 2

$$\int \arcsin x \, dx$$



## Integral 2

$$\int \arcsin x \, dx$$

**Method 1.** Try substituting the only thing possible:  $\theta = \arcsin x$ . This means  $x = \sin \theta$  and so  $dx = \cos \theta \, d\theta$ .

## Integral 2

$$\int \arcsin x \, dx$$

**Method 1.** Try substituting the only thing possible:  $\theta = \arcsin x$ . This means  $x = \sin \theta$  and so  $dx = \cos \theta \, d\theta$ . The integral becomes

$$\int \arcsin x \, dx = \int \theta \cos \theta \, d\theta$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 1.** Try substituting the only thing possible:  $\theta = \arcsin x$ . This means  $x = \sin \theta$  and so  $dx = \cos \theta \, d\theta$ . The integral becomes

$$\int \arcsin x \, dx = \int \theta \cos \theta \, d\theta$$

This should look like an integration by parts thing. Take

$$u = \theta$$

$$dv = \cos \theta \, d\theta$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 1.** Try substituting the only thing possible:  $\theta = \arcsin x$ . This means  $x = \sin \theta$  and so  $dx = \cos \theta \, d\theta$ . The integral becomes

$$\int \arcsin x \, dx = \int \theta \cos \theta \, d\theta$$

This should look like an integration by parts thing. Take

$$\begin{aligned} u &= \theta & v &= \sin \theta \\ du &= d\theta & dv &= \cos \theta \, d\theta \end{aligned}$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 1.** Try substituting the only thing possible:  $\theta = \arcsin x$ . This means  $x = \sin \theta$  and so  $dx = \cos \theta \, d\theta$ . The integral becomes

$$\int \arcsin x \, dx = \int \theta \cos \theta \, d\theta$$

This should look like an integration by parts thing. Take

$$\begin{aligned} u &= \theta & v &= \sin \theta \\ du &= d\theta & dv &= \cos \theta \, d\theta \end{aligned}$$

and we end up with

$$\begin{aligned} \int \theta \cos \theta \, d\theta &= \theta \sin \theta - \int \sin \theta \, d\theta \\ &= \theta \sin \theta + \cos \theta \\ &= x \arcsin x + \cos \arcsin x \\ &= x \arcsin x + \sqrt{1 - x^2}. \end{aligned}$$

## Integral 2

$$\int \arcsin x \, dx$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 2.** Try integration by parts straight away: let

$$u = \arcsin x$$

$$dv = dx$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 2.** Try integration by parts straight away: let

$$\begin{aligned} u &= \arcsin x & v &= x \\ du &= (1 - x^2)^{-1/2} dx & dv &= dx \end{aligned}$$



## Integral 2

$$\int \arcsin x \, dx$$

**Method 2.** Try integration by parts straight away: let

$$\begin{aligned} u &= \arcsin x & v &= x \\ du &= (1 - x^2)^{-1/2} dx & dv &= dx \end{aligned}$$

so our integral becomes

$$\int \arcsin x \, dx = x \arcsin x - \int \frac{x}{\sqrt{1 - x^2}} \, dx. \quad (*)$$

## Integral 2

$$\int \arcsin x \, dx$$

**Method 2.** Try integration by parts straight away: let

$$\begin{aligned} u &= \arcsin x & v &= x \\ du &= (1 - x^2)^{-1/2} dx & dv &= dx \end{aligned}$$

so our integral becomes

$$\int \arcsin x \, dx = x \arcsin x - \int \frac{x}{\sqrt{1 - x^2}} \, dx. \quad (*)$$

Let  $w = 1 - x^2$ , so  $dw = -2x \, dx$  and

$$\begin{aligned} \int \frac{x}{\sqrt{1 - x^2}} \, dx &= \int \frac{\frac{1}{-2x} x}{\sqrt{w}} \, dw \\ &= -\frac{1}{2} \int \frac{1}{\sqrt{w}} \, dw \\ &= -\sqrt{w} = -\sqrt{1 - x^2}. \end{aligned}$$

## Integral 2

$$\int \arcsin x \, dx$$

....integrated by parts...

$$\int \arcsin x \, dx = x \arcsin x - \int \frac{x}{\sqrt{1-x^2}} \, dx. \quad (*)$$

Let  $w = 1 - x^2$ , so  $dw = -2x \, dx$  and

$$\begin{aligned} \int \frac{x}{\sqrt{1-x^2}} \, dx &= \int \frac{\frac{1}{-2x} x}{\sqrt{w}} \, dw \\ &= -\frac{1}{2} \int \frac{1}{\sqrt{w}} \, dw \\ &= -\sqrt{w} = -\sqrt{1-x^2}. \end{aligned}$$

## Integral 2

$$\int \arcsin x \, dx$$

....integrated by parts...

$$\int \arcsin x \, dx = x \arcsin x - \int \frac{x}{\sqrt{1-x^2}} \, dx. \quad (*)$$

Let  $w = 1 - x^2$ , so  $dw = -2x \, dx$  and

$$\begin{aligned} \int \frac{x}{\sqrt{1-x^2}} \, dx &= \int \frac{\frac{1}{-2x} x}{\sqrt{w}} \, dw \\ &= -\frac{1}{2} \int \frac{1}{\sqrt{w}} \, dw \\ &= -\sqrt{w} = -\sqrt{1-x^2}. \end{aligned}$$

Substituting this back into (\*) gives

$$\int \arcsin x \, dx = x \arcsin x + \sqrt{1-x^2}.$$

# Key concepts (integration)

- 1 Definite integral as a Riemann sum. Indefinite integral as anti-differentiation.
- 2 Fundamental theorem of calculus.
- 3 Area calculation.
- 4 Techniques of integration:
  - 1 Substitution (chain rule)
  - 2 By parts (product rule)
  - 3 By partial fractions
  - 4 By trig identities

## A special challenge integral

$$\int \sqrt{\tan x} \, dx$$

# MTH 1020 Week 12 tutorial

- 1 Get into groups of 3-4 people who all prepared a different question in advance.
- 2 Write your **preferred name** and **ID number** on the whiteboards so I can take attendance
- 3 Present your prepared question to each other as I come around, you should only take about 5min each for this.
- 4 Then get started on the other questions **in your groups**.
- 5 **At the end:** please erase the boards and return any markers etc that you used (you do not need to return the handouts)