

# Level Three Conic Sections

Alex Elzenaar

September 21, 2019



*O King, for traveling over the country, there are royal roads and roads for common citizens, but in geometry there is one road for all.*  
(Menaechmus)

# Contents

<b>1</b>	<b>Introduction and Historical Remarks</b>	<b>2</b>
<b>2</b>	<b>Basic results</b>	<b>3</b>
<b>3</b>	<b>Isometries</b>	<b>6</b>
<b>4</b>	<b>Cones</b>	<b>9</b>
<b>5</b>	<b>Orthogonal families of conics</b>	<b>12</b>
<b>6</b>	<b>Orbital mechanics</b>	<b>15</b>
<b>7</b>	<b>Unification</b>	<b>18</b>
<b>A</b>	<b>List of formulae</b>	<b>21</b>
<b>B</b>	<b>Further reading</b>	<b>22</b>

## Preface

The conic sections are the simplest curves in the plane which are not simply straight lines. However, most secondary school treatments of the subject tend to present a set of vaguely connected case-by-case results, rather than any kind of coherent story. These notes are my attempt to avoid this.


Most of the content is presented as a series of exercises; however, even an enthusiastic Y13 student will require a significant amount of guidance. As always, though, it is important that the student struggles with the material on their own!

Section 2 (basic results) contains almost all the ‘required’ material; section 3 (isometries) examines the role of transformations in geometry in a superficial way that tries to give some idea of the fundamental concepts which tie all of the different sub-fields of geometry together. Section 4 (cones) is a short description of the link between conic sections and cones, including an account of the ‘ice-cream proof’ which shows the geometric meaning of the foci of the ellipse in relation to the cone which it is a slice of. Section 5 (orthogonal families of curves) gives a few notes on some beautiful geometry which relates different families of conics together, and section 6 (applications to physics) derives Kepler’s laws of planetary motion using L3 calculus and the theory of conics. Finally, section 7 proves a rather pretty theorem which, in some sense, tells us everything special about the form of conic sections.

A short appendix is provided, which summarises the basic results.

## Prerequisites

These notes have perhaps the most ‘formal’ prerequisites out of all my Y13 notes. I will assume results from trigonometry, linear systems, calculus, and even algebra. However, this does not mean that the full power of these subjects are used. The most important prerequisites are actually Y11 and Y12 geometry because there are a number of results from there about triangles, circles, lines, and so forth that will be used here. My Y13 notes on trigonometry revise a number of these results, and so the reader is directed there initially.



---

## Section 1: Introduction and Historical Remarks

---

The study of the conic sections (as with much of elementary plane and solid geometry) goes back to the ancient Greeks. It is thought that the first person to study the slices of a cone was Menaechmus, who lived on the Gallipoli Peninsula in around 350 BCE, in his solution of the Delian problem: the construction of a cube whose volume is twice that of a given cube. Anecdotally, it is to Menaechmus that the famous quote on the cover of these notes is attributed (usually in his disputed role as the tutor to Alexander the Great).

Euclid of Alexandria published his book *Elements*, one of the most influential mathematical texts of all time, in around 300 BCE (for perspective, this is around thirty years after the death of Alexander the Great). In it, he set down the set of basic results which can be proved true about circles and lines on a plane: while modern geometry has gone much further than Euclid (in considering geometry in more than two dimensions, and on surfaces much more complicated than the plane), the basic results which are taught in any introductory geometry course are usually treated in the *Elements*.

As well as studying circles and lines, Euclid wrote a treatise on the conic sections. Unfortunately, it no longer survives; however, Apollonius of Perga (a city in what is now Turkey) expanded Euclid's work and published his *Conics* the century after Euclid's death. Apollonius used Euclidean geometry to prove most of the classical theorems on the conic sections, including the many applications to physics that the subject holds.

Despite these ancient roots, it turns out that the subject is deep enough to still hold interest in the modern day. Indeed, we will use techniques from calculus (in other words, from the 16th and 17th century) and from complex analysis (the 18th and 19th century) in these notes; and the proper setting for conics turns out to be projective geometry, which dates from the 19th and 20th century — although we will only touch on this briefly here.

Conic sections have many beautiful applications; later on, we will discuss applications to optics and to celestial mechanics. For example, in the 17th century the German astronomer Johannes Kepler showed that all astronomical bodies follow, due to gravity, paths in the shape of conic sections.

---

## Section 2: Basic results

---

**Definition.** Let  $O$  be a fixed point and  $\ell$  be some line not passing through  $O$ . A **conic**  $\mathcal{C}$  is the locus of a point  $P$  such that, if  $K$  is the point on  $\ell$  so that  $\ell \perp PK$ ,


$$\varepsilon = \frac{|OP|}{|PK|}$$

where  $\varepsilon$  is some fixed non-negative constant.

The point  $O$  is called the **focus**, the line  $\ell$  the **directrix**, and the constant  $\varepsilon$  the **eccentricity**. The **latus rectum** (Latin: straight side) is defined to be the chord of a conic through the focus and parallel to the directrix; let  $l$  be the length of the latus rectum.

The conic is variously called:

- an **ellipse** if  $\varepsilon < 1$ ;
- a **parabola** if  $\varepsilon = 1$ ; and
- an **hyperbola** if  $\varepsilon > 1$ .



### Exercise 1: Polar form.

Show that, if  $X$  is the point on the directrix  $\ell$  of a conic such that  $OX \perp \ell$ , then the polar equation of the conic with respect to the axis  $OX$  and origin  $O$  is

$$\frac{l}{r} = 1 + \varepsilon \cos \theta.$$

Conclude that:

1. Every conic is symmetric with respect to  $OX$ . (A curve is symmetric with respect to a line if it is unchanged after reflection across that line.)
2. The ellipse is a closed and bounded curve. (A curve is closed if when one walks along it, one eventually returns to one's starting place. A curve is bounded if there is some large square containing the entire curve.)
3. The parabola is unbounded, but is connected. (A curve is connected if, whenever  $P$  and  $Q$  are two points on the curve, one can walk from  $P$  to  $Q$  without leaving the curve.)
4. The hyperbola consists of two separate branches, each extending to infinity, given by  $-\alpha < \theta < \alpha$  and  $\alpha < \theta < 2\pi - \alpha$  (where  $\alpha = \operatorname{arcsec}(-\varepsilon)$ ).

### Exercise 2: Rectangular form.

By squaring the polar form equation, show that the Cartesian equation for a conic, taking suitable axes and origin, is

$$x^2 + y^2 = (l - \varepsilon x)^2.$$

1. If  $\varepsilon \neq 1$ , and a suitable origin is chosen, show that the conic equation can be written in the form

$$\frac{x^2}{a^2} \pm \frac{y^2}{b^2} = 1$$

for some real numbers  $a$  and  $b$ . The new location of the origin is called the **centre** of the conic. Conclude that:

- (a) Both the ellipse and hyperbola are symmetric across both Cartesian axes.
- (b) For an ellipse, the values  $2a$  and  $2b$  are the lengths of the chords through the origin along the  $x$ - and  $y$ -axes respectively. (The longer of these is called the **major axis**, and the shorter the **minor axis**.)
- (c) The two branches of a hyperbola lie in opposite regions bounded by the two lines (**asymptotes**)

$$\left(\frac{x}{a} - \frac{y}{b}\right)\left(\frac{x}{a} + \frac{y}{b}\right) = 0.$$

The value  $2a$  is the length of the transverse axis of the hyperbola and the value  $2b$  is the length of the conjugate axis: the two dimensions of the rectangle whose diagonals are the asymptotes and which is bounded by the hyperbola.

- (d) Further, show that as  $x \rightarrow \pm\infty$ , the branches of the hyperbola tend arbitrarily close to the asymptotes. (Hint: take limits.)
2. If  $\varepsilon = 1$ , show that the conic equation for the parabola can be written in the form

$$y^2 = 2l\left(\frac{1}{2}l - x\right).$$

By reflecting in a suitable vertical line, derive the standard form equation

$$y^2 = 2lx.$$

Given this latter equation, give the coordinates of the focus, and the Cartesian equation of the directrix.

### Exercise 3: Parametric form.

Show that:

- 1. The ellipse  $x^2/a^2 + y^2/b^2 = 1$  is parameterised by  $(a \cos t, b \sin t)$  for  $0 \leq t < 2\pi$ .
- 2. The parabola  $y^2 = 2lx$  is parameterised by  $(2lt^2, 2lt)$ .
- 3. The hyperbola  $x^2/a^2 - y^2/b^2 = 1$  is parameterised by  $(a \sec t, b \tan t)$  for  $0 \leq t < 2\pi$ .

Illustrate these parameterisations with a suitable computer program (e.g. Geogebra, or your favourite programming language).

### Exercise 4: Focii of rectangular conics.

Given the equation

$$\frac{x^2}{a^2} \pm \frac{y^2}{b^2} = 1,$$

compute the coordinates of the focus, equation of the directrix, and eccentricity of the conic. Hence

show that ellipses and hyperbolae have two foci and two directrices each. Where is the directrix of a circle?

Show that the following two properties can be used as alternative definitions for the ellipse and hyperbola:

AP1. Show that the ellipse with foci  $O_1$  and  $O_2$  is the locus of all points  $P$  such that  $d(O_1, P) + d(P, O_2) = R$  for some constant  $R$ . Give  $R$  in terms of  $a$  and  $b$ .

AP2. Show that the hyperbola with foci  $O_1$  and  $O_2$  is the locus of all points  $P$  such that  $d(O_1, P) - d(P, O_2) = R$  for some constant  $R$ . Give  $R$  in terms of  $a$  and  $b$ .

As an application of this definition, we will derive the ‘reflection property’ of the ellipse.

1. Suppose  $O_1$  and  $O_2$  are on the same side of a line  $\ell$ . Show that the shortest path from  $O_1$  to  $O_2$  which touches the line  $\ell$  is the broken line  $O_1XO_2$ , where  $X$  is on  $\ell$  and the angles between  $O_1X$  and  $\ell$  and between  $XO_2$  and  $\ell$  are equal. (Hint: you can do this with calculus, but that would be like killing a fly with a sledgehammer.)
2. Show that, if  $O_1$  and  $O_2$  are two foci of an ellipse and  $X$  is any point on the ellipse, then the broken line  $O_1XO_2$  makes equal angles with the tangent line of the ellipse at  $X$ . (Hint: show that this broken line is the shortest path from  $O_1$  to  $O_2$  that touches the tangent line at  $X$ .)
3. Thus, given the law of reflection for waves, show that if a light source is placed at  $O_1$  then every ray from the source will arrive at  $O_2$  at precisely the same time.
4. Derive some result of this kind for parabolas by treating a parabola as an ellipse with one focus ‘at infinity’. Suggest an application of this property related to, say, torches.

*Remark.* There is at least one interesting generalisation of the above exercise, known as the ‘elliptical billiards problem’: it is possible to show that the lines of the trajectory of a billiard ball on an elliptical table (or, more generally, a table with a conic boundary) are all tangent to a conic with the same foci as the table.

For a precise statement of the result, and a proof, see chapter 13 and appendix E of Leopold Flatto, *Poncelet’s theorem*. AMS (2009).

---

## Section 3: Isometries

---

In the previous section, we saw geometrically that every ellipse or hyperbola is just a translated and rotated version of the graph of an equation with the form

$$(1) \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$$

(Herein, in this section we will write  $p = 1/a^2$  and  $q = 1/b^2$  for convenience.)

Here, we will show this relation using coordinates — the advantage of this approach is that it allows us to calculate precisely from the equation of a conic its position and angle with respect to the standard axes. The subtle point here is basically that the same curve will have different equations depending on its position in the coordinate system — in the previous section we chose our coordinate system in the ‘nicest way possible’ by geometric means, and now we are given a conic already sitting in some coordinate system which we want to understand.

In order to pursue this programme, we need to study the effect of rotations and translations on coordinate systems. We begin with rotations because it turns out that rotating before translating is easier for our purposes.

Let  $\rho = \text{cis } \theta$  be a complex number such that  $|\rho| = 1$ . If  $z = x + yi$  is a point on the complex plane, then  $\rho$  acts by multiplication on  $z$  to rotate it around the origin by an angle  $\theta$ ; we can then calculate the resulting rectangular coordinates of  $\rho z$  to see how a rotation affects our normal coordinate system.

$$\begin{aligned} z &= |z| \text{cis}(\tan^{-1} x/y) \\ \rho z &= |z| \text{cis}(\tan^{-1} x/y + \theta) \\ &= |z| \left[ \cos(\tan^{-1} x/y + \theta) + i \sin(\tan^{-1} x/y + \theta) \right]. \end{aligned}$$

Using trig identities, we calculate

$$\begin{aligned} \cos(\tan^{-1} x/y + \theta) &= \cos(\tan^{-1} x/y) \cos \theta - \sin(\tan^{-1} x/y) \sin \theta \\ &= \frac{x}{\sqrt{x^2 + y^2}} \cos \theta - \frac{y}{\sqrt{x^2 + y^2}} \sin \theta \\ \sin(\tan^{-1} x/y + \theta) &= \sin(\tan^{-1} x/y) \cos \theta + \cos(\tan^{-1} x/y) \sin \theta \\ &= \frac{y}{\sqrt{x^2 + y^2}} \cos \theta + \frac{x}{\sqrt{x^2 + y^2}} \sin \theta \end{aligned}$$

and so

$$\begin{aligned} \rho z &= |z| \left[ \cos(\tan^{-1} x/y + \theta) + i \sin(\tan^{-1} x/y + \theta) \right] \\ &= (x \cos \theta - y \sin \theta) + i(y \cos \theta + x \sin \theta). \end{aligned}$$

In other words, if the point  $(x, y)$  is rotated about the origin by an angle  $\theta$ , then

$$(2) \quad (x, y) \xrightarrow{\rho} (x \cos \theta - y \sin \theta, y \cos \theta + x \sin \theta).$$

Considering  $ax^2 + bx + cxy + dy + ey^2$ , then, we want to rotate  $(x, y)$  by some  $\theta$  so that some terms vanish. Doing a long computation, we find that

$$\begin{aligned} ax^2 + bx + cxy + dx + ey^2 &\mapsto a(x \cos \theta - y \sin \theta)^2 + b(x \cos \theta - y \sin \theta) \\ &\quad + c(x \cos \theta - y \sin \theta)(y \cos \theta + x \sin \theta) \\ &\quad + d(y \cos \theta + x \sin \theta) + e(y \cos \theta + x \sin \theta)^2 \\ &= x^2(a \cos^2 \theta + \frac{c}{2} \sin 2\theta + e \sin^2 \theta) + x(b \cos \theta + d \sin \theta) \\ &\quad + xy((e - a) \sin 2\theta + c \cos 2\theta) \\ &\quad + y(-b \sin \theta + d \cos \theta) + y^2(a \sin^2 \theta - \frac{c}{2} \sin 2\theta + e \cos^2 \theta) \end{aligned}$$

and, looking at this, we see that an easy candidate we can try to get rid of is the  $xy$  term. In fact, if we want this term to be zero we need only solve

$$(3) \quad (e - a) \sin 2\theta + c \cos 2\theta = 0$$

which is easy:  $\frac{-c}{e-a} = \frac{\sin 2\theta}{\cos 2\theta} = \tan 2\theta$ , and so in order to remove the  $xy$  term we need only rotate our coordinate system by

$$(4) \quad \theta = \frac{1}{2} \tan^{-1} \frac{c}{a-e}.$$

Making the coordinate system change  $(x, y) \mapsto (x \cos \theta - y \sin \theta, y \cos \theta + x \sin \theta)$  therefore leaves us with something that looks like  $ax^2 + bx + dy + ey^2 = 1$ , where  $x$  and  $y$  are now coordinates in our rotated coordinate system and where the constants  $a$  to  $e$  are not necessarily the same as before.

We already know that if  $y = f(x)$  is graphed, then we can shift the graph up by  $x_0$  and to the right by  $y_0$  by suitable transformations of the coordinates:  $y - y_0 = f(x - x_0)$  has the shifted graph. Last year, we performed similar transformations on parabolas by completing the square, and so this is the technique which we will use now. We proceed as we did last year (but now completing the square in both  $x$  and  $y$ ):

$$ax^2 + bx + dy + ey^2 = a \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a} + e \left( \frac{d}{2e} + y \right)^2 - \frac{d^2}{4e} = 1.$$

If we now let  $(x, y) \mapsto \left( x - \frac{b}{2a}, y - \frac{d}{2e} \right)$ , then we have got a coordinate transformation that removes all the linear terms and leaves us with something like

$$ax^2 + ey^2 = 1 + \frac{b^2}{4a} + \frac{d^2}{4e}$$

and upon division of both sides by  $1 + \frac{b^2}{4a} + \frac{d^2}{4e}$  we end up, as promised, with something of the form  $px^2 + qy^2 = 1$ . Note that our coordinate system has completely changed: the new curve has the same shape as our original curve, but sits inside our new coordinate system much more naturally than it sat in our old coordinate system.

### Exercise 5: Further thoughts on this argument.

You are asked to think critically about the above argument (so reread it and check the computations) and to extend it (so not only must you read it, but understand it).

1. Clearly  $y = x^2$  cannot be written in the canonical form. Where does the proof above fall down for a parabola?
2. What happens if our ellipse is a circle?
3. We have now classified the quadratic equations in two variables corresponding to hyperbolae and ellipses ( $px^2 \pm qy^2 = 1$ ) and parabolas ( $y^2 = kx^2$ ). What other forms can the graph of such a quadratic equation take? (Hint: you should be able to find three other kinds, and show that every quadratic equation is of one of the six kinds.)

### Exercise 6: Computations.

First, read the following quote:

The attitude adopted in this book is that while we expect to get numbers out of the machine, we also expect to take action based on them, and, therefore we need to understand thoroughly what numbers may, or may not, mean. To cite the author's favorite motto,  
 "The purpose of computing is insight, not numbers," although some people claim,  
 "The purpose of computing numbers is not yet in sight."  
 There is an innate risk in computing because "to compute is to sample, and one then enters the domain of statistics with all its uncertainties."

— Richard W. Hamming, *Introduction to applied numerical analysis*, McGraw-Hill 1971, p.31.  
 (Quoted in <https://mathoverflow.net/a/7164>.)

1. Classify the following conics:

(a)  $\frac{10}{4}x^2 + 3xy + \frac{10}{4}y^2 = 1$

(b)  $4x^2 + 24xy + 11y^2 = 5$



2. What happens to the equation for the hyperbola  $x^2 - y^2 = a^2$  upon rotation by  $\pi/4$ ?
3. Give an equation for the parabola with vertex  $(0, 2)$ , and with axis  $y = x + 2$  (opening into the positive  $x$ -half plane).
4. Find the two foci of an ellipse passing through  $(0, 0)$ ,  $(0, 1)$ , and  $(1, 2)$ .

*Remark.* Felix Klein (a German mathematician who studied, among other things, geometric transformations) defined a geometry as a set of points together with a set of transformations of those points, such that two figures (two collections of points) are said to be ‘equivalent’ if there exists some invertible transformation mapping the figures onto each other.

For example, in normal Euclidean geometry (our current playground), we say two figures are ‘congruent’ if they can be placed on top of each other to exactly cover each other: in other words (and more precisely), two figures are congruent if one can be mapped to the other via a series of translations, rotations, and reflections — precisely the case which we have studied in this section. (Such transformations are called *isometries*.) Since all these maps preserve lengths and angles, these things are well-defined in Euclidean geometry.


A less familiar geometry is affine geometry, in which we also allow dilations (scaling) — so all circles are equivalent in affine geometry. This does mean that lengths become meaningless: any line segment can be mapped onto any other line segment, no matter its length. Angles also become meaningless (why?). However, enough properties are left to provide a rich venue for geometry: for example, parallel lines are still parallel after an affine transformation, and so any result which only uses parallelism in its proof makes sense here.

Other geometries include Möbius geometry, where figures are equivalent if they can be mapped onto each other by an isometry or a circle inversion (a reflection through a circle), and projective geometry, which is a further loosening of the conditions on an affine geometry such that even parallel lines make no sense and the only results left are those which just involve incidence and intersection. Perhaps surprisingly, projective geometry is in some sense the most rich geometry of all. This geometry is, in fact, the setting for section 7 of these notes.

---

## Section 4: Cones

---



We now move from the planar definitions of the conics to the ‘cone-slicing’ definitions. Essentially, our goal is to show that the conics are just slices of a cone (hence the name), as in the diagram above.<sup>1</sup>

### Exercise 7: Defining a cone.

We will define a cone to be the locus of all lines in three dimensional space which pass through the origin and through a point on some fixed curve in the plane  $z = 1$ . Note: to show a given object is a cone, one must guess a good defining curve (usually by intersecting it with the given plane) and then do two things: (N1) show that every line through the origin and a point on the guessed curve lies on the object, and (N2) show that every point on the object lies on such a line.

1. Give various non-examples of cones, especially surfaces satisfying criteria N1 but not N2 and those satisfying N2 but not N1.
2. Show that  $x^2 + y^2 = cz^2$  is a cone whose defining curve is a circle; show that the line through the origin and the centre of this circle cuts the  $z = 1$  plane at right angles. (This cone is called a **right circular cone**.)
3. Show that  $ax^2 + by^2 - cz^2 = 0$  is a cone if  $c > 0$  and  $a$  and  $b$  are real numbers. What is its defining curve?

### Exercise 8: Algebraic cone-slicing.

Show that all six kinds of conic sections which you found in exercise 5 are indeed obtained by slicing cones with planes.

*Hints:*

- Fix a cone, say  $x^2 + y^2 = cz^2$ , and then slice it with various planes.
- The plane through  $(x_0, y_0, z_0)$  which is orthogonal to the line between  $(0, 0, 0)$  and  $(a, b, c)$  is given by

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

If you use this fact, you need not prove it.

### Exercise 9: Geometric cone-slicing: the ellipse.


In the previous exercise, we had to use coordinates to prove that every slice of a cone is a conic. We will now discover a geometric proof that an ellipse is just a slice of a right circular cone, in such a way that we will understand why such slices obey the focii properties of the ellipse.

Let us take our plane and slice our cone with it. Pick any point  $P$  in the intersection of the two surfaces (i.e. on the curve we wish to show is an ellipse). In addition, we will place two spheres  $S_1$  and  $S_2$  into our cone such that they are tangent to both the cone and to the plane. Let  $F_1$  and  $F_2$  be the

---

<sup>1</sup>By Pbroks13 - Own work, CC BY 3.0, <https://commons.wikimedia.org/w/index.php?curid=5919064>

two points of contact between the plane and the spheres, and let  $C_1$  and  $C_2$  be the circles of contact between the cone and the spheres. The setup is illustrated below.<sup>a</sup>



1. Show that if  $X$  is a point outside some sphere, and if  $M$  and  $N$  are points on the sphere such that  $XM$  and  $XN$  are tangent lines to the sphere, then  $|XM| = |XN|$ . (Hint: consider the cross-section of the sphere obtained by slicing it with the plane of the triangle  $XMN$ .)
2. Suppose the vertex of our cone is  $O$ . Show that if  $A_1$  and  $A_2$  are the points of intersection between the line  $OP$  and the two circles  $C_1$  and  $C_2$  then  $|PF_1| = |PA_1|$  and  $|PF_2| = |PA_2|$ .
3. Show that  $|PA_1| + |PA_2|$  is independent of the point  $P$ .
4. Conclude that  $|PF_1| + |PF_2|$  is constant for every value of  $P$ , and therefore that the locus of  $P$  is an ellipse with foci  $F_1$  and  $F_2$ .

<sup>a</sup> Tom M. Apostol, *Linear Algebra: A first course* (pp.78–9). John Wiley & Sons (1997).

### Exercise 10: Geometric cone-slicing: other conics.

The method of proof in the previous exercise is called the **method of Dandelin spheres**, after the Belgian mathematician Germinal Dandelin.



By Unknown - [www-history.mcs.st-andrews.ac.uk](http://www-history.mcs.st-andrews.ac.uk), Public Domain,  
<https://commons.wikimedia.org/w/index.php?curid=958124>

1. Adapt the proof in the previous exercise to show that hyperbolae and parabolae are slices of right circular cones.
2. Show that the degenerate conics are also slices of right circular cones.

3. Let us construct the Dandelin spheres for a conic lying in a plane  $\mathcal{P}$ . Pick one of the two spheres arbitrarily; call the circle of tangency of this sphere to the cone  $\mathcal{Q}$ . Show that  $\mathcal{P}$  and  $\mathcal{Q}$  intersect in a line, and that this line is a directrix of the conic.

For additional details, references, and diagrams, see Berger §17.3.

---

## Section 5: Orthogonal families of conics

---

In this section, we will consider families of conics with the same foci, and we will find all the curves which are orthogonal to each family.

**Definition.** Two curves are said to be **orthogonal** if, at every point where they intersect, they have tangent lines which meet at right angles. Equivalently, two curves are orthogonal if at each point of intersection the normal line of one is the tangent line of the other.

Let us first revise a result which you should now have proved twice (both in Y12 coordinate geometry and in Y13 calculus).

### Exercise 11: Orthogonality.

Show that, if  $y = f(x)$  is a simple curve (i.e. a curve which does not cross itself) that is differentiable at  $P = (x_0, y_0)$ , then there is a unique line  $N$  passing through  $P$  that is orthogonal to the curve at  $P$ ; further, the slope of  $N$  is  $-\frac{1}{f'(x_0)}$ .

If the reader is feeling clever, she should prove it in two different ways. (Hint: you can prove it via drawing triangles — this is the proof given in the Y13 calculus notes — or by rotating your line.)

*Remark.* In the exercise above, we implicitly assumed in the statement that our curve was the graph of a function; however, this need not be the case. We only need the curve to be a function locally (i.e. for there to exist a small ball about  $(x_0, y_0)$  within which our curve is the graph of a function). You should check that your proof only uses local properties of the graph of  $f$ !

### Exercise 12: Homogenous differential equations.

Suppose  $y$  is a function of  $x$ . Recall that a differential equation is said to be **separable** if it can be written in the form  $\frac{dy}{dx} = P(x)Q(y)$ , where  $P$  and  $Q$  are functions of  $x$  and  $y$  alone.

1. Show, using the chain rule (i.e. without treating  $\frac{dy}{dx}$  as a fraction), that  $\int \frac{1}{Q(y)} dy = \int P(x) dx$ .
2. If a curve with derivative  $\frac{dy}{dx} = \frac{\sin x}{\cos y}$  passes through  $(0, 0)$ , which  $y$  values does it take when  $x = \pi$ ?
3. Find an explicit formula for a function  $f$  defined for all positive numbers such that when  $y = f(x)$ ,  $\frac{dy}{dx} = \frac{y^2}{1-x}$  and  $f(1) = 0$ .

The differential equations we wish to consider later in this section are, unfortunately, *not* separable. However, they are nice enough that we can apply a change of variables to *make* them separable.

We say that a function of two variables  $P(x, y)$  is **homogenous of degree  $n$**  if  $P(tx, ty) = t^n P(x, y)$  for all  $x$  and  $y$ .

4. Suppose that  $P(x, y)$  and  $Q(x, y)$  are both homogenous of degree  $n$ . Show that  $f(x, y) = \frac{P(x, y)}{Q(x, y)}$  is homogenous of degree 0.
5. Let us consider the equation  $\frac{dy}{dx} = f(x, y)$ , where  $f$  is as in the previous question. Let  $t = 1/x$ ; show that, using the substitution  $z = y/x$  and homogeneity, we obtain the separable equation

$$z + x \frac{dz}{dx} = f(1, z) = \frac{P(1, z)}{Q(1, z)}.$$

Find the solution of this equation.

6. Using this technique, solve  $\frac{dy}{dx} = \frac{x^2 - 2y^2}{xy}$ . (Be sure to check that the technique is applicable: i.e. that the right hand side is homogenous of degree zero.)

In a break from form, I will do a nice easy example to start with. Suppose we consider the family of curves consisting of all the circles whose centre is the origin; it should be intuitively clear that the only curves orthogonal to all of these circles at once are the lines through the origin.

As a first step, we know that every such circle has equation  $x^2 + y^2 = R^2$  for some constant  $R$ . Given any particular circle, its derivative is given by  $2x + 2y \frac{dy}{dx} = 0$  or  $\frac{dy}{dx} = -\frac{x}{y}$ ; since the parameter  $R$  is absent from this equation, it describes every such circle. (We have shown that every such circle has this derivative, but you should check yourself, by integrating, that every equation of this form is a circle centred at the origin.)

We want to find all the curves such that, at every point  $(x, y)$  on our curve, the tangent line of our curve is the normal line of the circle centred at the origin and passing through that point. To do this, we simply need to solve

$$\frac{dy}{dx} = \frac{y}{x}$$

and this is not only separable but easy:  $\int \frac{dy}{y} = \int \frac{dx}{x}$  and thus  $\ln|x| = \ln|y| + C$ , or  $x = Ky$ ; and this is just the family of all lines through the origin, as we guessed initially.

For convenience, if  $\mathcal{F}$  is a family of curves then we will call the family of curves orthogonal to  $\mathcal{F}$  the **orthogonal complement** of  $\mathcal{F}$ .

### Exercise 13: Orthogonal families of curves.

We (or rather you, the reader) will now show that when we take orthogonal complements of various families of conics, then we obtain other families of conics. As well as the results explicitly stated here, you should consider how the families are related to each other. Show that:

1. The family of NE-SW diagonal hyperbolae,  $xy = c$ , are orthogonal to the family of NW-SE diagonal hyperbolae,  $xy = -c$ .
2. The vertical parabolae,  $y = ax^2$ , are orthogonal to the family of ellipses  $x^2 + \frac{y^2}{b} = 1$ . (See `parabola-and-ellipse.ggb`.)
3. The family of all circles tangent to the  $y$ -axis at the origin (i.e.  $x^2 + y^2 = 2rx$ ) is orthogonal to the family of all circles tangent to the  $x$ -axis at the origin (i.e.  $x^2 + y^2 = 2ry$ ).

Hence, complete the following: orthogonal complements of  $X$  are  $Y$ , taking  $X$  to be hyperbolae, parabolae, circles, and ellipses. (We have found the orthogonal complements of families of curves in particular places, but consider the hint after exercise 14.)

Finally, we have one rather pretty example.

### Exercise 14: Circles of Apollonius.

Fix two points  $A$  and  $B$ . Let us consider the locus of all points  $P$  such that  $\frac{|AP|}{|BP|} = \rho$ , for positive numbers  $\rho$ ; we will examine one particular perpendicular family.

1. Show that if we fix  $A$  and  $B$  and vary  $\rho$  then we have a family of circles whose centres lie on the line  $AB$ .
2. Show that the orthogonal complement of this family is the set of circles passing through  $A$  and  $B$  with centres on the perpendicular bisector of the segment  $AB$ .
3. Use Geogebra (or similar software) to draw a nice picture.

Hint: without loss of generality, you can suppose  $A = (-1, 0)$  and  $B = (0, 1)$ . (Then the result follows for any choice of  $A$  and  $B$ , simply because dilations, translations, and rotations all preserve circles and orthogonality of lines.)

There are three ways to do this exercise that the author is aware of: perhaps the easiest for us now is to apply calculus like we have been doing throughout this section. For a geometric view on this exercise, see Coxeter §6.6; for a view from complex calculus, see Fisher §1.2.

### Exercise 15: Bipolar coordinates (difficult).

(Prerequisite: you do need to understand exercise 14 quite well. You should also read the epilogue on hyperbolic trigonometric functions in the trigonometry notes.)

From your pictures in the previous exercise, you should notice that every point in the plane is at the

intersection of two circles of Apollonius. Let us fix  $A = (-1, 0)$  and  $B = (1, 0)$ , as in the hint for the previous exercise; we will use these circles to define a new coordinate system for the plane.

Suppose  $P = (x, y)$  lies at the intersection of two circles; then we define  $\sigma$  to be the angle  $APB$ , and  $\tau = \ln \left| \frac{PA}{PB} \right|$ . We then define the **bipolar coordinates** of  $P$  to be  $(\sigma, \tau)$ .

1. Show that

$$\begin{cases} x = \frac{\sinh \tau}{\cosh \tau - \cos \sigma} \\ y = \frac{\sin \sigma}{\cosh \tau - \cos \sigma} \end{cases}.$$

2. Identifying the plane  $\mathbb{R}^2$  with the complex numbers  $\mathbb{C}$ , show that this is equivalent to  $x + iy = ai \cot \left( \frac{\sigma + i\tau}{2} \right)$ . (Recall that we define  $\cot z = i \frac{e^{iz} + e^{-iz}}{e^{iz} - e^{-iz}}$ .)
3. Identify the locus of all  $P = (\sigma, \tau)$  such that (a)  $\sigma$  is held constant, or (b)  $\tau$  is held constant.
4. Translate the standard equations of the conics into bipolar coordinates.

---

## Section 6: Orbital mechanics

---

We will now look briefly at some applications to the theory of gravity. Recall from stage 2 physics (or earlier) the following definitions and results:

**Definition.**

1. If an object's position  $x$  is given as a function of time  $t$ , then  $v := \frac{dx}{dt}$  is known as the velocity of the object and  $a := \frac{dv}{dt} = \frac{d^2x}{dt^2}$  is known as the acceleration of the object.
2. Newton's second law of motion: An object with mass  $m$ , when acted upon by a force  $F$ , undergoes an acceleration  $a = F/m$  in the same direction as the force; the force does not affect the motion of the object in any other direction.
3. Newton's law of gravity: If two objects have masses  $M$  and  $m$  and their centres are a distance  $R$  apart, then they impose on each other equal and opposite forces of magnitude  $\frac{GMm}{R^2}$ , where  $G$  is a constant.
4. If an object of mass  $m$  is close to the surface of the earth, then the gravitational force upon it due to the earth is  $mg$  towards the centre of the planet, where  $g \approx 9.81$ .

As a warmup, we will derive some simple results from stage 2 physics.

### Exercise 16: Projectile motion.

An object of small mass  $m$  flying through the air can often be modelled as a **projectile**: a particle with constant speed  $v_x = \frac{dx}{dt}$  along the ground, and constant acceleration  $a_y = mg$  pointing towards the centre of the earth.

1. Show that the motion of the mass is a parabola.
2. If the object is launched skywards at an angle of  $\pi/4$  to the horizontal, with a velocity  $v = 10$  along its direction of travel, how far along the ground will it travel?
3. With the conditions of problem (2), how is the focus of the parabola traced positioned?
4. At what angle must the object be launched for the focus to lie precisely on the ground?

So, to a first approximation, objects follow a parabolic path due to gravity. We made two main simplifications here:

1. We assumed that the force of gravity on the object was constant (in both magnitude and direction) regardless of position.
2. We assumed that the earth was fixed in space relative to the object.

Because of the positioning of our object, these approximations are actually quite valid: the force of gravity is roughly perpendicular to the surface of the earth, the surface of the earth is approximately flat when you are close to it, and the object was far enough away from the centre of the earth for a small increase or decrease in height to have a negligible impact on the magnitude force felt by the object. Further, the earth is massive compared to our object and so would feel a negligible force due to it.

We will now relax the first of these two approximations, and consider the new path followed.<sup>2</sup>

In order to make our discussion more concrete, we will fix some large mass  $M$  at the origin  $O$ , and we will consider some smaller planet, located at  $\mathbf{r}$  with mass  $m$ . The planet  $P$  will accelerate due to the gravitational force exerted on it by  $O$ ; we want to study the motion of  $P$ . It is convenient to use polar coordinates.

**Exercise 17: Kepler's second law.** 1. Suppose that  $\mathbf{u}_r = (\cos \theta, \sin \theta)$  is the unit vector in the direction of  $\mathbf{r}$ , so  $\mathbf{r} = r\mathbf{u}_r$ . Let us define  $\mathbf{u}_\theta = (-\sin \theta, \cos \theta)$  to be the unit vector perpendicular to  $\mathbf{u}_r$  in the direction of increasing  $\theta$ . Show that  $\frac{d\mathbf{u}_r}{d\theta} = \mathbf{u}_\theta$  and  $\frac{d\mathbf{u}_\theta}{d\theta} = -\mathbf{u}_r$ , and hence (using the

---

<sup>2</sup>For a detailed exposition of this derivation, see Simmons §21.



product and chain rules) deduce that:

$$\begin{aligned}\mathbf{v} &= \frac{d\mathbf{r}}{dt} = r \frac{d\theta}{dt} \mathbf{u}_\theta + \frac{dr}{dt} \mathbf{u}_r \\ \mathbf{a} &= \frac{d\mathbf{v}}{dt} = \left( 2 \frac{dr}{dt} \frac{d\theta}{dt} + r \frac{d^2\theta}{dt^2} \right) \mathbf{u}_\theta + \left( \frac{d^2r}{dt^2} - r \left( \frac{d\theta}{dt} \right)^2 \right) \mathbf{u}_r.\end{aligned}$$

2. Hence, using Newton's second law of motion and (a) above, we can write expressions for the components of the force  $\mathbf{F} = F_\theta \mathbf{u}_\theta + F_r \mathbf{u}_r$  acting on an object of mass  $m$ :

$$\begin{aligned}F_\theta &= m \left( 2 \frac{dr}{dt} \frac{d\theta}{dt} + r \frac{d^2\theta}{dt^2} \right) \\ F_r &= m \left( \frac{d^2r}{dt^2} - r \left( \frac{d\theta}{dt} \right)^2 \right).\end{aligned}$$

Our force of gravity is a **central force**: it only acts in the direction of  $\mathbf{r}$ . Hence the  $\theta$ -component of the force acting on the mass  $m$  is zero. Show that

$$r^2 \frac{d\theta}{dt} = h$$

for some constant  $h$ .

3. Suppose  $A(t)$  gives the area 'swept out' by the radius vector  $\mathbf{r}$  between some fixed time  $t_0$  and the time  $t$ . Using the fact that  $\frac{dA}{d\theta} = \frac{r^2}{2}$  (since  $dA = \pi r^2 \cdot \frac{d\theta}{2\pi}$ ), show that  $A(t_2) - A(t_1) = \frac{1}{2}h(t_2 - t_1)$ . Conclude **Kepler's second law**: if  $P$  and  $Q$  are planets with associated masses  $m_P$  and  $m_Q$ , then over a time  $\Delta t$  the planets will sweep out equal area.

**Exercise 18: Kepler's first law.** 1. Let  $z = 1/r$ . Express  $\frac{dr}{dt}$  and  $\frac{d^2r}{dt^2}$  in terms of  $\frac{dz}{d\theta}$  and  $\frac{d^2z}{d\theta^2}$ .

2. By Newton's law of gravity,  $F_r = -\frac{GMm}{r^2}$ . Thus, we have that  $\frac{d^2r}{dt^2} - r \left( \frac{d\theta}{dt} \right)^2 = -\frac{GM}{r^2}$ . Use the expression in (1b) above to show that  $\frac{d^2r}{dt^2} - \frac{h^2}{r^3} = -\frac{GM}{r^2}$ , and then apply the substitution in (a) to show that  $\frac{d^2z}{d\theta^2} + z = \frac{GM}{h^2}$ .

3. Solve the differential equation in (b). (Hint: the general solution is  $z = A \sin \theta + B \cos \theta + \frac{GM}{h^2}$ .)

4. Let us shift and rotate our coordinate axis such that  $\mathbf{r}$  is minimised when  $\theta = 0$ : in other words, we pick our axis such that the mass  $m$  is closest to the origin at  $\theta = 0$ . Since  $1/r$  is maximised when  $r$  is minimised, we have that  $\left. \frac{dz}{d\theta} \right|_{\theta=0} = 0$  and  $\left. \frac{d^2z}{d\theta^2} \right|_{\theta=0} < 0$ . Show that  $A = 0$  and  $B > 0$ .

5. Further, show that

$$r = \frac{h^2/GM}{1 + (Bh^2/GM) \cos \theta};$$

6. Write  $\varepsilon = Bh^2/GM$  in the equation of (e), and conclude **Kepler's first law** by comparison with exercise 1: the locus of  $P$  is a conic section with one focus  $O$ .<sup>a</sup>

<sup>a</sup>It can be shown that  $\varepsilon = \sqrt{1 + E \frac{2h^2}{G^2 M^2 m}}$ , where  $E$  is the potential energy associated with the mass  $m$ ; so the particular conic traced depends on the energy associated with the mass  $m$ . Thus the planets in the solar system, which have negative energy (they are trapped by the sun, and work must be done to them to pull them away), trace elliptical orbits; while bodies like comets trace parabolic and hyperbolic trajectories.

**Exercise 19: Kepler's third law.** 1. Suppose our mass traces an elliptical orbit, such that

$Bh^2/GM < 1$  in the equation  $r = \frac{h^2/GM}{1 + (Bh^2/GM) \cos \theta}$ . Let us write this in rectangular form as  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ . Using results from exercises 1 and 2 (collected in the appendix), show that  $a = \frac{h^2 a^2}{GM b^2}$  and thus that  $b^2 = \frac{h^2 a}{GM}$ . (Hint:  $a$ , the semi-major axis length, is half the sum of the maximum

and minimum values for  $r$ .)

2. If  $T$  is the period of revolution of our mass  $m$ , use Kepler's second law (in particular the formula derived in (1c)) to show that  $\pi ab = \frac{hT}{2}$  and thus, using (a), that  $T^2 = \left(\frac{4\pi^2}{GM}\right) a^3$ . Noting that this is independent of the mass  $m$ , conclude **Kepler's third law**: the square of the period of revolution of any planet is proportional to the cube of its average distance from the sun, where the constant of proportionality depends only on the mass  $M$  and some constants of nature.

---

## Section 7: Unification

---

*You know, for a mathematician, he did not have enough imagination. But he has become a poet and now he is fine.* — Hilbert (on an ex-student)

Consider the three regular conics: the ellipse, parabola, and hyperbola. In some sense we would like to be able to say that a parabola is ‘an ellipse ‘with a focus at infinity’, and that a hyperbola is what you get ‘when you push the focus past infinity and it comes back around’.

In order to make this idea make some kind of sense, we will develop a sequence of maps from the plane to a smaller space that is easier to visualise.

We will start with some technical facts.

### Exercise 20: The line.

The equation of a line in  $\mathbb{R}^2$  (the plane) is  $ax + by = c$  where at least one of  $a$  or  $b$  is nonzero. Show that the line has slope  $-a/b$ , and that it can be parameterised by a function of the form  $\phi(t) = (\lambda t + r, \mu t + s)$  for some real numbers  $\lambda, \mu, r$ , and  $s$ . (Hint: treat the cases  $a = 0$  and  $a \neq 0$  separately.)

### Exercise 21: Diffeomorphisms.

A function which has the property that for every possible output there is exactly one input producing it is called a **bijection**; the properties ‘ $f$  is a bijection’ and ‘ $f$  has an inverse’ are equivalent (every function with either property has both).

A bijective function which is differentiable and has a differentiable inverse is called a **diffeomorphism**; if two sets can be mapped onto each other by a diffeomorphism then they are called **diffeomorphic**. We shall write  $X \simeq Y$  if  $X$  is diffeomorphic to  $Y$ .

Show that:

1. Every set is diffeomorphic to itself. (Reflexivity)
2. If  $X \simeq Y$  then  $Y \simeq X$ . (Symmetry)
3. If  $X \simeq Y$  and  $Y \simeq Z$  then  $X \simeq Z$ . (Transitivity)
4. The real line  $\mathbb{R}$  is diffeomorphic to the open interval  $-\pi < x < \pi$ . (Hint: consider  $\arctan$ .)
5. The real line  $\mathbb{R}$  is diffeomorphic to the circle  $x^2 + y^2 = 1$  with a point removed. Here are two methods:
  - (a) Define  $\phi$  from the interval  $-\pi < x < \pi$  to the circle minus the point  $(-1, 0)$  to be the usual parameterisation  $\phi(t) = (\cos t, \sin t)$ . This is differentiable; show it has a differentiable inverse and is therefore a diffeomorphism. Hence the circle and the interval are diffeomorphic; use transitivity and (4) above to finish.
  - (b) Show that for every real  $t \neq 0$  the line  $y = t(x + 1)$  intersects the circle in a single point other than  $(-1, 0)$ , and intersects the line  $x = 1$  precisely once as well. Prove that every  $P$  on the circle, except for  $(-1, 0)$ , corresponds to a distinct point on the line in this way (that is, if  $Q$  is on the line then there is precisely one  $P$  on the circle which corresponds to it). Prove that the function  $\psi$  that sends each  $P$  on the circle to its corresponding point on the line is the required diffeomorphism.

### Exercise 22: Projective 2-space.

We want to be able to say that a parabola is an ellipse with one point ‘sent to infinity’, and that the hyperbola is an ellipse with two such ‘points at infinity’. We therefore need to extend our set of points to include points at infinity.

1. In  $\mathbb{R}^3$ , let  $\Pi$  be the *plane*  $z = 1$ . Clearly  $\Pi \simeq \mathbb{R}^2$ . Let  $\Sigma$  be the sphere  $x^2 + y^2 + z^2 = 1$  centred at  $O = (0, 0, 0)$ .

Show that for every point  $P$  in  $\Pi$  there are precisely two points on the sphere that lie on the line  $OP$ , and these points are diametrically opposite. Conversely, for every pair of diametrically

opposite points that lie on  $\Sigma$  but do not lie in the plane  $z = 0$  there is precisely one point lying in  $\Pi$ .

2. Further, convince yourself that if a point  $P$  moves in a straight line out from the origin of  $\mathbb{R}^2$  (that is, the point  $(0, 0, 1)$  in the plane  $\Pi$ ) then the corresponding line  $OP$  approaches the  $z = 0$  plane. More formally, prove that if  $L$  is any line in  $\Pi$ , parameterised by  $L(t) = (\lambda t + r, \mu t + s, 1)$  for some real numbers  $\lambda, \mu, r$ , and  $s$ , then as  $t \rightarrow \infty$ ,  $OL(t)$  (that is, the line joining  $O$  and  $L(t)$ ) approaches a line in the  $z = 0$  plane (and the points on  $\Sigma$  corresponding to  $L(t)$  approach a single diametrically opposed pair of points of  $\Sigma$  in the  $z = 0$  plane).

Hence pairs of diametrically opposed points on the sphere (minus the equator) correspond to points in the plane (by 1.), and diametrically opposed points on the equator correspond to ‘directions’ (possible slopes of lines) in the plane. But this is precisely what we wanted: a new space which ‘contains’ the old space, together with a set of new points that correspond to directions ‘towards infinity’.

Define  $\mathbb{P}^2$ , **projective 2-space**, to be the set of pairs of diametrically opposed points of the sphere  $\Sigma$ . If one such pair is  $\{P, Q\}$  we will denote this point of projective 2-space by either  $[P]$  or  $[Q]$ . (Clearly if we know  $P$  then we can compute  $Q$  easily, and vice-versa, since we know they are opposite.)

3. Show that the correspondence described in 1. above is precisely given by

$$\begin{aligned} \varphi : \mathbb{R}^2 &\rightarrow \mathbb{P}^2 \\ (x, y) &\mapsto \left[ \frac{1}{\sqrt{x^2 + y^2 + 1}}(x, y, 1) \right]. \end{aligned}$$

4. If we ignore the lower hemisphere (because it doesn’t give us any new information) and ignore the equator (because it corresponds to the points at infinity), show that the (restricted) map

$$\begin{aligned} \varphi : \mathbb{R}^2 &\rightarrow \mathbb{P}^2 \\ (x, y) &\mapsto \frac{1}{\sqrt{x^2 + y^2 + 1}}(x, y, 1) \end{aligned}$$

is a diffeomorphism.

Hence the embedding of  $\mathbb{R}^2$  into our new space is a *true* embedding — it conveys the same information.

Continuing on with the idea of 4. above, we can stop thinking about the lower hemisphere and just focus on the upper hemisphere of  $\Sigma$  (which is an embedding of our usual plane) together with the equator (which contains the points at infinity, as long as we remember that opposite points are ‘glued together’).

It is clear that can draw the hemisphere ‘from above’ by just collapsing the  $z$ -coordinate (setting it to zero so that the entire hemisphere lies in one plane).

5. Show that the projection map  $\pi$  from the upper-hemisphere-with-equator to the disc  $x^2 + y^2 \leq 1$  given by  $\pi(x, y, z) = (x, y)$  is a diffeomorphism. (You simply need to exhibit an inverse: but note that since  $x^2 + y^2 + z^2 = 1$  must hold then you can find  $z^2$  given  $x$  and  $y$ ; and then take the positive square root, since we care about the upper hemisphere.)

Since the new disc space is diffeomorphic to  $\mathbb{P}^2$  we will denote both by the same name (i.e.  $\mathbb{P}^2$  will be used to refer to the disc as well as to the sphere model).

We therefore obtain the map from  $\mathbb{R}^2$  to our new space given by

$$\begin{aligned} \psi : \mathbb{R}^2 &\rightarrow \mathbb{P}^2 \\ (x, y) &\mapsto \pi(\varphi(x, y)) = \left[ \frac{1}{\sqrt{x^2 + y^2 + 1}}(x, y) \right]. \end{aligned}$$

This map ‘collapses’ (the ‘real’ term is **compactifies**, but the definition is quite technical) the plane-with-points-at-infinity to the disc  $x^2 + y^2 \leq 1$ , and allows us to see all the points on our conics at once.

**Exercise 23: Examples of conics in  $\mathbb{P}^2$ .**

**Exercise 24: Bezout’s theorem for conics.**

---

## Appendix A: List of formulae

---

**Polar form of conic with focus at  $(0, 0)$**

$$\frac{l}{r} = 1 + \varepsilon \cos \theta.$$

**Rectangular form of ellipse or hyperbola with centre at  $(0, 0)$**

$$\frac{x^2}{a^2} \pm \frac{y^2}{b^2} = 1, \text{ where } 2a \text{ is the } x\text{-dimension and } 2b \text{ is the } y\text{-dimension}$$

**Relation of rectangular form to eccentricity**

$$a^2 \varepsilon^2 = a^2 - \beta^2, \text{ where } \beta = b \text{ for an ellipse and } \beta = ib \text{ for a hyperbola.}$$

**Relation of rectangular form to latus rectum**

$$l = b^2/a.$$

**Relation of rectangular form to foci**

$$(\pm c, 0) \text{ where } c^2 = a^2 - \beta^2.$$

**Rectangular form of parabola with vertex at  $(0, 0)$**

$$y^2 = 4ax, \text{ where } (0, a) \text{ is the focus and } x = -a \text{ is the directrix.}$$

---

## Appendix B: Further reading

---

See references in text, as well as:

- Marcel Berger, *Geometry I and II*. Springer-Verlag (1987).
- H.S.M. Coxeter, *Introduction to geometry*. John Wiley & Sons (1961).
- Stephen D. Fisher, *Complex variables*. Dover (1999).
- Robin Hartshorne, *Geometry: Euclid and beyond*. Springer (2000).
- Keith Kendig, *Conics*. Mathematical Association of America (2005).
- Morris Kline, *Mathematics for the nonmathematician*. Dover (1985).
- John M. Lee, *Axiomatic geometry*. American Mathematical Society (2017).
- George Salmon, *A treatise on conic sections*. Longmans, Green (1900).
- George F. Simmons, *Differential equations with applications and historical notes*. McGraw-Hill (1991).
- Marta Sved, *Journey into geometries*. Mathematical Association of America (1991).