

Every compound in the AXIOM catalogue in one line each — what it does, how it is given, and how strong the evidence is. For mechanism, trial figures, safety and sources, see the full compound guide.

REGISTRATION-GRADE

Approved medicine or completed Phase 3

HUMAN TRIALS

Controlled human data; not FDA/EMA approved

PRECLINICAL

Cell and animal models only

01 GLP-1s & Weight Loss

3

Cagrilintide

HUMAN TRIALS

Satiety through the amylin pathway — weight reduction without touching a GLP-1 receptor.

Subcutaneous · 10 mg

Retatrutide

HUMAN TRIALS

The strongest weight-loss result yet recorded in clinical trials — nearly 29% of body weight.

Subcutaneous · 5 mg · 10 mg · 15 mg · 20 mg · 30 mg · 60 mg

Tirzepatide

REGISTRATION-GRADE

The approved benchmark of the class — about a fifth of body weight in registration trials.

Subcutaneous · 10 mg · 30 mg · 40 mg

02 GH Secretagogues

5

CJC-1295 (No DAC) + Ipamorelin

PRECLINICAL

A synchronised growth-hormone pulse driven from two receptors at once.

Subcutaneous · 10 mg · 20 mg

HGH 191AA (Somatropin)

REGISTRATION-GRADE

Growth hormone itself — the reference molecule for the entire axis.

Subcutaneous · 36 IU · 40 IU

Ipamorelin

PRECLINICAL

The cleanest pulse in the secretagogue class — GH without cortisol, prolactin or appetite signal.

Subcutaneous · 10 mg

Ipamorelin + Tesamorelin

PRECLINICAL

The selective ghrelin pulse paired with the one GHRH analogue that carries registration-grade evidence.

Subcutaneous · 18 mg

Tesamorelin

REGISTRATION-GRADE

The registration-grade GHRH analogue — documented visceral-fat reduction in Phase 3 trials.

Subcutaneous · 10 mg · 20 mg

03 Healing & Repair

8

BPC-157

PRECLINICAL

The most-studied repair peptide in the catalogue — soft tissue, tendon and gut lining in animal models.

Subcutaneous (orally stable) · 10 mg

BPC-157 + TB-500 (Wolverine)

PRECLINICAL

Angiogenesis and cell migration in one preparation — the two repair axes that don't overlap.

Subcutaneous · 20 mg

GHK-Cu

HUMAN TRIALS

Collagen remodelling with human trial data behind it.

Topical or subcutaneous · 50 mg · 100 mg

KPV

PRECLINICAL

The anti-inflammatory signal of alpha-MSH, without its pigmentary activity.

Subcutaneous, oral or topical · 10 mg

KLOW (BPC-157 + TB-500 + GHK-Cu + KPV)

PRECLINICAL

Four repair mechanisms in a single preparation.

Subcutaneous · 80 mg

LL-37

HUMAN TRIALS

The body's own antimicrobial — wound closure where healing has stalled.

Topical or local · 5 mg

Peg MGF

PRECLINICAL

The repair signal of loaded muscle, extended to a research-usable half-life.

Subcutaneous · 2 mg

TB-500

PRECLINICAL

Cell migration to the injury site — the actin-binding arm of repair research.

Subcutaneous · 10 mg

04 Brain Health & Nootropics

6

Adamax

PRECLINICAL

A designed successor to Semax — with no published evidence of its own yet.

Intranasal · 10 mg

Cerebrolysin

HUMAN TRIALS

The most clinically documented neurotrophic in the catalogue — stroke and dementia trial data.

Intravenous or intramuscular · 60 mg

Pinealon

PRECLINICAL

A gene-level neuronal bioregulator hypothesis — cell and rodent data only.

Subcutaneous or intranasal · 20 mg

Selank

HUMAN TRIALS

Anxiety reduction comparable to benzodiazepines in Russian trials — without the sedation.

Intranasal · 10 mg

Semax

HUMAN TRIALS

Two decades of clinical use in stroke recovery and cognition.

Intranasal (or subcutaneous) · 10 mg

Selank + Semax

HUMAN TRIALS

Calm and focus studied side by side — the two Russian regulatory peptides together.

Intranasal · 20 mg

05 Energy & Endurance

5

5-Amino-1MQ

PRECLINICAL

An NAD+ -preserving switch aimed specifically at fat tissue.

Oral · 50 mg

AICAR

PRECLINICAL

The original exercise mimetic — endurance gains in untrained mice.

Intravenous or subcutaneous · 50 mg

L-Carnitine (Injectable)

REGISTRATION-GRADE

A fat-oxidation shuttle, delivered at full systemic dose.

Intravenous or intramuscular · 5000 mg

MOTS-c

PRECLINICAL

A mitochondrial signal the body raises during exercise — and loses with age.

Subcutaneous · 10 mg · 40 mg

SLU-PP-332 (Injectable)

PRECLINICAL

The newest exercise-mimetic axis — endurance biology through the ERR receptors.

Subcutaneous · 5 mg

06 Immunity

1

Thymosin Alpha-1

HUMAN TRIALS

An immune-system modulator with controlled trial data in infection and sepsis.

Subcutaneous · 10 mg

07 Sexual Health

5

HCG

REGISTRATION - GRADE

An LH mimic — the standard tool for driving gonadal testosterone.
Subcutaneous or intramuscular · 10000 IU

HMG

REGISTRATION - GRADE

Both gonadotropins in one preparation — the FSH complement to HCG.
Subcutaneous or intramuscular · 75 IU

Kisspeptin

HUMAN TRIALS

The master switch at the top of the reproductive axis.
Subcutaneous, IV or intranasal · 10 mg

Oxytocin Acetate

HUMAN TRIALS

The bonding hormone — a large but contested social-behaviour literature.
Intranasal, IV or intramuscular · 10 mg

PT-141

REGISTRATION - GRADE

Sexual desire acted on centrally, in the brain — not through the vasculature.
Subcutaneous (autoinjector) · 10 mg

08 Longevity & Cellular Repair

5

DSIP

HUMAN TRIALS

A sleep-pressure modulator — most active exactly when sleep is disturbed.
Subcutaneous or intravenous · 5 mg · 10 mg

Epithalon

PRECLINICAL

Telomerase activation — demonstrated in human cells, not yet in humans.
Subcutaneous or intranasal · 50 mg

Humanin

PRECLINICAL

A mitochondrial cytoprotectant that tracks with human longevity.
Subcutaneous or intraperitoneal · 10 mg

NAD+

HUMAN TRIALS

Refilling the coenzyme pool that halves between your 20s and 50s.
Intravenous (or subcutaneous) · 500 mg · 1000 mg

SS-31

REGISTRATION - GRADE

The registration-grade mitochondrial peptide — approved for a genetic energy disorder.
Subcutaneous · 50 mg

Full detail — mechanism, trial figures, safety signals, regulatory status, glossary and sources — is in the AXIOM Compound Guide. Pricing is published separately in the AXIOM Price List.

COMPLIANCE

Research Use Only. Peptides listed here are intended exclusively for in-vitro laboratory research — not drugs, food, cosmetics, or supplements, and not for human or veterinary consumption, diagnosis, or treatment. No dosing or usage guidance is provided. Evidence grades reflect the state of published research at the time of writing and are not claims of efficacy. Several compounds are controlled or regulated in many jurisdictions; buyers are solely responsible for confirming the laws that apply to them.