

# 7

## 7.1

We prove  $M \wedge N \leq M \times N$ . Notice for  $M \wedge N$  if  $\Sigma_m \neq \Sigma_n$  then  $M \wedge N = \emptyset$  which is always a subset of any set. Therefore, we assume from now on  $\Sigma_m = \Sigma_n$ . Also checkout these definitions:

$$(q_m, q_n)(\delta_m \wedge \delta_n)_a = (q_m \delta_a^m, q_n \delta_a^n) \quad (1)$$

$$(q_m, q_n)(\delta_m \times \delta_n)_{(a_m, a_n)} = (q_m \delta_{a_m}^m, q_n \delta_{a_n}^n) \quad (2)$$

We choose for the covering  $(\eta, \xi) \text{ id}_{(q_m, q_n)}$  as  $\eta$  and define  $\xi : \Sigma_n \rightarrow \Sigma_m \times \Sigma_n; a_m \mapsto (a_m, a_m)$ . which is well-defined due the assumption  $\Sigma_n = \Sigma_m$ .

Now we show  $\eta((q_m, q_n))(\delta^m \wedge \delta^n)_{a_m} \subseteq \eta((q_m, q_n))(\delta^m \times \delta^n)_{\xi(a_m)}$ :

$$\eta((q_m, q_n))(\delta^m \wedge \delta^n)_{a_m} = (q_m, q_n)(\delta^m \wedge \delta^n)_{a_m} \quad (\text{def. } \eta, 1)$$

$$= (q_m \delta_{a_m}^m, q_n \delta_{a_m}^n) \quad (2)$$

$$= (q_m, q_n)(\delta_m \times \delta_n)_{(a_m, a_m)} \quad (\text{def. } \xi)$$

$$= (q_m, q_n)(\delta_m \times \delta_n)_{\xi(a_m)} \quad (\text{def. } \eta)$$

$$= \eta((q_m, q_n))(\delta^m \times \delta^n)_{\xi(a_m)}$$

□

## 7.2

## 7.3

We prove  $P \leq N \omega R$  under the assumption  $M \leq N$  and  $P \leq M \omega R$ . Let  $\eta_1 : Q_n \rightarrow Q_m$  and  $\eta_2 : Q_m \times Q_r \rightarrow Q_p$  be the  $\eta$ -functions of  $M \leq N$  and  $P \leq M \omega R$ . Furthermore let  $\xi_1 : \Sigma_p \rightarrow \Sigma_r$  be the  $\xi$ -function of  $P \leq M \omega R$ .

We define  $\eta : Q_n \times Q_r \rightarrow Q_p; (q_n, q_r) \mapsto \eta_2(\eta_1(q_n), q_r)$ . Moreover we choose  $\xi_1$  as  $\xi$ .

Finally, we show  $\eta((q_n, q_r))\delta_{a_p}^p \subseteq \eta((q_n, q_r))\delta_{\xi(a_p)}^\omega$ .

$$\begin{aligned}
\eta((q_n, q_r))\delta_{a_p}^p &= \eta_2((\eta_1(q_n), q_r))\delta_{a_p}^p && (\text{def. } \eta, \eta_1) \\
&= \eta_2(q_m, q_r)\delta_{a_p}^p && (\text{def. } \eta_2) \\
&= q_p\delta_{a_p}^p && (P \leq M\omega R) \\
&\subseteq \eta_2((q_m, q_r)\delta_{\xi(a_p)}^\omega) && (\text{def. } \xi) \\
&= \eta_2((q_m, q_r)\delta_{a_r}^\omega) && (\text{def. } \delta^\omega) \\
&= \eta_2((q_m\delta_{\omega_{a_r}(q_r)}^m, q_r\delta_{a_r}^r)) && (\text{def. } \omega) \\
&= \eta_2((q_m\delta_{a_m}^m, q_r\delta_{a_r}^r)) && (M \leq N) \\
&= \eta_2((\eta_1(q_n\delta_{a_n}^n), q_r\delta_{a_r}^r)) && (\text{def. } \eta) \\
&= \eta((q_n\delta_{a_n}^n, q_r\delta_{a_r}^r)) && (\text{def. } \omega) \\
&= \eta((q_n\delta_{\omega_{a_r}(q_r)}^n, q_r\delta_{a_r}^r)) && (\text{def. } \delta^\omega) \\
&= \eta((q_n, q_r)\delta_{a_r}^\omega) && (\text{def. } \xi) \\
&= \eta((q_n, q_r)\delta_{\xi(a_p)}^\omega)
\end{aligned}$$

□

## 7.4