

Simplified Discharge Summary

What's Wrong

*You have pneumonia in the lower part of your right lung. This kind of pneumonia is called community-acquired pneumonia. You also have mild asthma that comes and goes, and allergies from seasons like springtime. When you came to the hospital, your oxygen was low, but it's better now. **You were discharged from the hospital on Feb 18th and should follow the instructions below.***

Your Medicines

Azithromycin

250mg

Once daily

- 1. Take 1 tablet every day at the same time.*
- 2. You can take it with or without food.*
- 3. Take all 4 days of medicine. Do not stop early, even if you feel better.*
- 4. Do not take antacids with aluminum or magnesium for 1 hour before or 2 hours after this medicine.*
- 5. Call your doctor if you have bad diarrhea or blood in your poop.*
- ~~*6. Do not take this medicine if you are allergic to azithromycin, erythromycin, or macrolides.*~~
- ~~*7. Do not take this medicine if you use heart medicines like amiodarone or sotalol.*~~

Prednisone

40mg

Once daily (morning, with food)

- 1. Take 1 tablet in the morning with food or milk.*
- 2. Take it for 3 days exactly and finish all doses.*
- 3. Do not take it on an empty stomach.*
- ~~*4. Stay away from anyone who has chickenpox or shingles.*~~

~~5. Tell your doctor before getting any vaccines.~~

~~6. This medicine may make you hungrier, cranky, have trouble sleeping, make your face red, or increase blood sugar.~~

~~7. Do not take this medicine if you have an untreated fungal infection or are allergic to corticosteroids.~~

Salbutamol

2 puffs (90mcg/puff)

Every 4–6 hours as needed

1. Shake the inhaler well before each use.

2. Breathe out fully.

3. Inhale 2 puffs slowly and hold your breath for 10 seconds each time.

4. Wait 1 minute between each puff.

5. Rinse your mouth after using the inhaler if you experience discomfort in your throat.

6. Go to the emergency room if you use more than 8 puffs in 24 hours and still feel bad, or if your symptoms do not get better in 20 minutes.

7. Do not use the inhaler too much.

Cetirizine

10mg

Once daily

1. Take 1 tablet every day at the same time.

2. You can take it with or without food.

3. This medicine can make you sleepy. Do not drive or use machines if you feel sleepy.

4. Do not drink alcohol while on this medicine because it will make you drowsier.

~~5. Do not take this medicine if you have severe kidney problems or are allergic to cetirizine or levocetirizine.~~

Your Appointments

Appointment 1: You need a check-up after pneumonia **in your GP clinic**. The doctor will make sure you are better, look at your chest X-ray, check that you have taken all your antibiotics, ask if you have symptoms, and go over your asthma plan.

On or before February 28, 2026

To be confirmed

General Doctor (GP)

Not specified

(555) 204-9300 (Call GP to book)

Appointment 2: You need to check your breathing and asthma **in an outpatient respiratory clinic**. The doctor will do tests to see how your lungs work and decide if you need long-term asthma medicine.

To be confirmed within 4-6 weeks

To be scheduled

Lung Clinic (Outpatient Respiratory Clinic)

Not specified

(555) 204-7700 (Contact number if not called by March 7)

What to Do

- 1. Rest but increase your activity a little bit when you feel stronger.*
- 2. Sleep with your head raised to help you breathe.*
- 3. Drink 6–8 glasses of water each day. Avoid alcohol until you finish all of your antibiotics.*
- 4. Do breathing exercises three times a day to help your lungs. Breathe slowly through your nose for 4 seconds, hold for 2 seconds, then exhale slowly through your mouth for 6 seconds. Repeat ten times.**
- 5. Only go back to work or heavy activities when the doctor says it's okay.*
- 6. Stay away from cigarette smoke, dust, and strong smells while you are recovering.*
- 7. Call your doctor or go to the hospital if your symptoms get worse or you are not getting better.*

Warning Signs

- 1. Call 911 or go to the **emergency room** right away for:*
 - Severe shortness of breath while resting.*
 - Blue or grey lips or fingers.*
 - Sharp chest pain that moves to your arm or jaw.*

- *Fever over 39.5°C that doesn't go down with medicine.*
- *Coughing up a lot of blood.*
- *Feeling confused or passing out.*

2. Call your doctor if:

- *You still have a fever over 38°C after 2 days on antibiotics.*
- *You get a rash or reaction to medicine.*
- *Your wheezing does not stop even after using an inhaler.*
- *Your cough gets worse or you have other new symptoms like swelling in your leg.*