

MATH 1104 Linear Algebra

Study Session Questions

December 28, 2025

Learning Objectives

- You should be confident solving linear systems using row reduction and the matrix equation $Ax = b$, and you should know what linear dependence and span are.
 - You should be confident applying matrix operations, including inverses, and you should know how to calculate determinants and use Cramer's rule.
 - You should be confident analyzing vector spaces by identifying subspaces, computing dimension/rank, and finding bases for a space.
 - You should be confident working with complex numbers and computing eigenvalues and eigenvectors to study diagonalizability.
 - You should be confident calculating inner products, applying projections, and implementing the Gram-Schmidt process for orthonormal bases.
1. Find the general solution to the corresponding homogenous system of linear equations as defined below.

$$x_1 + x_2 - x_3 = 1$$

$$2x_1 + 3x_2 - 4x_3 = 4$$

$$x_1 + 2x_2 - 3x_3 = 2$$

- A. no solutions
 - B. $(x_1, x_2, x_3) = t(1, 0, 1) + s(0, 1, -2)$
 - C. $(x_1, x_2, x_3) = t(-1, 2, 1)$
 - D. $(x_1, x_2, x_3) = (1, 1, 0)$
2. How can we know the size of the solution set of a linear system? Select all true statements.
 - ☐ An inconsistent system (i.e $[0 \ 0 \ 0 \mid 3]$) has no solution
 - ☐ If the system is square and $\det(A) = 0$, the solution is unique
 - ☐ If all columns in the coefficient matrix are pivots, the solution is unique
 - ☐ If there are any free variables, the system has infinite solutions
 - ☐ If the rank is less than the number of variables, there are no solutions
 - ☐ A consistent system with fewer equations than variables has infinite sols.

3. Suppose $b = x_1a_1 + x_2a_2$ as defined below. What is the sum $x_1 + x_2$?

$$a_1 = \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}, \quad a_2 = \begin{bmatrix} -1 \\ -1 \\ -2 \end{bmatrix}, \quad b = \begin{bmatrix} 9 \\ 2 \\ 4 \end{bmatrix}$$

- A. 3
- B. 6
- C. 9
- D. 12

4. Given the vectors defined below, which of the following statements are true? Select all that apply.

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}, \quad v_3 = \begin{bmatrix} -2 \\ 0 \\ 0 \end{bmatrix}$$

- ☐ $(3, -1, 3) \in \text{Span}\{v_1, v_2\}$
- ☐ The zero vector is in $\text{Span}\{v_1, v_3\}$
- ☐ $(0, 1, 2) \in \text{Span}\{v_1, v_3\}$
- ☐ $(7, -12, 4) \in \text{Span}\{v_1, v_2, v_3\}$
- ☐ All three vectors are linearly DEPENDENT.
- ☐ All three vectors span \mathbb{R}^3 .

5. For what values (k, h) does the system defined below have a unique solution?

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 3 & 4 \\ 2 & 1 & k \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} h \\ 5 \\ 7 \end{bmatrix}$$

- A. $k = 2, h = -3$
- B. $k \neq -2$, for all $h \in \mathbb{R}$
- C. $k \neq -3$, for all $h \in \mathbb{R}$
- D. $k \neq 3, k \neq 2$
- E. $h = 4$, for all $k \in \mathbb{R}$
- F. $k \neq 4$, for all $h \in \mathbb{R}$

6. For what values of m is the set $\{u, v\}$ linearly dependent? Type them in numerical order, comma and space separated.

$$u = \begin{bmatrix} m \\ 1 \end{bmatrix}, \quad v = \begin{bmatrix} 2 \\ m+1 \end{bmatrix}$$

7. Consider the following augmented matrix of a system of linear equations below. Select all statements that are TRUE.

$$\left[\begin{array}{ccc|c} 1 & 3 & 2 & 5 \\ 0 & 1 & 4 & -1 \\ 0 & 0 & a^2 - 4a & a - 4 \end{array} \right]$$

- ☐ When $a = 0$, there are infinitely many solutions
 - ☐ When $a \neq 0$, the system is consistent
 - ☐ When $a = 2$, the system has no solutions
 - ☐ When $a = 2$, the columns of the coefficient matrix are linearly independent
 - ☐ When $a = 4$, there are infinitely many solutions
 - ☐ When $a \neq 4$, the columns of the coefficient matrix are linearly independent
8. Compute the original matrix A given its inverse, A^{-1} , defined below.

$$A^{-1} = \begin{bmatrix} 1 & -2 & 5 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix}$$

A. $\begin{bmatrix} -1 & 2 & -5 \\ 0 & -1 & 4 \\ 0 & 0 & -1 \end{bmatrix}$

B. $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

C. $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}$

D. $\begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 2 & 5 & 1 \end{bmatrix}$

E. the original matrix is the identity matrix

F. the original matrix does not exist

9. Which of the following matrices are invertible? Select all that apply.

☐ $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

☐ $\begin{bmatrix} 3 & 1 & 2 \\ 6 & 2 & 4 \\ 0 & -1 & -2 \end{bmatrix}$

☐ $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

☐ $\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

10. Let A , B , and C each be 3×3 matrices. If $\det(A) = 2$, $\det(B) = 4$, and $\det(C) = 8$, select all responses that must be true.

- ☐ $\det(3A) = 18$
☐ $\det(ABC) = 64$
☐ $\det(B^{-1}C^T) = 2$
☐ $\det((2B)^{-1}) = \frac{1}{8}$
☐ $\det[A^{-1}(B^T)^2] = 8$
☐ $\det(AI) = 2$

11. Let A and B be 3×3 matrices as defined below. If $\det(A) = 4$, determine $\det(B)$.

$$A = \begin{bmatrix} a & b & c \\ u & v & w \\ x & y & z \end{bmatrix}, \quad B = \begin{bmatrix} a & 3b + 5a & c \\ u & 3v + 5u & w \\ x & 3y + 5x & z \end{bmatrix}$$

- A. $\det(B) = -4$
B. $\det(B) = 60$
C. $\det(B) = 4$
D. $\det(B) = 8$
E. $\det(B) = 12$
F. $\det(B) = 20$

12. Let A and B be matrices as defined below. Find the matrix X such that $3X - B = AX + 2I$.

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}, \quad 3X + B = AX + 2I$$

- A. $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
B. $\begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$
C. $\begin{bmatrix} 3 & 2 \\ 1 & 2 \end{bmatrix}$
D. $\begin{bmatrix} 2 & 2 \\ 1 & 2 \end{bmatrix}$

13. Find all values of k for which the matrix A defined below is NOT invertible.

$$A = \begin{bmatrix} 2 & 1 & k \\ 4 & k & 9 \\ k & 0 & 0 \end{bmatrix}$$

- A. $k = -3$
B. $k = 3$

- C. $k = -1$
- D. $k = 1$
- E. $k = 0$
- F. such a k does not exist

14. How can Cramer's Rule be used to solve linear systems? Briefly explain in your own words.

15. What is the coordinate vector of $x = (5, 0, 4)$ relative to basis B as defined below?

$$B = \left\{ \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

- A. $[x]_B = (5, 3, -3)$
- B. $[x]_B = (3, -2, 2)$
- C. $[x]_B = (1, -2, -4)$
- D. $[x]_B = (-5, 0, -4)$

16. Which of the following statements about properties of transformations are TRUE? Select all that apply.

- ☐ A linear map is invertible only if it is both one-to-one and onto
- ☐ The kernel is the set of all vectors that map to the zero vector
- ☐ The range of a transformation is the span of the columns of its matrix
- ☐ A transformation from \mathbb{R}^n to \mathbb{R}^m is always invertible if $n > m$
- ☐ Any set of independent vectors from the domain can form the standard matrix
- ☐ A transformation is one-to-one if its kernel contains only the zero vector

17. Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the linear transformation as defined below. Select the most true response.

$$T \left(\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \right) = \begin{bmatrix} x + 3w \\ 2x - y + z \\ 4z - 5w \end{bmatrix}$$

- A. T is one-to-one (injective).
- B. T is onto (surjective).

- C. T is neither injective nor surjective.
- D. T is bijective, and thus, also invertible.
18. Let A be a 6×9 matrix such that row echelon form has 5 pivot positions. Which of these statements are TRUE?
- ☐ $\dim(\text{Nul } A) = 4$
 - ☐ $\text{Rank } A = 5$
 - ☐ $\text{Nul } A = \mathbb{R}^4$
 - ☐ $\dim(\text{Col } A) = 5$
19. Let matrix A be as defined below. Which of the following sets is a basis for the null space of A ?

$$A = \begin{bmatrix} 1 & 0 & -3 & 2 \\ 0 & 1 & 2 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

A. $\left\{ \begin{bmatrix} 3 \\ -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 4 \\ 0 \\ 1 \end{bmatrix} \right\}$

B. $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$

C. $\left\{ \begin{bmatrix} 3 \\ 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -4 \\ 1 \\ 0 \end{bmatrix} \right\}$

D. $\left\{ \begin{bmatrix} -3 \\ 2 \\ 1 \\ 1 \end{bmatrix} \right\}$

20. Select all of the sets below that are vector spaces.

☐ $\left\{ \begin{bmatrix} p \\ q \\ r \end{bmatrix} : 3p - 4q = r, 2q = p + 3q \right\}$

☐ $\left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : x - 2y + z = 5 \right\}$

☐ $\left\{ \begin{bmatrix} 3 \\ c \\ d \end{bmatrix} : c, d \text{ real} \right\}$

☐ $\left\{ \begin{bmatrix} -a + 2b \\ a - 2b \\ 3a - 6b \end{bmatrix} : a, b \text{ real} \right\}$

21. Let A be defined below and $w = (0, 0, 4, -2)$. Select the most correct response.

$$A = \begin{bmatrix} 3 & 0 & 4 & 8 \\ 1 & 0 & 3 & 6 \\ 4 & -2 & 2 & 4 \\ 4 & 1 & 0 & 0 \end{bmatrix}$$

- A. w is in $\text{Col } A$.
 - B. w is in $\text{Nul } A$.
 - C. w is in both $\text{Col } A$ and $\text{Nul } A$.
 - D. w is in neither $\text{Col } A$ nor $\text{Nul } A$.
22. Let $v = (0, 1, 1)$, an eigenvector of A . What is the corresponding eigenvalue?

$$A = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix}$$

- A. -4
 - B. 4
 - C. -3
 - D. 3
 - E. -2
 - F. 2
23. What is the polar form of the complex number $z = 3 - 3i$?
- A. $6(\cos(-\pi/4) - i \sin(-\pi/4))$
 - B. $3\sqrt{2}(\cos(-\pi/4) + i \sin(-\pi/4))$
 - C. $3\sqrt{2}(\cos(7\pi/4) - i \sin(7\pi/4))$
 - D. $3\sqrt{2}(\sin(\pi/4) + i \cos(\pi/4))$
24. Which of the following equations involving complex numbers are true? Select all that apply.
- ☐ The modulus of $3 + 4i$ is 5.
 - ☐ $(2i)^2 = 4i$
 - ☐ $(2 + 3i)(2 - 3i) = 13$
 - ☐ $i^6 = i$
 - ☐ $\frac{3 + 4i}{1 + 2i} = i - 1$
 - ☐ $\frac{4 + 3i}{2 - i} = 5$
25. Suppose A is a 4×4 NON-invertible matrix. Which of the following cannot be the characteristic polynomial of A ?
- A. $\lambda^4(\lambda - 2)$
 - B. $-(\lambda + 3)^2$

- C. $\lambda^2 + 4\lambda$
D. $-\lambda(\lambda^3 - 5)$

26. $\lambda = 4$ is an eigenvalue for the matrix B as defined below. Which one of the following sets is a basis for E_4 ?

$$B = \begin{bmatrix} 6 & -2 & 4 \\ 2 & 2 & 4 \\ 2 & -2 & 8 \end{bmatrix}$$

- A. $\left\{ \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$
B. $\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} \right\}$
C. $\left\{ \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix} \right\}$
D. $\left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \right\}$

27. Exactly one of the following statements involving diagonalizability is TRUE. Which one?

- A. Every invertible matrix is diagonalizable.
B. Every diagonalizable matrix is invertible.
C. All $n \times n$ diagonalizable matrices have n distinct eigenvalues.
D. All $n \times n$ diagonalizable matrices have n linearly independent eigenvectors.

28. Select the statement that is most true involving the diagonalizability of matrices A and B .

$$A = \begin{bmatrix} 3 & -1 \\ 1 & 5 \end{bmatrix}, \quad B = \begin{bmatrix} 3 & 6 \\ 4 & 1 \end{bmatrix}$$

- A. A is diagonalizable.
B. B is diagonalizable.
C. Both A and B are diagonalizable.
D. Neither A nor B are diagonalizable.

29. Let $u = (1, 2, 3)$ and $v = (3, 2, 1)$. Select all statements that are TRUE.

- ☐ $u \cdot v = 10$
☐ $\|u\| = \|v\|$
☐ The vectors u and v are orthogonal
☐ The sum $u + v$ is orthogonal to the difference $u - v$

30. Determine the value k for which $v = (k, 5, 7)$ lies in W^\perp with W as defined below.

$$W = \text{Span} \left\{ \begin{bmatrix} 3 \\ 4 \\ -2 \end{bmatrix}, \begin{bmatrix} 7 \\ 0 \\ 2 \end{bmatrix} \right\}$$

- A. $k = -1$
- B. $k = 1$
- C. $k = -2$
- D. $k = 2$
- E. any k
- F. no such k exists

31. Let $u = (2, 1)$ and $v = (1, 3)$. What is the angle between u and v ?

- A. $\pi/2$
- B. $\pi/3$
- C. $\pi/4$
- D. $\pi/6$

32. Let $W = \text{Span}\{(1, 2, 2)\}$ and $x = (2, 3, 5)$. What is the distance from x to W ?

- A. $3\sqrt{3}$
- B. $-\frac{7}{2}$
- C. $2\sqrt{5}$
- D. 2

33. Briefly explain how to use the Gram-Schmidt process to turn linearly-independent vectors into an orthonormal basis.

34. Suppose the columns of B form an orthogonal basis for \mathbb{R}^4 . If $x = c_1b_1 + c_2b_2 + c_3b_3 + c_4b_4$, what is the value of c_2 ?

$$B = \begin{bmatrix} 1 & 2 & 4 & 1 \\ 2 & -1 & 8 & 2 \\ 3 & 0 & -4 & 9 \\ 4 & 0 & -2 & 8 \end{bmatrix}, \quad x = \begin{bmatrix} 7 \\ 5 \\ -1 \\ 4 \end{bmatrix}$$

A. $\frac{5}{4}$

B. $-\frac{4}{25}$

C. $\frac{9}{5}$

D. 1