

Quantum Field Theory

Arya Farahi

Copyright ©2012 by Arya Farahi

All rights reserved.

ISBN ...

... Publications

To my professor

Contents

1	Classical Scalar Field	3
1.1	Scalar field theory	3
1.2	Classical Scalar Field Theory	4
1.2.1	Linear (free) theory	4
1.2.2	Nonlinear (interacting) theory	4
1.3	Dimensional analysis and scaling	5
1.3.1	Scaling Dimension	5
1.4	Scale Invariance	6
1.4.1	Conformal Invariance	6
1.5	ϕ^4 theory	7
1.5.1	Spontaneous symmetry breaking	7
1.5.2	Kink solutions	8
1.6	Complex scalar field theory	8
1.7	$O(N)$ theory	9
1.8	Quantum scalar field theory	9
1.8.1	Feynman Path Integral	10
1.8.2	Renormalization	12
1.9	Scalar Field equation for general metrics	13
1.9.1	Schwarzschild Metric	15
1.9.2	FRW Metric	15
1.9.3	Anti de Sitter space	16
1.9.4	Scalar Field in Anti de Sitter space	17
1.10	Questions	22
2	Scalar Field in AdS space	23
2.1	n -dimensional AdS space	23
2.2	Black hole in Anti de Sitter space	26

3	Scalar Field in Black Hole Anti de Sitter	31
3.1	Introduction	31
3.2	Charged Black Hole in AdS	31
3.3	Zero Temperature	35
3.4	Charged Scalar Field in Charged Black Hole	37
3.4.1	Exeptional Cases	40
3.5	Numerical Approach	41
3.5.1	Adaptive grid size	43
4	Dirac Equation in Black Hole AdS	45
4.1	Introduction	45
4.2	Lorentz Lie Algebra	45
4.2.1	Gamma matrices in n-dimentional space	47
4.2.2	Gamma matrices in Curved space-time	47
4.3	Dirac Equation for curved space-time	49
4.3.1	Diran Equation solution for simple AdS_5 space	50
4.3.2	General Note on Diagonal Mertric which depends on just one variable	53
4.3.3	Dirac equation in Charged Field	54
4.4	solving the Dirac equation	57
4.4.1	Green function	58
4.4.2	Zero Temprature	59
4.4.3	Looking more carefully at zero temprature and field without mass	61
5	Spin 3/2 particles	63
5.1	Introduction	63
5.2	Rarita-Schwinger Formalism	63
5.3	Equation of motion for 3/2 spin particles	64
5.4	RaritaSchwinger equation	65
5.4.1	RaritaSchwinger equation in 4-dimentional Minkowskispac	66
5.5	Spin 3/2 in curved space	68
6	NOT RELATED TO THE PROJECT	71
6.1	SELF NOTES FOR MY RECORD	71
6.1.1	Is the nature Lorentz Invariance ?	71
6.1.2	Black Rings in 5-dimention	71

Quantum Field Theory
©2012 by Arya Farahi

Chapter 1

Classical Scalar Field

1.1 Scalar field theory

In theoretical physics, scalar field theory can refer to a classical or quantum theory of scalar fields. A field which is invariant under any Lorentz transformation is called a "scalar", in contrast to a vector or tensor field. The quanta of the quantized scalar field are spin-zero particles, and as such are bosons.

No fundamental scalar fields have been observed in nature, though the Higgs boson may yet prove the first example. However, scalar fields appear in the effective field theory descriptions of many physical phenomena. An example is the pion, which is actually a "pseudoscalar", which means it is not invariant under parity transformations which invert the spatial directions, distinguishing it from a true scalar, which is parity-invariant. Because of the relative simplicity of the mathematics involved, scalar fields are often the first field introduced to a student of classical or quantum field theory.

In this article, the repeated index notation indicates the Einstein summation convention for summation over repeated indices. The theories described are defined in flat, D-dimensional Minkowski space, with (D-1) spatial dimension and one time dimension and are, by construction, relativistically covariant. The Minkowski space metric, $\eta_{\mu\nu}$, has a particularly simple form: it is diagonal, and here we use the $[-+++]$ sign convention.

1.2 Classical Scalar Field Theory

1.2.1 Linear (free) theory

The most basic scalar field theory is the linear theory. The action for the free relativistic scalar field theory is:

$$\mathcal{S} = \int d^{D-1}x dt \mathcal{L} = \int d^{D-1}x dt \left[\frac{1}{2} \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} m^2 \phi^2 \right] = \int d^{D-1}x dt \left[\frac{1}{2} (-\partial_t \phi)^2 + \right. \quad (1.1)$$

where \mathcal{L} is known as a Lagrangian density. This is an example of a quadratic action, since each of the terms is quadratic in the field, ϕ . The term proportional to m^2 is sometimes known as a mass term, due to its interpretation in the quantized version of this theory in terms of particle mass.

The equation of motion for this theory is obtained by extremizing the action above. It takes the following form, linear in ϕ :

$$\eta^{\mu\nu} \partial_\mu \partial_\nu \phi - m^2 \phi = -\partial_t^2 \phi + \nabla^2 \phi - m^2 \phi = 0 \quad (1.2)$$

Note that this is the same as the Klein-Gordon equation, but that here the interpretation is as a classical field equation, rather than as a quantum mechanical wave equation.

1.2.2 Nonlinear (interacting) theory

The most common generalization of the linear theory above is to add a scalar potential $V(\phi)$ to the equations of motion, where typically, V is a polynomial in ϕ of order 3 or more (often a monomial). Such a theory is sometimes said to be interacting, because the Euler-Lagrange equation is now nonlinear, implying a self-interaction. The action for the most general such theory is

$$\begin{aligned} \mathcal{S} &= \int d^{D-1}x dt \mathcal{L} = \int d^{D-1}x dt \left[\frac{1}{2} \eta^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] \\ &= \int d^{D-1}x dt \left[\frac{1}{2} (-\partial_t \phi)^2 + \frac{1}{2} \delta^{ij} \partial_i \phi \partial_j \phi - \frac{1}{2} m^2 \phi^2 - \sum_{n=3}^{\infty} \frac{1}{n!} g_n \phi^n \right] \quad (1.3) \end{aligned}$$

The $n!$ factors in the expansion are introduced because they are useful in the Feynman diagram expansion of the quantum theory, as described below. The corresponding Euler-Lagrange equation of motion is:

$$\eta^{\mu\nu} \partial_\mu \partial_\nu \phi + V'(\phi) = -\partial_t^2 \phi + \nabla^2 \phi + V'(\phi) = \square \phi + V'(\phi) = 0 \quad (1.4)$$

We can define the new operator, let say box or \square , which is useful when we want to use general relativity ideas in quantum feild theory.

$$\square \phi = \eta^{\mu\nu} \partial_\mu \partial_\nu \phi = -\partial_t^2 \phi + \nabla^2 \phi \quad (1.5)$$

1.3 Dimensional analysis and scaling

Physical quantities in these scalar field theories may have dimensions of length, time or mass, or some combination of the three. However, in a relativistic theory, any quantity t , with dimensions of time, can be 'converted' into a length, $l = ct$, by using the velocity of light, c .

Similarly, any length l is equivalent to an inverse mass, $l = \frac{\hbar}{mc}$, using Planck's constant, \hbar . Heuristically, one can think of a time as a length, or either time or length as an inverse mass. In short, one can think of the dimensions of any physical quantity as defined in terms of just one independent dimension, rather than in terms of all three. This is most often termed the mass dimension of the quantity.

One objection is that this theory is classical, and therefore it is not obvious that Planck's constant should be a part of the theory at all. In a sense this is a valid objection, and if desired one can indeed recast the theory without mass dimensions at all. However, this would be at the expense of making the connection with the quantum scalar field slightly more obscure. Given that one has dimensions of mass, Planck's constant is thought of here as an essentially arbitrary fixed quantity with dimensions appropriate to convert between mass and inverse length.

1.3.1 Scaling Dimension

The classical scaling dimension, or mass dimension, Δ , of ϕ describes the transformation of the field under a rescaling of coordinates:

$$x \rightarrow \lambda x \quad (1.6)$$

$$\phi \rightarrow \lambda^{-\Delta} \phi \quad (1.7)$$

The units of action are the same as the units of \hbar , and so the action itself has zero mass dimension. This fixes the scaling dimension of ϕ to be

$$\Delta = \frac{D-2}{2}. \quad (1.8)$$

1.4 Scale Invariance

There is a specific sense in which some scalar field theories are scale-invariant. While the actions above are all constructed to have zero mass dimension, not all actions are invariant under the scaling transformation

$$x \rightarrow \lambda x \quad (1.9)$$

$$\phi \rightarrow \lambda^{-\Delta} \phi \quad (1.10)$$

The reason that not all actions are invariant is that one usually thinks of the parameters m and g_n as fixed quantities, which are not rescaled under the transformation above. The condition for a scalar field theory to be scale invariant is then quite obvious: all of the parameters appearing in the action should be dimensionless quantities. In other words, a scale invariant theory is one without any fixed length scale (or equivalently, mass scale) in the theory.

For a scalar field theory with D spacetime dimensions, the only dimensionless parameter g_n satisfies $n = \frac{2D}{D-2}$. For example, in $D = 4$ only g_4 is classically dimensionless, and so the only classically scale-invariant scalar field theory in $D = 4$ is the massless ϕ^4 theory. Classical scale invariance normally does not imply quantum scale invariance. See the discussion of the beta function below.

1.4.1 Conformal Invariance

A transformation:

$$x \rightarrow \tilde{x}(x) \quad (1.11)$$

is said to be conformal if the transformation satisfies:

$$\frac{\partial \tilde{x}^\mu}{\partial x^\rho} \frac{\partial \tilde{x}^\nu}{\partial x^\sigma} \eta_{\mu\nu} = \lambda^2(x) \eta_{\rho\sigma} \quad (1.12)$$

for some function $\lambda^2(x)$. The conformal group contains as subgroups the isometries of the metric $\eta_{\mu\nu}$ (the Poincar group) and also the scaling transformations considered above. In fact, the scale-invariant theories in the previous section are also conformally-invariant.

1.5 ϕ^4 theory

Massive ϕ^4 theory illustrates a number of interesting phenomena in scalar field theory.

The Lagrangian density is:

$$\mathcal{L} = \frac{1}{2}(\partial_t \phi)^2 - \frac{1}{2}\delta^{ij}\partial_i \phi \partial_j \phi - \frac{1}{2}m^2 \phi^2 - \frac{g}{4!}\phi^4. \quad (1.13)$$

1.5.1 Spontaneous symmetry breaking

This Lagrangian has a Z_2 symmetry under the transformation $\phi \rightarrow -\phi$.

This is an example of an internal symmetry, in contrast to a space-time symmetry.

If m^2 is positive, the potential $V(\phi) = \frac{1}{2}m^2\phi^2 + \frac{g}{4!}\phi^4$ has a single minimum, at the origin. The solution $\phi = 0$ is clearly invariant under the Z_2 symmetry. Conversely, if m^2 is negative, then one can readily see that the potential $V(\phi) = \frac{1}{2}m^2\phi^2 + \frac{g}{4!}\phi^4$ has two minima. This is known as a double well potential, and the lowest energy states (known as the vacua, in quantum field theoretical language) in such a theory are not invariant under the Z_2 symmetry of the action (in fact it maps each of the two vacua into the other). In this case, the Z_2 symmetry is said to be spontaneously broken.

1.5.2 Kink solutions

The ϕ^4 theory with a negative m^2 also has a kink solution, which is a canonical example of a soliton. Such a solution is of the form:

$$\phi(\vec{x}, t) = \pm \frac{m}{2\sqrt{g}} \tanh\left(\frac{m(x-x_0)}{\sqrt{2}}\right) \quad (1.14)$$

where x is one of the spatial variables (ϕ is taken to be independent of t , and the remaining spatial variables). The solution interpolates between the two different vacua of the double well potential. It is not possible to deform the kink into a constant solution without passing through a solution of infinite energy, and for this reason the kink is said to be stable. For $D > 2$, i.e. theories with more than one spatial dimension, this solution is called a domain wall.

Another well-known example of a scalar field theory with kink solutions is the sine-Gordon theory.

1.6 Complex scalar field theory

In a complex scalar field theory, the scalar field takes values in the complex numbers, rather than the real numbers. The action considered normally takes the form:

$$\mathcal{S} = \int d^{D-1}x dt \mathcal{L} = \int d^{D-1}x dt [\eta^{\mu\nu} \partial_\mu \phi^* \partial_\nu \phi - V(|\phi|^2)] \quad (1.15)$$

This has a $U(1)$ symmetry, whose action on the space of fields rotates $\phi \rightarrow e^{i\alpha} \phi$, for some real phase angle α .

As for the real scalar field, spontaneous symmetry breaking is found if m^2 is negative. This gives rise to a Mexican hat potential which is analogous to the double-well potential in real scalar field theory, but now the choice of vacuum breaks a continuous $U(1)$ symmetry instead of a discrete one. This leads to a Goldstone boson.

1.7 $O(N)$ theory

One can express the complex scalar field theory in terms of two real fields, $\phi^1 = \text{Re}\phi$ and $\phi^2 = \text{Im}\phi$ which transform in the vector representation of the $U(1) = O(2)$ internal symmetry. Although such fields transform as a vector under the internal symmetry, they are still Lorentz scalars. This can be generalised to a theory of N scalar fields transforming in the vector representation of the $O(N)$ symmetry. The Lagrangian for an $O(N)$ -invariant scalar field theory is typically of the form:

$$\mathcal{L} = \frac{1}{2} \eta^{\mu\nu} \partial_\mu \phi \cdot \partial_\nu \phi - V(\phi \cdot \phi) \quad (1.16)$$

using an appropriate $O(N)$ -invariant inner product.

1.8 Quantum scalar field theory

In quantum field theory, the fields, and all observables constructed from them, are replaced by quantum operators on a Hilbert space. This Hilbert space is built on a vacuum state, and dynamics are governed by a Hamiltonian, a positive operator which annihilates the vacuum. A construction of a quantum scalar field theory may be found in the canonical quantization article, which uses canonical commutation relations among the fields as a basis for the construction. In brief, the basic variables are the field ϕ and its canonical momentum. Both fields are Hermitian. At spatial points \vec{x}, \vec{y} at equal times, the canonical commutation relations are given by

$$[\phi(\vec{x}), \phi(\vec{y})] = [\pi(\vec{x}), \pi(\vec{y})] = 0, \text{ and } [\phi(\vec{x}), \pi(\vec{y})] = i\delta(\vec{x} - \vec{y}) \quad (1.17)$$

and the free Hamiltonian is:

$$H = \int d^3x \left[\frac{1}{2} \pi^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{m^2}{2} \phi^2 \right] \quad (1.18)$$

A spatial Fourier transform leads to momentum space fields:

$$\tilde{\phi}(\vec{k}) = \int d^3x e^{-i\vec{k} \cdot \vec{x}} \phi(\vec{x}), \text{ and } \tilde{\pi}(\vec{k}) = \int d^3x e^{-i\vec{k} \cdot \vec{x}} \pi(\vec{x}) \quad (1.19)$$

which are used to define annihilation and creation operators:

$$a(\vec{k}) = \left(E\tilde{\phi}(\vec{k}) + i\tilde{\pi}(\vec{k}) \right), a^\dagger(\vec{k}) = \left(E\tilde{\phi}(\vec{k}) - i\tilde{\pi}(\vec{k}) \right), \quad (1.20)$$

where $E = \sqrt{k^2 + m^2}$. These operators satisfy the commutation relations:

$$[a(\vec{k}_1), a(\vec{k}_2)] = [a^\dagger(\vec{k}_1), a^\dagger(\vec{k}_2)] = 0, [a(\vec{k}_1), a^\dagger(\vec{k}_2)] = (2\pi)^3 2E \delta(\vec{k}_1 - \vec{k}_2) \quad (1.21)$$

The state $|0\rangle$ annihilated by all of the operators is identified as the bare vacuum, and a particle with momentum \vec{k} is created by applying $a^\dagger(\vec{k})$ to the vacuum. Applying all possible combinations of creation operators to the vacuum constructs the Hilbert space. This construction is called Fock space. The vacuum is annihilated by the Hamiltonian:

$$H = \int \frac{d^3k}{(2\pi)^3} \frac{1}{2} a^\dagger(\vec{k}) a(\vec{k}) \quad (1.22)$$

where the zero-point energy has been removed by Wick ordering.

Interactions can be included by adding an interaction Hamiltonian. For a ϕ^4 theory, this corresponds to adding a Wick ordered term $g : \phi^4 : / 4!$ to the Hamiltonian, and integrating over x . Scattering amplitudes may be calculated from this Hamiltonian in the interaction picture. These are constructed in perturbation theory by means of the Dyson series, which gives the time-ordered products, or n-particle Green's functions $\langle 0 | \mathcal{T} \{ \phi(x_1) \cdots \phi(x_n) \} | 0 \rangle$ as described in the Dyson series article. The Green's functions may also be obtained from a generating function that is constructed as a solution to the Schwinger-Dyson equation.

1.8.1 Feynman Path Integral

The Feynman diagram expansion may be obtained also from the Feynman path integral formulation. The time ordered vacuum expectation values of polynomials in ϕ , known as the n-particle Green's functions, are constructed by integrating over all possible fields, normalized by the vacuum expectation value with no external fields:

$$\langle 0 | \mathcal{T} \{ \phi(x_1) \cdots \phi(x_n) \} | 0 \rangle = \frac{\int \mathcal{D}\phi \phi(x_1) \cdots \phi(x_n) e^{i \int d^4x \left(\frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{m^2}{2} \phi^2 - \frac{g}{4!} \phi^4 \right)}}{\int \mathcal{D}\phi e^{i \int d^4x \left(\frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{m^2}{2} \phi^2 - \frac{g}{4!} \phi^4 \right)}} \quad (1.23)$$

All of these Green's functions may be obtained by expanding the exponential in $J(x)\phi(x)$ in the generating function

$$Z[J] = \int \mathcal{D}\phi e^{i \int d^4x \left(\frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{m^2}{2} \phi^2 - \frac{g}{4!} \phi^4 + J\phi \right)} = Z[0] \sum_{n=0}^{\infty} \frac{i^n J(x_1) \cdots J(x_n)}{n!} \langle 0 | \mathcal{T} \{ \phi(x_1) \cdots \phi(x_n) \} | 0 \rangle \quad (1.24)$$

A Wick rotation may be applied to make time imaginary. Changing the signature to $(++++)$ then turns the Feynman integral into a statistical mechanics partition function in Euclidean space:

$$Z[J] = \int \mathcal{D}\phi e^{-\int d^4x \left(\frac{1}{2} (\nabla\phi)^2 + \frac{m^2}{2} \phi^2 + \frac{g}{4!} \phi^4 + J\phi \right)} \quad (1.25)$$

Normally, this is applied to the scattering of particles with fixed momenta, in which case, a Fourier transform is useful, giving instead:

$$\tilde{Z}[\tilde{J}] = \int \mathcal{D}\tilde{\phi} e^{-\int d^4p \left(\frac{1}{2} (p^2 + m^2) \tilde{\phi}^2 + \frac{\lambda}{4!} \tilde{\phi}^4 - \tilde{J}\tilde{\phi} \right)} \quad (1.26)$$

The standard trick to evaluate this functional integral is to write it as a product of exponential factors, schematically:

$$\tilde{Z}[\tilde{J}] \sim \int \mathcal{D}\tilde{\phi} \prod_p \left[e^{-(p^2 + m^2) \tilde{\phi}^2 / 2} e^{-g \tilde{\phi}^4 / 4!} e^{\tilde{J}\tilde{\phi}} \right] \quad (1.27)$$

The second two exponential factors can be expanded as power series, and the combinatorics of this expansion can be represented graphically. The integral with $\lambda = 0$ can be treated as a product of infinitely many elementary Gaussian integrals, and the result may be expressed as a sum of Feynman diagrams, calculated using the following Feynman rules:

- * Each field $\tilde{\phi}(p)$ in the n -point Euclidean Green's function is represented by an external line (half-edge) in the graph, and associated with momentum p .
- * Each vertex is represented by a factor $-g$.
- * At a given order g^k , all diagrams with n external lines and k vertices are constructed such that the momenta flowing into each vertex is zero. Each internal

line is represented by a propagator $1/(q^2 + m^2)$, where q is the momentum flowing through that line.

* Any unconstrained momenta are integrated over all values.

* The result is divided by a symmetry factor, which is the number of ways the lines and vertices of the graph can be rearranged without changing its connectivity.

* Do not include graphs containing "vacuum bubbles", connected subgraphs with no external lines.

The last rule takes into account the effect of dividing by $\tilde{Z}[0]$. The Minkowski-space Feynman rules are similar, except that each vertex is represented by $-ig$, while each internal line is represented by a propagator $\frac{1}{q^2 - m^2 + i\epsilon}$, where the term represents the small Wick rotation needed to make the Minkowski-space Gaussian integral converge.

1.8.2 Renormalization

The integrals over unconstrained momenta, called "loop integrals", in the Feynman graphs typically diverge. This is normally handled by renormalization, which is a procedure of adding divergent counter-terms to the Lagrangian in such a way that the diagrams constructed from the original Lagrangian and counter-terms is finite. A renormalization scale must be introduced in the process, and the coupling constant and mass become dependent upon it.

The dependence of a coupling constant g on the scale λ is encoded by a beta function, $\beta(g)$, defined by the relation:

$$\beta(g) = \lambda \frac{\partial g}{\partial \lambda} \quad (1.28)$$

This dependence on the energy scale is known as the running of the coupling parameter, and theory of this kind of scale-dependence in quantum field theory is described by the renormalization group.

Beta-functions are usually computed in an approximation scheme, most commonly perturbation theory, where one assumes that the coupling constant is small. One can then make an expansion in powers of the coupling parameters and truncate the higher-order terms (also known as higher loop contributions, due to the number of loops in the corresponding Feynman graphs).

The beta-function at one loop (the first perturbative contribution) for the ϕ^4 theory is:

$$\beta(g) = \frac{3}{16\pi^2}g^2 + O(g^3) \quad (1.29)$$

The fact that the sign in front of the lowest-order term is positive suggests that the coupling constant increases with energy. If this behavior persists at large couplings, this would indicate the presence of a Landau pole at finite energy, or quantum triviality. The question can only be answered non-perturbatively, since it involves strong coupling.

A quantum field theory is trivial when the running coupling, computed through its beta function, goes to zero when the cutoff is removed. Consequently, the propagator becomes that of a free particle and the field is no longer interacting. Alternatively, the field theory may be interpreted as an effective theory, in which the cutoff is not removed, giving finite interactions but leading to a Landau pole at some energy scale. For a ϕ^4 interaction, Michael Aizenman proved that the theory is indeed trivial for space-time dimension $D \geq 5$. For $D = 4$ the triviality has yet to be proven rigorously, but lattice computations have confirmed this. (See Landau pole for details and references.) This fact is relevant as the Higgs field, for which triviality bounds are used to set limits on the Higgs mass, based on the new physics must enter at a higher scale (perhaps the Planck scale) to prevent the Landau pole from being reached.

1.9 Scalar Field equation for general metrics

Until here we learned how to deal with scalar field equation in Minkowski-space. Now we want to focus more about scalar field equation in curved spaces, and solve the equation for some well known curved spaces which has specific metrics, such as Schwarzschild, FRW, and Anti de Sitter metric.

The metric tensor is such an important object in curved space that is given a new symbol, let say $g_{\mu\nu}$, while $\eta_{\mu\nu}$ specifically be used for Minkowski space as a metric symbol. $g_{\mu\nu}$ should obey some rules to be considered as metric. First of all it should be a symmetric (0,2) matrix, or more generally tensor. Second, it usually taken non-degenerate, it means that its determinant does not vanish, $g = |g_{\mu\nu}|$. If the determinant is not zero, then obviously the matrix, tensor, has its own inverse, which is unique:

$$g^{\mu\nu} g_{\nu\sigma} = \delta_\sigma^\nu \quad (1.30)$$

In minkowski-space we defined the line element:

$$ds = \eta_{\mu\nu} dx^\mu dx^\nu \quad (1.31)$$

Respectively, it is possible to define the line element in the same way, which is more general and it can be use for any kind of curved space or flat space:

$$ds = g_{\mu\nu} dx^\mu dx^\nu \quad (1.32)$$

Based on the new metric, we want to developpe the new scalar field equation of motion. Lagrangian has the form of,

$$\mathcal{L} = \sqrt{-|g_{\mu\nu}|} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} m^2 \phi^2 \right] \quad (1.33)$$

The next step, we have to derive the equation of motion from Lagrangian, equation 1.33,

$$\begin{aligned} 0 &= \frac{\partial \mathcal{L}}{\partial \phi} - \frac{\partial}{\partial \mu} \frac{\mathcal{L}}{\partial \phi_\mu} \rightarrow \\ 0 &= \frac{\partial}{\partial \mu} \sqrt{-|g_{\mu\nu}|} g^{\mu\nu} \frac{\partial \phi}{\partial \nu} - \sqrt{-|g_{\mu\nu}|} m^2 \phi \end{aligned} \quad (1.34)$$

So the motion equation for scalar field has the form of,

$$0 = \left[\frac{1}{\sqrt{-|g_{\mu\nu}|}} \partial_\mu \sqrt{-|g_{\mu\nu}|} g^{\mu\nu} \partial_\nu - m^2 \right] \phi \quad (1.35)$$

For few sections we just introduce some famous metrics and their tensors and after that we would try to solve the equation of motion for scalar field, equation 2.3.

1.9.1 Schwarzschild Metric

$$ds^2 = -V(r)dt^2 + V(r)^{-1}dr^2 + r^2d\Omega_{D-2}^2 \quad (1.36)$$

For example the Schwartzschild black hole of mass M is described by the metric:

$$ds^2 = -\left(1 - \frac{2M}{r}\right)dt^2 + \left(1 - \frac{2M}{r}\right)^{-1}dr^2 + r^2d\Omega^2 \quad (1.37)$$

The Schwarzschild metric describes the spacetime around a spherically symmetric body, such as a planet, or a black hole. With coordinates $(x^0, x^1, x^2, x^3) = (ct, r, \theta, \phi)$, we can write the metric as:

$$(g_{\mu\nu}) = \begin{pmatrix} -\left(1 - \frac{2GM}{rc^2}\right) & 0 & 0 & 0 \\ 0 & \left(1 - \frac{2GM}{rc^2}\right)^{-1} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix} \quad (1.38)$$

The interesting point about the metric of Schwarzschild is that if one puts the mass of black hole equal to zero the metric would become similar to Minkowski's metric. Because when the mass is equal to zero it means that there is no black hole so the metric should be the same as Minkowski's metric.

1.9.2 FRW Metric

Friedmann-Rabertson-Walker, or simply FRW, metric describe homogeneous, isotropic universe, including, to a good degree of approximation, the portion we have seen of our own universe. The metric, in spherical coordinate, has the form of:

$$ds^2 = -dt^2 + a(t)^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right) \quad (1.39)$$

and the tensor of FRW metric has the form of:

$$(g_{\mu\nu}) = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & \frac{a(t)^2}{1 - k^2 r^2} & 0 & 0 \\ 0 & 0 & a(t)^2 r^2 & 0 \\ 0 & 0 & 0 & a(t)^2 r^2 \sin^2(\theta) \end{pmatrix} \quad (1.40)$$

In these equations $a(t)$ is the scale factor, cosmic scale factor or sometimes the Robertson-Walker scale factor parameter of the Friedmann equations is a function of time which represents the relative expansion of the universe. Modern observations suggests that the universe is expanding and introducing $a(t)$ is the way the help us to consider the expansion of our universe in our equations. And the relation of scale factor and Hubble parameter has the form of:

$$H = \frac{\dot{a}(t)}{a(t)} \quad (1.41)$$

1.9.3 Anti de Sitter space

In mathematics and physics, n-dimensional anti de Sitter space, sometimes written AdS_n , is a maximally symmetric Lorentzian manifold with constant negative scalar curvature. It is the Lorentzian analogue of n-dimensional hyperbolic space, just as Minkowski space and de Sitter space are the analogues of Euclidean and elliptical spaces respectively.

It is best known for its role in the AdS/CFT correspondence.

In the language of general relativity, anti de Sitter space is a maximally symmetric, vacuum solution of Einstein's field equation with a negative (attractive) cosmological constant Λ (corresponding to a negative vacuum energy density and positive pressure).

In mathematics, anti de Sitter space is sometimes defined more generally as a space of arbitrary signature (p, q) . Generally in physics only the case of one timelike dimension is relevant. Because of differing sign conventions, this may correspond to a signature of either $(n1, 1)$ or $(1, n1)$.

A coordinate patch covering part of the space gives the half-space coordinatization of anti de Sitter space. The metric for this patch is:

$$ds^2 = \frac{1}{z^2} (-dt^2 + dx^2 + dy^2 + dz^2) \quad (1.42)$$

We easily see that this metric is conformally equivalent to a flat half-space Minkowski spacetime.

The constant time slices of this coordinate patch are hyperbolic spaces in the Poincar half-plane metric. In the limit as $z = 0$, this half-space metric reduces

to a Minkowski metric $dz^2 = dt^2 - dx^2 - dy^2$; thus, the anti-de Sitter space contains a conformal Minkowski space at infinity ("infinity" having y-coordinate zero in this patch).

In AdS space time is periodic, and the universal cover has non-periodic time. The coordinate patch above covers half of a single period of the spacetime.

Because the conformal infinity of AdS is timelike, specifying the initial data on a spacelike hypersurface would not determine the future evolution uniquely (i.e. deterministically) unless there are boundary conditions associated with the conformal infinity.

The "half-space" region of anti de Sitter space and its boundary.

Another commonly used coordinate system which covers the entire space is given by the coordinates t , $r \geq 0$ and the hyperpolar coordinates α , θ and ϕ .

$$\begin{aligned} ds^2 &= -(k^2 r^2 + 1) dt^2 + \frac{1}{k^2 r^2 + 1} dr^2 + r^2 d\Omega^2 \\ &= -(k^2 r^2 + 1) dt^2 + \frac{1}{k^2 r^2 + 1} dr^2 + r^2 (d\theta^2 + \sin^2(\theta) d\phi^2) \end{aligned} \quad (1.43)$$

The image on the right represents the "half-space" region of anti deSitter space and its boundary. The interior of the cylinder corresponds to Anti de Sitter spacetime, while its cylindrical boundary corresponds to its conformal boundary. The green shaded region in the interior corresponds to the region of AdS covered by the half-space coordinates and it is bounded by two null, aka lightlike, geodesic hyperplanes; the green shaded area on the surface corresponds to the region of conformal space covered by Minkowski space.

The green shaded region covers half of the AdS space and half of the conformal spacetime; the left ends of the green discs will touch in the same fashion as the right ends.

1.9.4 Scalar Field in Anti de Sitter space

Now we want to find the general solution of the equation of motion, 2.3, in Anti de Sitter space, from now AdS, space. First we focus on general solution in Cartesian coordinate system and after that we figure out how to find the general solution of equation of motion for AdS space with Spherical coordinate basis.

Equation 1.42 shows the metric of AdS space in Cartesian coordinate system. Based on equation 1.42 the metric tensor has the form of:

$$(g_{\mu\nu}) = \begin{pmatrix} -\frac{1}{z^2} & 0 & 0 & 0 \\ 0 & \frac{1}{z^2} & 0 & 0 \\ 0 & 0 & \frac{1}{z^2} & 0 \\ 0 & 0 & 0 & \frac{1}{z^2} \end{pmatrix} \quad (1.44)$$

Determinant of tensor $g_{\mu\nu}$, equation 1.44, is $|g_{\mu\nu}| = -\frac{1}{z^8}$. So the for the scalar fields and in AdS space with cartesian coordinate system the equation of motion, equation 2.3, would be,

$$\left[-z^2 \frac{\partial^2}{\partial t^2} + z^2 \frac{\partial^2}{\partial x^2} + z^2 \frac{\partial^2}{\partial y^2} + z^2 \frac{\partial^2}{\partial z^2} - 2z \frac{\partial}{\partial z} - m^2 \right] \phi = 0 \quad (1.45)$$

and then we can divide both side of equaion by z^2 ,

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} - \frac{2}{z} \frac{\partial \phi}{\partial z} - \frac{m^2}{z^2} \phi = \frac{\partial^2 \phi}{\partial t^2} \quad (1.46)$$

Now, we can use seperation of variables method to find the solution of equation of motion in AdS space with cartesian coordinate system, equation 1.46. Based on this method, $\phi(t, x, y, z) = T(r)X(x)Y(y)Z(z)$, and by substituting the new function in our equation, equation 1.46, and then divide both side of equation by ϕ , $T(r)X(x)Y(y)Z(z)$, we would have:

$$\frac{\frac{\partial^2 X(x)}{\partial x^2}}{X(x)} + \frac{\frac{\partial^2 Y(y)}{\partial y^2}}{Y(y)} + \frac{\frac{\partial^2 Z(z)}{\partial z^2} - \frac{2}{z} \frac{\partial Z(z)}{\partial z} - \frac{m^2}{z^2} Z(z)}{Z(z)} = \frac{\frac{\partial^2 T(t)}{\partial t^2}}{T(t)} \quad (1.47)$$

There are four terms, three spatial terms and one time term, which that each ones depend just on one varibale. So we can easily conclude that, as you may learned on basic courses like partial differential equation, each term should be constant. Lest say C_1 , C_2 , and C_3 for spatial terms and C for time term. Then we have to solve each term separately fo find the answers. This equation is very similar to equation of wave, like what we have in electrodynamics. The differenc is that one of our spatial terms is not similar what we had before on wave problems. But we use the same method to find the solution for this partial differencial equation. Let's do the calculations.

For time term, it would be like,

$$C = \frac{\frac{\partial^2 T(t)}{\partial t^2}}{T(t)} \quad (1.48)$$

and then,

$$T(t) = a_1 e^{i\omega t} + a_2 e^{i\omega' t} \quad (1.49)$$

which ω and ω' can be complex number. Here, for simplicity, we assume that $T(t) = a_1 e^{i\omega t}$, then the constant, C , would be equal to $-\omega^2$ and a_1 is just a normalization factor. For the next step we try to find the solution of spatial terms. The first two spatial terms, $X(x)$ and $Y(y)$, are the same as time term and we can solve them like time term, so for this two term we have:

$$\begin{aligned} X(x) &= b_1 e^{ik_1 x} + b_2 e^{k_{12} x} \\ Y(y) &= b_3 e^{ik_2 y} + b_4 e^{k_{22} y} \end{aligned} \quad (1.50)$$

and like the time term for simplicity we can assume that the second terms are zero. So, C_1 and C_2 are k_1^2 and k_2^2 respectively. b_1 and b_3 are just normalization factors. The most important part of the equation is the last spatial time which make difference between Minkowski's space and AdS space. The differential equation for the second term has the form of,

$$\frac{\partial^2 Z(z)}{\partial z^2} - \frac{2}{z} \frac{\partial Z(z)}{\partial z} - (C_3 + \frac{m^2}{z^2}) Z(z) = 0 \quad (1.51)$$

and C_3 is a constant that related with C_2 , C_3 , and C with the following relationship,

$$C_3 = C - C_1 - C_2 = -[\omega^2 - k_1^2 - k_2^2] = -k^2 \quad (1.52)$$

If we multiply equation by z^2 then it becomes famous Bessel differential equation, which its answers are Bessel functions.

$$z^2 \frac{\partial^2 Z(z)}{\partial z^2} - 2z \frac{\partial Z(z)}{\partial z} - (z^2 C_3 + m^2) Z(z) = 0 \quad (1.53)$$

and the answer of this equation has the form of,

$$Z(z) = d_1 z^{\frac{3}{2}} J_m(\sqrt{k^2} z) + d_2 z^{\frac{3}{2}} Y_m(\sqrt{k^2} z) \quad (1.54)$$

That d_1 and d_2 are just normalization factors. So the complete solution of motion equation for scalar field in AdS space with cartesian coordinate basis has the form of,

$$\phi(t, x, y, z) = A_1 e^{ik_\mu x_\mu} z^{\frac{3}{2}} J_m(\sqrt{k^2} z) + A_2 e^{ik_\mu x_\mu} z^{\frac{3}{2}} Y_m(\sqrt{k^2} z) \quad (1.55)$$

That A_1 and A_2 are just normalization factors and $k_\mu x_\mu$ is $\omega t + k_1 x + k_2 y$ and ω , k_1 , and k_2 can be complex numbers.

Now, lets use the same method to find the solution of motion equation for scalar field in AdS space with Spherical coordinate basis. Equation 1.43 shows the metric of AdS space in Spherica coordinate system. Based on equation 1.43 the metric tensor is:

$$(g_{\mu\nu}) = \begin{pmatrix} -(k^2 r^2 + 1) & 0 & 0 & 0 \\ 0 & \frac{1}{k^2 r^2 + 1} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin(\theta)^2 \end{pmatrix} \quad (1.56)$$

Determinant of tensor $g_{\mu\nu}$, equation 1.56, is $|g_{\mu\nu}| = -r^4 \sin(\theta)^2$. So the equation of motion for scalar field in AdS space with spherical coordinate system basis, equation 2.3, has the form of,

$$-\frac{1}{(k^2 r^2 + 1)} \frac{\partial^2 \phi}{\partial t^2} + (k^2 r^2 + 1) \frac{\partial^2 \phi}{\partial r^2} + (4k^2 r + \frac{2}{r}) \frac{\partial \phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \Omega^2} - m^2 \phi = 0$$

In case of spherical symmetry we can separate the angular varibales from radial and time variables, r and t . In basic quantum mechanics courses and partial differential courses it is proved that for spherical symmetry the eigenvalue of angular part is: $l(l+1)$. So we can divide ϕ into two part, angular part and time-radial part, $\phi(t, r, \theta, \Phi) = X(t, r) Y_l^m(\theta, \Phi)$, also is is possible to show that $\frac{\partial^2 Y_l^m(\theta, \Phi)}{\partial \Omega^2} = l(l+1)$. On the other hand we can assume that function of $X(t, r)$ are sparetable into to function of $R(r)$ and $T(t)$. Then simply we can rewrite equation of motion,

$$\begin{aligned}
& -\frac{R(r)}{(k^2 r^2 + 1)} \frac{\partial^2 T(t)}{\partial t^2} + (k^2 r^2 + 1) T(t) \frac{\partial^2 R(r)}{\partial r^2} \\
& + (4k^2 r + \frac{2}{r}) T(t) \frac{\partial R(r)}{\partial r} + (\frac{l(l+1)}{r^2} - m^2) R(r) T(t) = 0
\end{aligned} \tag{1.57}$$

or,

$$\begin{aligned}
& (k^2 r^2 + 1)^2 \frac{\partial^2 R(r)}{\partial r^2} + (k^2 r^2 + 1) (4k^2 r + \frac{2}{r}) \frac{\partial R(r)}{\partial r} \\
& + \frac{l(l+1)}{r^2} + (k^2 l(l+1) - m^2) = \frac{\partial^2 T(t)}{T(t)}
\end{aligned} \tag{1.58}$$

Because one side of the equation is function of time and the other side is function of r then, for having solution, both side should be constant, like C . The solution for time term is:

$$T(t) = A_1 e^{i\omega t} + A_2 e^{i\omega' t} \tag{1.59}$$

Which for simplicity we assume the the second term is zero then, we have $T(t) = A_1 e^{i\omega t}$ that A_1 is a normalization factor, and ω can be a complex number. On the other hand, our constant, C , would be $-\omega^2$. So differential equation for radial term is,

$$\begin{aligned}
& (k^2 r^2 + 1)^2 \frac{\partial^2 R(r)}{\partial r^2} + (k^2 r^2 + 1) (4k^2 r + \frac{2}{r}) \frac{\partial R(r)}{\partial r} \\
& + \left[\frac{l(l+1)}{r^2} + (k^2 l(l+1) + \omega^2 - m^2) \right] R(r) = 0
\end{aligned} \tag{1.60}$$

We can solve this differential equation with series method. In series method we assume that, $R(r) = \sum_{n=-\infty}^{+\infty} a_n r^n$, and all n 's should be integer. First we substitute a_0 in our differential equation and we get $a_0 = \text{constant}$. We do the same thing for $n > 0$ and we get $a_n = 0$, for $n > 0$, because based on boundary condition in large radius the function should approaches to zero, based on this argument all for positive n 's we should get zero. Now let's focus on $n < 0$. By substituting $\sum a_{-n} r^{-n}$ in equation 1.60 we get,

$$\begin{aligned}
0 = \sum_{n=0}^{\infty} & a_{-n}n(n+1)k^4r^{-n+2} + a_{-n}2k^2n(n+1)r^{-n} + a_{-n}n(n+1)r^{-n-2} \\
& - a_{-n}4nk^2r^{-n+2} - a_{-n}6nk^2r^{-n} - a_{-n}nr^{-n-2} \\
& - a_{-n}l(l+1)k^4r^{-n-2} + a_{-n} \left[k^2l(l+1) + \omega^2 - m^2 \right] r^{-n}
\end{aligned} \tag{1.61}$$

Then the coefficient of each r^n should be zero, so we have,

$$\begin{aligned}
0 = & a_{-n-2}(n+2)(n+3)k^4r^{-n} + a_{-n}2k^2n(n+1)r^{-n} + a_{-n}(n-2)(n-1)r^{-n} \\
& - a_{-n-2}4(n+2)k^2r^{-n} - a_{-n}6nk^2r^{-n} - a_{-n}(n-2)r^{-n} \\
& - a_{-n+2}l(l+1)k^4r^{-n} + a_{-n} \left[k^2l(l+1) + \omega^2 - m^2 \right] r^{-n}
\end{aligned} \tag{1.62}$$

Finally, we can find all a_n 's, if we have boundari condition,

$$a_{-n-2} = \frac{a_{-n} \left[(-2n^2 + 4n - l(l+1))k^2 + m^2 - \omega^2 \right] + a_{-n+2} \left[l(l+1) - n^2 + 3n - 2 \right]}{(n^2 + n - 2)k^4} \tag{1.63}$$

1.10 Questions

1. Derive the partial differential equation for equation of motion in Schwarzschild space with spherical coordinate basis.

2. Derive the partial differential equation for equation of motion in FRW space with spherical coordinate basis.

Chapter 2

Scalar Field in AdS space

2.1 n -dimensional AdS space

In chapter 1 we learned how to solve the scalar field in AdS space in cartesian and spherical coordinate basis. In this chapter, we are looking into this problem further and we discuss about the different features of AdS space in detail.

The metric of n -dimensional AdS space for m timelike dimension and $n - m$ spacelike dimension has the form of,

$$ds^2 = \frac{1}{z^2} \left[- \sum_{i=1}^m dt_i^2 + \sum_{i=1}^{n-m} dx_i^2 \right] \quad (2.1)$$

Which in this equation we assume that x_n would be z coordinate, so the metric tensor has the form of,

$$g_{\mu\nu} = \frac{1}{z^2} \begin{pmatrix} \begin{pmatrix} -1 & 0 & \cdots & 0 \\ 0 & -1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -1 \end{pmatrix}_{m \times m} & 0 \\ 0 & \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}_{(n-m) \times (n-m)} \end{pmatrix} \quad (2.2)$$

And the determinant of this tensor is $|g_{\mu\nu}| = \frac{(-1)^m}{z^n}$. Let's use equation of motion for scalar scalar, equation 2.3 which was introduced in chapter 2, to derive the differential equation of evolution of field in n -dimensional AdS space,

$$0 = \left[\frac{1}{\sqrt{-|g_{\mu\nu}|}} \partial_\mu \sqrt{-|g_{\mu\nu}|} g^{\mu\nu} \partial_\nu - m^2 \right] \phi \quad (2.3)$$

$$\left[-z^2 \sum_{i=1}^m \frac{\partial^2}{\partial t_i^2} + z^2 \sum_{i=1}^{n-m-1} \frac{\partial^2}{\partial x_i^2} + z^2 \frac{\partial^2}{\partial z^2} + (2-n)z \frac{\partial}{\partial z} - m^2 \right] \phi = 0 \quad (2.4)$$

and then we can divide both side of equation by z^2 ,

$$\sum_{i=1}^{n-m-1} \frac{\partial^2 \phi}{\partial x_i^2} + \frac{\partial^2 \phi}{\partial z^2} + \frac{2-n}{z} \frac{\partial \phi}{\partial z} - \frac{m^2}{z^2} \phi = \sum_{i=1}^m \frac{\partial^2 \phi}{\partial t_i^2} \quad (2.5)$$

As we saw in chapter 1 if we use separation of variables method then there is a simple solution for time and all spatial terms, except z term for Z spatial coordinate. The time and spatial, except z , part of field ϕ has the form of,

$$\phi(x_\mu) = e^{ik \cdot x} Z(z) = e^{ik_\mu x_\mu} Z(z) \quad (2.6)$$

By substituting $\phi(x_\mu)$ in equation 2.5 we have,

$$\frac{\partial^2 Z(z)}{\partial z^2} + \frac{2-n}{z} \frac{\partial Z(z)}{\partial z} + \left[k^2 - \frac{m^2}{z^2} \right] Z(z) = 0 \quad (2.7)$$

Let's say $Z(z) = z^\alpha f(z)$, then $\frac{\partial Z(z)}{\partial z} = \alpha z^{\alpha-1} f(z) + z^\alpha \frac{\partial f(z)}{\partial z}$ and $\frac{\partial^2 Z(z)}{\partial z^2} = \alpha(\alpha-1)z^{\alpha-2} f(z) + 2\alpha z^{\alpha-1} \frac{\partial f(z)}{\partial z} + z^\alpha \frac{\partial^2 f(z)}{\partial z^2}$. Then by substituting these three equation in equation 2.7 we have,

$$\begin{aligned} \alpha(\alpha-1)z^{\alpha-2} f(z) + 2\alpha z^{\alpha-1} \frac{\partial f(z)}{\partial z} + z^\alpha \frac{\partial^2 f(z)}{\partial z^2} \\ + \frac{2-n}{z} \left[\alpha z^{\alpha-1} f(z) + z^\alpha \frac{\partial f(z)}{\partial z} \right] + \left[k^2 - \frac{m^2}{z^2} \right] z^\alpha f(z) = 0 \end{aligned} \quad (2.8)$$

And then,

$$z^2 \frac{\partial^2 f(z)}{\partial z^2} + (2\alpha + 2 - n)z \frac{\partial f(z)}{\partial z} + [k^2 z^2 - (m^2 + \alpha(n-1-\alpha))] f(z) = 0 \quad (2.9)$$

By choosing $\alpha = \frac{n-1}{2}$, we get the Bessel differential equation and the general answer has the form of,

$$\begin{aligned} \phi(x_i, z) = & A_1 e^{(ik_\mu x_\mu)} z^{\frac{n-1}{2}} J_\nu(kz) \\ & + A_2 e^{(ik_\mu x_\mu)} z^{\frac{n-1}{2}} Y_\nu(kz) \quad , \quad \nu = \sqrt{m^2 + \frac{(n-1)^2}{4}} \end{aligned} \quad (2.10)$$

Which k_μ 's can be complex numbers and A_1 and A_2 are just normalization facors which depends on boundary condition.

Let's see what happened for our field when it approached to $z = 0$ and $z \rightarrow \infty$. Bsed of definition of Bessel function in the Literature,

$$z^{\frac{n-1}{2}} J_\nu(kz) = z^{\frac{n-1}{2}} \sum_{i=0}^{\infty} \frac{(-1)^i}{i! \Gamma(i + \nu + 1)} \left(\frac{k}{2} z\right)^{(2i+\nu)} \quad (2.11)$$

and,

$$z^{\frac{n-1}{2}} Y_\nu(kz) = z^{\frac{n-1}{2}} \frac{J_\nu(kz) \cos(\nu\pi) - J_{-\nu}(z)}{\sin(\nu\pi)} \quad (2.12)$$

Then,

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} J_\nu(kz) = z^{\frac{n-1}{2} + \nu} \times \frac{k^\nu}{2^\nu \Gamma(\nu + 1)} \quad , \quad \nu = \sqrt{m^2 + \frac{n^2 - 2n + 1}{4}} \quad (2.13)$$

and,

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} Y_\nu(kz) = -z^{\frac{n-1}{2} - \nu} \times \frac{\Gamma(\nu) 2^\nu}{k^\nu \pi} \quad , \quad \nu = \sqrt{m^2 + \frac{n^2 - 2n + 1}{4}} \quad (2.14)$$

or simply,

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} J_\nu(kz) = C z^{\frac{n-1}{2} + \sqrt{m^2 + \frac{(n+1)^2}{4}}} \quad (2.15)$$

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} Y_\nu(kz) = C' z^{\frac{n-1}{2} - \sqrt{m^2 + \frac{(n-1)^2}{4}}} \quad (2.16)$$

For case $n = 4$, it would be,

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} J_\nu(kz) = C z^{\frac{3}{2} + \sqrt{m^2 + \frac{9}{4}}} \quad (2.17)$$

and,

$$\lim_{z \rightarrow 0} z^{\frac{n-1}{2}} Y_V(kz) = C' z^{\frac{3}{2} - \sqrt{m^2 + \frac{9}{4}}} \quad (2.18)$$

2.2 Black hole in Anti de Sitter space

In theoretical physics, an AdS black hole is a black hole solution of general relativity or its extensions which represents an isolated massive object, but with a negative cosmological constant. Such a solution asymptotically approaches Anti de Sitter space at spatial infinity, and is a generalization of the Kerr vacuum solution, which asymptotically approaches Minkowski spacetime at spatial infinity.

In $3 + 1$ dimensions, the metric is given by,

$$ds^2 = - \left(k^2 r^2 + 1 - \frac{C}{r} \right) dt^2 + \frac{1}{k^2 r^2 + 1 - \frac{C}{r}} dr^2 + r^2 d\Omega^2 \quad (2.19)$$

where t is the time coordinate, r is the radial coordinate, Ω are the polar coordinates, C is a constant and k is the AdS curvature.

In general, in $d + 1$ dimensions, the metric is given by,

$$ds^2 = - \left(k^2 r^2 + 1 - \frac{C}{r^{d-2}} \right) dt^2 + \frac{1}{k^2 r^2 + 1 - \frac{C}{r^{d-2}}} dr^2 + r^2 d\Omega^2 \quad (2.20)$$

According to the AdS/CFT correspondence, if gravity were quantized, an AdS black hole would be dual to a thermal state on the conformal boundary. In the context of say, AdS/QCD, this would correspond to the deconfinement phase of the quark-gluon plasma.

Also we can write the black hole's metric in $d + 1$ dimension AdS space in another format which may be more useful,

$$ds^2 = \frac{u^2}{L^2} \left[- \left(1 - \left(\frac{u_0}{u} \right)^d \right) dt^2 + \sum_{i=1}^{d-1} dx_i^2 \right] + \frac{L^2 du^2}{u^2 \left(1 - \left(\frac{u_0}{u} \right)^d \right)} \quad (2.21)$$

From now we assume that we have just one time like coordinate, though it would be easy to expand this idea into, let's say m -dimensional time like space.

That $u = u_0$ is its horizon and $u \rightarrow \infty$ is its boundary. And the metric tensor has the form of,

$$g_{\mu\nu} = \begin{pmatrix} -\frac{u^2}{L^2} \left(1 - \left(\frac{u_0^2}{u^2}\right)^d\right) & & & & 0 \\ & \begin{pmatrix} \frac{u^2}{L^2} & 0 & \cdots & 0 \\ 0 & \frac{u^2}{L^2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{u^2}{L^2} \end{pmatrix}_{d-1 \times d-1} & & \\ 0 & & & & \frac{L^2}{u^2 \left(1 - \left(\frac{u_0^2}{u^2}\right)^d\right)} \end{pmatrix} \quad (2.22)$$

The determinant of this matrix is $-\left(\frac{u}{L}\right)^{d-1}$. Now we want to solve the equation of motion for scalar field for this metric.

$$0 = -\frac{L^2}{u^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right)} \frac{\partial^2 \phi}{\partial t^2} + \frac{L^2}{u^2} \frac{\partial^2 \phi}{\partial x_i^2} + \frac{u^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right)}{L^2} \frac{\partial^2 \phi}{\partial u^2} \\ + \left(\frac{L}{u}\right)^{d-1} \left[\frac{u^{d-4}(d-3)}{L^{d-3} \left(1 - \left(\frac{u_0}{u}\right)^d\right)} - d \frac{1}{u} \left(\frac{u_0}{u}\right)^d \frac{u^{d-3}}{L^{d-3} \left(1 - \left(\frac{u_0}{u}\right)^d\right)^2} \right] \frac{\partial \phi}{\partial u} \quad (2.23)$$

We use the same method to solve equation . And we can assume that $\frac{\partial^2 \phi}{\partial t^2}$ and $\frac{\partial^2 \phi}{\partial x_i^2}$ are all constant. It is the same as we say we want to solve the d -dimensional wave equation for AdS black hole. So we have,

$$0 = \left[\frac{k_t^2 L^2}{u^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right)} - k_i^2 \frac{L^2}{u^2} \right] f(u) + \frac{u^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right)}{L^2} \frac{d^2 f(u)}{du^2} \\ + \left(\frac{L}{u}\right)^{d-1} \left[\frac{L^2 (d-3)}{u^3 \left(1 - \left(\frac{u_0}{u}\right)^d\right)} - d \frac{1}{u} \left(\frac{u_0}{u}\right)^d \frac{L^2}{u^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right)^2} \right] \frac{df(u)}{du} \quad (2.24)$$

and then,

$$0 = \left[k_t^2 - k_i^2 \left(1 - \left(\frac{u_0}{u}\right)^d\right) \right] f(u) + \frac{u^4 \left(1 - \left(\frac{u_0}{u}\right)^d\right)^2}{L^4} \frac{d^2 f(u)}{du^2} \\ + L^{d-1} \left[\frac{(d-3)}{u^d} - \frac{du_0^d}{u^{2d}} \frac{1}{\left(1 - \left(\frac{u_0}{u}\right)^d\right)} \right] \frac{df(u)}{du} \quad (2.25)$$

Now let's change the variable u into $\frac{u_0}{w}$. This form would be more useful in this context, because our horizon and boundary changes from $w = 1$ to $w = 0$.

And one can expand the answer near $w = 1$ and/or $w = 0$. The metric has the form of,

$$ds^2 = L^2 \left[\frac{u_0^2}{L^4 w^2} \left[-(1 - w^d) dt^2 + \sum_{i=1}^{d-1} dx_i^2 \right] + \frac{d^2 w}{w^2 (1 - w^d)} \right] \quad (2.26)$$

After rescaling time, and x_i coordinates we get,

$$ds^2 = \frac{L^2}{w^2} \left[-(1 - w^d) dt^2 + \sum_{i=1}^{d-1} dx_i^2 + \frac{d^2 w}{w^2 (1 - w^d)} \right] \quad (2.27)$$

and the metric tensor would be,

$$g_{\mu\nu} = \begin{pmatrix} -\frac{L^2}{w^2} (1 - w^d) & & & & 0 \\ & \begin{pmatrix} \frac{L^2}{w^2} & 0 & \cdots & 0 \\ 0 & \frac{L^2}{w^2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{L^2}{w^2} \end{pmatrix}_{d-1 \times d-1} & & \\ 0 & & & & \frac{L^2}{w^2 (1 - w^d)} \end{pmatrix} \quad (2.28)$$

The determinant of this matrix is $-(\frac{L}{w})^{2d+2}$. Now we want to solve the equation of motion of scalar field for this metric.

$$\begin{aligned} 0 = & -\frac{w^2}{L^2 (1 - w^d)} \frac{\partial^2 \phi}{\partial t^2} + \frac{w^2}{L^2} \frac{\partial^2 \phi}{\partial x_i^2} + \frac{w^2 (1 - w^d)}{L^2} \frac{\partial^2 \phi}{\partial w^2} \\ & + \frac{w}{L^2} [1 - d - w^d] \frac{\partial \phi}{\partial w} - m^2 \phi \end{aligned} \quad (2.29)$$

We use the separation of variables method to solve equation . And we can assume that $\frac{\partial^2 \phi}{\partial t^2}$ and $\frac{\partial^2 \phi}{\partial x_i^2}$ are all constant. Let's say $\phi(t, x_i, w) = A e^{ik \cdot x} f(w) + B e^{-ik \cdot x} g(w)$. For finding $f(w)$ and $g(w)$, first we assume that $B = 0$ and then we can find $f(w)$ then we assume that $A = 0$ and we can find $g(w)$. By substituting $\phi(t, x_i, w)$ in the equation and assuming $B = 0$ we get,

$$\begin{aligned} 0 = & \frac{w^2}{L^2 (1 - w^d)} k_i^2 f(w) - \sum_{i=1}^{d-1} \frac{w^2}{L^2} k_{x_i}^2 f(w) + \frac{w^2 (1 - w^d)}{L^2} \frac{\partial^2 f(w)}{\partial w^2} \\ & + \frac{w}{L^2} [1 - d - w^d] \frac{\partial f(w)}{\partial w} - m^2 f(w) \end{aligned} \quad (2.30)$$

Now let's focus on the behaviour of our function near boundary and near horizon. When $w \rightarrow 0$ it approaches to boundary. And when $w \rightarrow 1$ it approaches to horizon. Near boundary our ordinary differential equation has the form of,

$$0 = w^2 \frac{\partial^2 f(w)}{\partial w^2} + w(1-d) \frac{\partial f(w)}{\partial w} + \left[(k_t^2 - \sum_{i=1}^{d-1} k_{x_i}^2) w^2 - L^2 m^2 \right] f(w) \quad (2.31)$$

And this is a Bessel function which we have solved it before in section 2.1 we solved this equation and showed that it has two independent answer, $w^{\frac{d}{2}} J_\nu(kw)$ and $w^{\frac{d}{2}} Y_\nu(kw)$ that $\nu = \sqrt{m^2 + \frac{d^2}{4}}$, and also we proved that for $w \rightarrow 0$ the function behave,

$$\lim_{w \rightarrow 0} w^{\frac{d}{2}} J_\nu(kw) \rightarrow w^{\frac{d}{2} + \sqrt{m^2 + \frac{d^2}{4}}} \quad (2.32)$$

$$\lim_{w \rightarrow 0} w^{\frac{d}{2}} Y_\nu(kw) \rightarrow w^{\frac{d}{2} - \sqrt{m^2 + \frac{d^2}{4}}} \quad (2.33)$$

Near horizon it is a little more tricky. First we need to expand the coefficient of $f(w)$ and $\frac{\partial f(w)}{\partial w}$ near $w = 1$ and then one should try to find the behaviour of differential equation and function of $f(w)$. Let's divide both side of equation 2.30 by coefficient of $\frac{\partial^2 f(w)}{\partial w^2}$,

$$0 = \frac{\partial^2 f(w)}{\partial w^2} + \left[\frac{1-d-w^d}{w(1-w^d)} \right] \frac{\partial f(w)}{\partial w} + \left[\frac{1}{(1-w^d)^2} k_t^2 - \frac{1}{1-w^d} \sum_{i=1}^{d-1} k_{x_i}^2 - \frac{L^2 m^2}{w^2(1-w^d)} \right] f(w) \quad (2.34)$$

Now one should try to expand the coefficient of $f(w)$ and $\frac{\partial f(w)}{\partial w}$ near $w = 1$. And ignore the terms which approach to zero,

$$0 = \frac{\partial^2 f(w)}{\partial w^2} + \left[-\frac{d}{1-w^d} + \frac{1-d}{2d}(1-w^d) + \dots \right] \frac{\partial f(w)}{\partial w} + \left[\frac{1}{(1-w^d)^2} k_t^2 - \frac{1}{1-w^d} \sum_{i=1}^{d-1} k_{x_i}^2 - L^2 m^2 \left(-\frac{1}{1-w^d} - \frac{2}{d} + \dots \right) \right] f(w) \quad (2.35)$$

As one takes the limit of this function when $w \rightarrow 1$, $\frac{1}{(1-w^d)^2}$ approaches to ∞ faster than $\frac{1}{1-w^d}$. So close to $w = 1$ the differential equation has the form of,

$$0 = \frac{\partial^2 f(w)}{\partial w^2} - \frac{d}{1-w^d} \frac{\partial f(w)}{\partial w} + \frac{k_t^2}{(1-w^d)^2} f(w) \quad (2.36)$$

or,

$$0 = \frac{\partial^2 f(w)}{\partial w^2} - \frac{1}{1-w} \frac{\partial f(w)}{\partial w} + \frac{k_t^2}{(1-w)^2 d^2} f(w) \quad (2.37)$$

So the answer has the form of,

$$f(w) = A(1-w)^{ik_t} + B(1-w)^{-ik_t} = Ae^{ik_t \ln(1-w)} + Be^{-ik_t \ln(1-w)} \quad (2.38)$$

$e^{-ik_t \ln(1-w)}$ is incoming wave which go into the black hole and $e^{ik_t \ln(1-w)}$ is outgoing wave which coming out of black hole. If we choose $A = 0$ then there is no out coming term so all information, wave, is just going into the black hole and nothing come back.

Chapter 3

Scalar Field in Black Hole Anti de Sitter

3.1 Introduction

In chapter 2 we learned how to deal with AdS space and in last section, section ??, we solved the scalar field equation for simple black hole in Anti de Sitter space and studied the behavior of waves near horizon and boundary of black hole. In this chapter we want to study black holes in AdS space more generally. For example in the following section we are going to derive the equations for charged black hole and then we are going to study the behavior of wave function, general solution of scalar field near horizon and boundary. Then we apply this idea to some real systems and try to study the behavior of some condens matter systems by applying this idea to our system.

3.2 Charged Black Hole in AdS

In section ?? we saw that the metric for simple black hole has the form of,

$$ds^2 = - \left(k^2 r^2 + 1 - \frac{C}{r^{d-2}} \right) dt^2 + \frac{1}{k^2 r^2 + 1 - \frac{C}{r^{d-2}}} dr^2 + r^2 d\Omega^2 \quad (3.1)$$

or,

$$ds^2 = \frac{u^2}{L^2} \left[- \left(1 - \left(\frac{u_0}{u} \right)^d \right) dt^2 + \sum_{i=1}^{d-1} dx_i^2 \right] + \frac{L^2 du^2}{u^2 \left(1 - \left(\frac{u_0}{u} \right)^d \right)} \quad (3.2)$$

Now we can add charge to our black hole and make it charged black hole. So the metric changes as,

$$ds^2 = \frac{r^2}{R^2}(-f dt^2 + d\vec{x}^2) + \frac{R^2}{r^2} \frac{dr^2}{f}, \quad f = 1 + \frac{Q^2}{r^{2d-2}} - \frac{M}{r^d} \quad (3.3)$$

and we can find the radius of horizon,

$$f(r_0) = 0 \rightarrow M = r_0^d + \frac{Q^2}{r_0^{d-2}} \quad (3.4)$$

Also charge, energy, and entropy density have forms of,

$$\rho = \frac{2(d-2)}{c_d} \frac{Q}{\kappa^2 R^{d-1} g_F}, \quad \varepsilon = \frac{d-1}{2\kappa^2} \frac{M}{R^{d+1}}, \quad s = \frac{2\pi}{\kappa^2} \left(\frac{r_0}{R}\right)^{d-1} \quad (3.5)$$

And temprature of this system is,

$$T = \frac{dr_0}{4\pi r^2} \left(1 - \frac{(d-2)Q^2}{dr_0^{2d-2}}\right) \quad (3.6)$$

For zero temprature, $r_0^{2d-2} = \frac{(d-2)Q^2}{d}$, the redshift factor f gets the form of,

$$f = 1 - \left(\frac{r_0}{r}\right)^d + \frac{d}{d-2} \left(\frac{r_0}{r}\right)^d \left(\left(\frac{r_0}{r}\right)^{d-2} - 1 \right) \quad (3.7)$$

Near the horizon, $f \rightarrow 0$ or $r \rightarrow r_0$, we can write the taylor series, so we have,

$$f = d(d-1) \left(\frac{r-r_0}{r_0}\right)^2 + \dots \quad (3.8)$$

By changing the varibales, $r - r_0 = \frac{\lambda R_2^2}{\xi}$ and $t = \lambda^{-1} \tau$, which $R_2^2 = \frac{R^2}{d(d-1)}$ we find that the metric becomes,

$$ds^2 = \frac{R_2^2}{\xi^2} (-d\tau^2 + d\xi^2) + \frac{r_0^2}{R^2} d\vec{x}^2 \quad (3.9)$$

and for low temprature, by replacing $r_0 - r_* = \lambda \frac{R_2^2}{\xi_0}$, the metric has the form of,

$$ds^2 = \frac{R_2^2}{\xi^2} \left(-\left(1 - \frac{\xi}{\xi_0}\right) d\tau^2 + \frac{d\xi^2}{1 - \frac{\xi}{\xi_0}} \right) + \frac{r_0^2}{R^2} d\vec{x}^2 \quad (3.10)$$

and the temprature is,

$$\begin{aligned}
 T &= \frac{dr_0}{4\pi R^2} \left(1 - \frac{r_*^{2d-2}}{r_0^{2d-2}} \right) \\
 &= \frac{dr_0}{4\pi R^2} \frac{(r_0 - r_*)(r_0^{2d-1} + r_* r_0^{2d-2} + \dots + r_*^{2d-1} r_0 + r_*^{2d-1})}{r_0^{2d-2}} \\
 &= \frac{dr_0}{4\pi R^2} \frac{2(d-1)(r_0 - r_*) r_0^{2d-1}}{r_0^{2d-2}} = \frac{1}{2\pi \zeta_0}
 \end{aligned} \tag{3.11}$$

For zero temprature and for charged balck hole in AdS space near the horizon the metric tensor has the form of,

$$g_{\mu\nu} = \begin{pmatrix} -\frac{R_2^2}{\zeta^2} & & & & 0 \\ & \begin{pmatrix} \frac{r_0^2}{R^2} & 0 & \dots & 0 \\ 0 & \frac{r_0^2}{R^2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \frac{r_0^2}{R^2} \end{pmatrix} & & \\ & & & d-1 \times d-1 & \\ 0 & & & & \frac{R_2^2}{\zeta^2} \end{pmatrix} \tag{3.12}$$

and for low temprature has the form of,

$$g_{\mu\nu} = \begin{pmatrix} -\frac{R_2^2}{\zeta^2} \left(1 - \frac{\zeta^2}{\zeta_0^2} \right) & & & & 0 \\ & \begin{pmatrix} \frac{r_0^2}{R^2} & 0 & \dots & 0 \\ 0 & \frac{r_0^2}{R^2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \frac{r_0^2}{R^2} \end{pmatrix} & & \\ & & & d-1 \times d-1 & \\ 0 & & & & \frac{R_2^2}{\zeta^2} \frac{1}{1 - \frac{\zeta^2}{\zeta_0^2}} \end{pmatrix} \tag{3.13}$$

and equation of plain wave for zero and low temprature respectively would be,

$$\frac{\zeta^2}{R_2^2} \frac{\partial^2 f(\zeta)}{\partial \zeta^2} + \left[\frac{\zeta^2 k_t^2}{R_2^2} - m^2 - \frac{R^2 k_x^2}{r_0^2} \right] f(\zeta) = 0 \tag{3.14}$$

and,

$$\frac{\zeta^2(1 - \frac{\zeta^2}{\zeta_0^2})}{R_2^2} \frac{\partial^2 f(\zeta)}{\partial \zeta^2} - 2 \frac{\zeta^3}{R_2^2 \zeta_0^2} \frac{\partial f(\zeta)}{\partial \zeta} + \left[\frac{\zeta^2 k_t^2}{R_2^2(1 - \frac{\zeta^2}{\zeta_0^2})} - m^2 - \frac{R^2 k_x^2}{r_0^2} \right] f(\zeta) = 0 \quad (3.15)$$

Now, let's focus more on our original metric without doing any changing variables or taking any limits. By defining $f = 1 + \frac{Q^2}{r^{2d-2}} - \frac{M}{r^d}$, the metric tensor of equation 3.3 is,

$$g_{\mu\nu} = \begin{pmatrix} -\frac{r^2}{R^2} f & & & & 0 \\ & \begin{pmatrix} \frac{r^2}{R^2} & 0 & \cdots & 0 \\ 0 & \frac{r^2}{R^2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{r^2}{R^2} \end{pmatrix}_{d-1 \times d-1} & & \\ 0 & & & \frac{R^2}{r^2 f} & \end{pmatrix} \quad (3.16)$$

Now we are able to write the equation of motion and solve it for scalar field, which describes the behaviour of plain wave in Anti de Sitter space with existence of a charged black hole. Then we can study the behaviour of plain wave near the horizon and boundary and then we are able to find the Green function. So let's take a look at its differential equation,

$$0 = \frac{1}{R^2} \left[(d+1)r - \frac{(d-3)Q^2}{r^{2d-3}} - \frac{M}{r^{d-1}} \right] \frac{\partial \phi(r)}{\partial r} + \frac{r^2 f}{R^2} \frac{\partial^2 \phi(r)}{\partial r^2} + \left[\frac{R^2}{r^2 f} k_t^2 - \frac{R^2}{r^2} k_x^2 - m^2 \right] \phi(r) \quad (3.17)$$

Now we are ready to study the behaviour of our wave function near horizon, $r \rightarrow r_0$, and near boundary, $r \rightarrow \infty$. For $d > 1$, near boundary redshift factor f approaches to 1 and the equation 3.17 would be simplified as,

$$0 = r^2 \frac{\partial^2 \phi(r)}{\partial r^2} + (d+1)r \frac{\partial \phi(r)}{\partial r} - R^2 m^2 \phi(r) \quad (3.18)$$

By substituting $\phi(r) = r^\alpha$ in our differential equation we get, $0 = \alpha^2 + \alpha d - R^2 m^2$. So the answer near boundary is looks like,

$$\phi(r) = A(k_\mu) r^{\Delta-d} + B(k_\mu) r^{-\Delta} \quad , \quad \Delta = \frac{d}{2} + \sqrt{\frac{d^2}{4} + R^2 m^2} \quad (3.19)$$

Horizon dose not have a zero area so its radius is not zero, too. Equation 3.4 gives us the radius of horizon. Also instead of mass of black hole M in redshift factor f we can substitute $r_0^d + \frac{Q^2}{r_0^{d-2}}$, so the redshift factor becomes,

$$f = \left(1 - \frac{r_0^d}{r^d}\right) + \frac{Q^2}{r^d} \left[\frac{1}{r^{d-2}} - \frac{1}{r_0^{d-2}} \right] \quad (3.20)$$

$$= (r - r_0) \left[\frac{r^{d-1} + r^{d-2}r_0 + \dots + r_0^{d-2}r + r_0^{d-1}}{r^d} - \frac{Q^2}{r^d} \frac{r^{d-3} + r^{d-4}r_0 + \dots + r_0^{d-4}r + r_0^{d-3}}{r^{d-2}r_0^{d-2}} \right] \quad (3.21)$$

By defining $Q^2 \equiv \frac{d}{d-2} r_*^{2d-2}$, near horizon we can simplify our differential equation when $r \rightarrow r_0$, and $f \rightarrow (r - r_0) \left[\frac{d}{r_0} - \frac{dr_*^{2d-2}}{r_0^{2d-1}} \right]$,

$$0 = \frac{\partial^2 \phi(r)}{\partial r^2} + \frac{1}{(r - r_0)} \frac{\partial \phi(r)}{\partial r} + \frac{1}{(r - r_0)^2} \times \frac{R^4 k_t^2}{d^2 r_0^2 \left(1 - \frac{r_*^{2d-2}}{r_0^{2d-2}}\right)^2} \phi(r) \quad (3.22)$$

or simply,

$$0 = \frac{\partial^2 \phi(r)}{\partial r^2} + \frac{1}{(r - r_0)} \frac{\partial \phi(r)}{\partial r} + \frac{\beta^2}{(r - r_0)^2} \phi(r) \quad (3.23)$$

when

$$\beta^2 = \frac{R^4 k_t^2}{d^2 r_0^2 \left(1 - \frac{r_*^{2d-2}}{r_0^{2d-2}}\right)^2} \quad (3.24)$$

And answer of this differential equation is:

$$\phi(r) = A(r - r_0)^{\sqrt{-\beta^2}} + B(r - r_0)^{-\sqrt{-\beta^2}} = A e^{i\beta \ln(r - r_0)} + B e^{-i\beta \ln(r - r_0)} \quad (3.25)$$

3.3 Zero Temperature

The last equations are true when we are far from zero temprature. In this part, we are dealing with zero temperature, and then we will study these equations in low temprature, close to zero temperature. Equation 3.20 is the red shift factor, f , in finit temprature, but in zero temprature its form changes and, as we will

see, we get factor of $(r - r_0)^2$. For simplicity, for our purpose we just focus on case $d = 4$, 3 conventional space direction and one conventional time direction and an other spacelike direction for having AdS_5/CFT_4 theory. Zero temprature is when $r_0 = r_*$, so by substituting $Q = \frac{d}{d-2}r_0^{2d-2}$ for red shift factor one gets,

$$\begin{aligned} f &= \frac{(d-2)r^{2d-2} - (2d-2)r_0^d r^{d-2} + dr_0^{2d-2}}{(d-2)r^{2d-2}}, \quad d = 4 \\ &= \frac{(r-r_0)^2(r+r_0)^2(r^2+2r_0^2)}{r^6} \end{aligned} \quad (3.26)$$

And equation of motion for scalar field has the form of,

$$\begin{aligned} 0 &= \frac{1}{R^2} \left[\frac{(r-r_0)(r+r_0)(5r^4+5r_0^2r^2+2r_0^4)}{r^5} \right] \frac{\partial \phi(r)}{\partial r} \\ &\quad + \frac{r^2 f}{R^2} \frac{\partial^2 \phi(r)}{\partial r^2} + \left[\frac{R^2}{r^2 f} k_t^2 - \frac{R^2}{r^2} k_x^2 - m^2 \right] \phi(r) \end{aligned} \quad (3.27)$$

Note that in this case, zero temprature, r_0 is exactly equal to r_* . Now we want to find the Green function and behaviour of our function near boundart, $r \rightarrow \infty$ and close to horizon, $r \rightarrow r_0$. Near boundary nothing would change from before, everything we had in last section is true for zero temprature too and we have the same form of answer for zero tamprature. Because Q , charge of black hole wich is related to temprature, and M , mass of black hole which is related to radius of horizon, do not play any role in the equation of motion, equation 3.17, near boundary, in case of $r \rightarrow \infty$. Close to horizon, $r \rightarrow r_0$, redshift factor has the form of $f = \frac{12}{r_0^2}(r-r_0)^2$, equation of motion 3.27 gets the form of,

$$0 = (r-r_0)^4 \frac{\partial^2 \phi(r)}{\partial r^2} + 2(r-r_0)^3 \frac{\partial \phi(r)}{\partial r} + \frac{R^4 k_t^2}{144} \phi(r) \quad (3.28)$$

and the answer has the form of,

$$\phi(r, \mu) = C(k_\mu) e^{\frac{i\beta}{r-r_0}} + D(k_\mu) e^{\frac{-i\beta}{r-r_0}}, \quad \beta = \frac{R^2 k_t}{12} \quad (3.29)$$

$e^{\frac{i\beta}{r-r_0}}$ is incoming wave which is going through black hole and $e^{\frac{-i\beta}{r-r_0}}$ is out going wave which is going out of black hole. In our case, we are mostly interested in case that we have just incoming wave and there is no out going wave which go out of black hole. So one need to choos $B = 0$. Then we are able to find the Green's function,

$$G(k_\mu) \equiv \frac{B(k_\mu)}{A(k_\mu)}, \quad \phi(r, k_\mu) = A(k_\mu) r^{\Delta-d} + B(k_\mu) r^{-\Delta} \quad (3.30)$$

3.4 Charged Scalar Field in Charged Black Hole

In this section we try to study behaviour of charged scalar field with charged black hole geometry in AdS space. Then we will learn more about the low frequencies, and then one should suggest low frequency limit to avoid breaking of the theory. Low frequency is breaking because of infinite number of degrees of freedom. But let's first look at the action of charged scalar field

$$S = - \int d^{d+1}x \sqrt{-|g|} [(D_\mu \phi)^* (D^\mu \phi) + m^2 \phi^* \phi] \quad (3.31)$$

where,

$$D_\mu = \partial_\mu - iqA_\mu \quad (3.32)$$

Note that action 3.31 just depends on q through:

$$\mu_q \equiv \mu q \quad (3.33)$$

which is the effective chemical potential for the field of charge q . We are interested to work on frequency space so we should take Fourier transformation of $\phi(r, x^\mu)$ and writing,

$$\phi(r, x^\mu) = \int \frac{d^d k}{\sqrt{(2\pi)^d}} \phi(r, k_\mu) e^{ik_\mu x^\mu} \quad (3.34)$$

And from now we work only on frequency space. After taking Fourier transformation we have,

$$\frac{-1}{\sqrt{-|g|}} \partial_r (\sqrt{-|g|} g^{rr} \partial_r \phi) + [g^{ii} (k^2 - u^2) + m^2] \phi = 0, \quad (3.35)$$

where,

$$u(r) = \sqrt{\frac{-g_{ii}}{g_{rr}}} \left(\omega + \mu_q \left(1 - \frac{r_0^{d-2}}{r^{d-2}} \right) \right), \quad (3.36)$$

If the action is defined by equation 3.31. By considering all above equations then for motion equation we get,

$$\begin{aligned} 0 = & \frac{r^2 f}{R^2} \frac{\partial^2 \phi(r)}{\partial r^2} + \frac{1}{R^2} \left[(d+1)r - \frac{(d-3)Q^2}{r^{2d-3}} - \frac{M}{r^{d-1}} \right] \frac{\partial \phi(r)}{\partial r} \\ & + \left[\frac{R^2}{r^2 f} \left(k_t + \mu_q \left(1 - \frac{r_0^{d-2}}{r^{d-2}} \right) \right)^2 - \frac{R^2}{r^2} k_x^2 - m^2 \right] \phi(r) \end{aligned} \quad (3.37)$$

Now we are ready to study $\phi(r, k_\mu)$ in finit temprature near boundary and close to horizon. As we have seen the radius of horizon is r_0 and boundary is located at $r \rightarrow \infty$. What ever we had on the last sections is true here. And nothing would be changed in our equation near boundary and close to horizon. By analyzing equation of 3.37 we will get the following answers near boundary and close to horizon respectively,

$$\phi(r, k_\mu) = A(k_\mu)r^{\Delta-d} + B(k_\mu)r^{-\Delta} \quad , \quad for \quad r \rightarrow \infty \quad (3.38)$$

where $\Delta = \frac{d}{2} + \sqrt{\frac{d^2}{4} + m^2 R^2}$ and,

$$\phi(r) = C(k_\mu)e^{i\beta \ln(r-r_0)} + D(k_\mu)e^{-i\beta \ln(r-r_0)} \quad (3.39)$$

where $\beta = \frac{R^2 k_t}{d(1 - \frac{r_*^2}{r_0^2})}$. In the above equation, as we have seen before, $e^{-i\beta \ln(r-r_0)}$ and $e^{i\beta \ln(r-r_0)}$ are incoming wave, which goes into the black hole, and out going wave, which is going out of black hole. And finally Green's function is defined,

$$G(k_\mu) \equiv \frac{B(k_\mu)}{A(k_\mu)} \quad (3.40)$$

Let's be more accurate close to the boundary and horizon. We want to expand the equations near boundary and horizon and find more terms to make our theory more accurate. Let's do it first for boundary. But becasue we are intrested in cases which $d = 4$, 4-dimentional cases, so here we do not solve the problem for general d -dimentional case. We can rewrite redshift factor,

$$f(r) = \frac{r-r_0}{r^6 r_0^2} \left[(4r_0^7 - 4r_*^6 r_0) + (14r_0^6 - 2r_*^6)(r-r_0) + 20r_0^5(r-r_0)^2 + 15r_0^4(r-r_0)^3 + 6r_0^3(r-r_0)^4 + r_0^2(r-r_0)^5 \right] \quad (3.41)$$

and the coefficient of $\frac{\partial \phi}{\partial r}$, let's call it $g(r)$, is

$$g(r) = \frac{1}{r^5 r_0^2} \left[(4r_0^8 - 4r_0^2 r_*^6) + (28r_0^7 - 4r_0 r_*^6)(r-r_0) + (74r_0^6 - 2r_*^6)(r-r_0)^2 + 100r_0^5(r-r_0)^3 + 75r_0^4(r-r_0)^4 + 30r_0^3(r-r_0)^5 + 5r_0^2(r-r_0)^6 \right] \quad (3.42)$$

In all of our equation ϕ is a function of r , $k_{\vec{x}}$, and $k_t = \omega$ and for simplicity we just write $\phi(r)$. But we are aware that actualy it is function of all of these

terms through fourier transformation. So now we are able to rewrite the equation of motion in $(r - r_0)^n$ terms, when $r \rightarrow r_0$,

$$\begin{aligned}
0 = & \frac{(r - r_0)^2}{R^2 r_0^{14}} \left[16(r_0^7 - r_*^6 r_0)^2 + 16(r_0^7 - r_*^6 r_0)(14r_0^6 - 2r_*^6)(r - r_0) \right. \\
& \left. + (14r_0^6 - 2r_*^6)^2 (r - r_0)^2 + \dots \right] \frac{\partial^2 \phi(r)}{\partial r^2} \\
& + \frac{(r - r_0)}{R^2 r_0^{14}} \left[16(r_0^7 - r_*^6 r_0)^2 + 12(r_0^7 - r_*^6 r_0)(14r_0^6 - 2r_*^6)(r - r_0) \right. \\
& \left. + 2(14r_0^6 - 2r_*^6)^2 (r - r_0)^2 + \dots \right] \frac{\partial \phi(r)}{\partial r} \\
& + \left[\frac{R^2}{r_0^2} \left(\omega^2 + 2\omega\mu_q \frac{2}{r_0} (r - r_0) + \mu_q^2 \frac{4}{r_0^2} (r - r_0)^2 + \dots \right) \right. \\
& \left. - \frac{r - r_0}{r_0^8} \left[(4r_0^7 - 4r_*^6 r_0) + (14r_0^6 - 2r_*^6)(r - r_0) + \dots \right] \left(\frac{R^2}{r_0^2} k_x^2 + m^2 \right) \right] \phi(r)
\end{aligned} \tag{3.43}$$

The solution has the form of,

$$\begin{aligned}
\phi(r) = & A(k_\mu, \omega) \left[e^{-i\beta \ln(r - r_0)} + a_1 e^{-i\beta \ln(r - r_0)} (r - r_0) + a_2 e^{-i\beta \ln(r - r_0)} (r - r_0)^2 \right] \\
& + B(k_\mu, \omega) \left[e^{i\beta \ln(r - r_0)} + a_1 e^{i\beta \ln(r - r_0)} (r - r_0) + a_2 e^{i\beta \ln(r - r_0)} (r - r_0)^2 \right]
\end{aligned} \tag{3.44}$$

if $\tilde{M} = \frac{R^4}{r_0^2} k_x^2 + R^2 m^2$, $\tilde{r} = \frac{r_*}{r_0}$, and $n = \mp i\beta$ then,

$$\begin{aligned}
a_1 = & - \frac{8(7 - \tilde{r}^6)(n^2 - n) + 6(7 - \tilde{r}^6)n + \frac{\omega\mu_q R^4}{r_0^2(1 - \tilde{r}^6)} - \tilde{M}}{16(n^2 + n) + 16(n + 1) + \frac{R^4 \omega^2}{r_0^2(1 - \tilde{r}^6)}} \times \frac{4}{r_0} \\
a_2 = & - \left[4(7 - \tilde{r}^6)^2(n^2 - n) + 8(7 - \tilde{r}^6)^2 n + 4 \frac{R^4 \mu_q^2}{r_0^2} - 2(7 - \tilde{r}^6)\tilde{M} \right. \\
& \left. + 4r_0 a_1 \left((1 - \tilde{r}^6)(7 - \tilde{r}^6)(8n^2 + 14n + 6) + \omega\mu_q \frac{R^4}{r_0^2} - (1 - \tilde{r}^6)\tilde{M} \right) \right] \\
& \times \left[16r_0^2(1 - \tilde{r}^6)^2(n^2 + 4n + 4) + \omega^2 R^4 \right]^{-1}
\end{aligned} \tag{3.45}$$

We can continue what we have done above to find the higher order terms of $\phi(r)$ near horizon. For our purpose these terms are more than enough. For the

next step we are interested to calculate more terms near boundary. So let's look at equation of motion one more time when we are in n -dimensional space,

$$\begin{aligned}
 0 = & \left[r^4 - \frac{2M}{r^{d-4}} + \frac{(d-3)2Q^2}{r^{2d-6}} + \frac{M^2}{r^{2d-4}} - \frac{Q^2M}{r^{3d-6}} + \frac{(d-3)Q^4}{r^{4d-8}} \right] \frac{\partial^2 \phi(r)}{\partial r^2} \\
 & \left[(d+1)r^3 - \frac{(d+2)M}{r^{d-3}} + \frac{4Q^2}{r^{2d-5}} + \frac{(d-4)MQ^2}{r^{3d-3}} + \frac{M^2}{r^{2d-3}} - \frac{(d-3)Q^4}{r^{3d-5}} \right] \frac{\partial \phi(r)}{\partial r} \\
 & \left[\left(R^4 \left(k_t + \mu_q \left(1 - \frac{r_0^{d-2}}{r^{d-2}} \right) \right)^2 - R^4 k_{\vec{x}}^2 \right) - R^2 m^2 \left(r^2 + \frac{Q^2}{r^{2d}} - \frac{M}{r^{d-2}} \right) \right] \phi(r)
 \end{aligned} \tag{3.46}$$

For case $d = 4$ answer has the form of,

$$\begin{aligned}
 \phi(r) = & A(k_\mu, \omega) r^{\Delta-4} \left[1 + a_1 \frac{1}{r^2} + a_2 \frac{1}{r^4} + \dots \right] \\
 & B(k_\mu, \omega) r^{-\Delta} \left[1 + a'_1 \frac{1}{r^2} + a'_2 \frac{1}{r^4} + \dots \right]
 \end{aligned} \tag{3.47}$$

where, $n = \Delta - d$ and $-\Delta$ respectively for a_1 , a_2 and a'_1 , a'_2 .

$$\begin{aligned}
 a_1, a'_1 = & - \frac{R^4(k_t + \mu_q)^2 - R^4 k_{\vec{x}}^2}{n^2 - 4 - R^2 m^2} \\
 a_2, a'_2 = & \frac{2M(n^2 + 2n) - a_2(R^4(k_t + \mu_q)^2 - R^4 k_{\vec{x}}^2) + MR^2 m^2}{n^2 - 4n - R^2 m^2}
 \end{aligned} \tag{3.48}$$

3.4.1 Exeptional Cases

In this part we talk about some exeptional cases, for example when $\Delta - d = -\Delta$ then our both independent answer become dependent and they are not different solutions of our differential equation. And we know that for second order differential equation we have to get two independent solution.

So for case $\Delta - d = -\Delta$, or in other word $\Delta = \frac{d}{2}$ or $-\frac{d^2}{4} = R^2 m^2$, our solution bear boundary has the form of,

$$\phi(r) = A(k_\mu, \omega) r^{-\Delta} + B(k_\mu, \omega) r^{-\Delta} \ln(r) \tag{3.49}$$

and for more general cases, when we are looking for more terms near boundary and $\Delta = k$ and k in an integer number and $d = 4$, answer has the form of,

$$\begin{aligned}
\phi(r) = & A_1(k_\mu, \omega)r^n + r^{n-2} [A_{21}(k_\mu, \omega) + A_{22}(k_\mu, \omega) \ln(r)] \\
& + r^{n-4} [A_{31}(k_\mu, \omega) + A_{32}(k_\mu, \omega) \ln(r) + A_{33}(k_\mu, \omega) \ln^2(r)] \\
& + \dots
\end{aligned} \tag{3.50}$$

wehre $n = \max(-\Delta, \Delta - 4)$.

In equation 3.50 there are just two independent coefficient, lets say A_1 and A_{21} , and the other are dependent to these two coefficient. For the first dependent coefficient we get,

$$A_{22}(k_\mu, \omega) = \frac{R^4(k_x^2 - k_t^2)}{2n}, \quad n = \max(-\Delta, \Delta - 4) \tag{3.51}$$

3.5 Numerical Approach

Equation of motion of form 3.37 is hard, or in other word is impossible, to solve analytically so far, consequently for studying the features of Green function one needs to solve the motion equation by numerical methods. In this section, we try to introduce an appropriate numerical method for solving this equation and then we use that method for studying the Green function under different conditions. Our aim is developing a numerical method which use near horizon condition to calculate the Green function.

One of the best numerical method with hight accuracy is Runge-Kutta method. For the first time, this technique were developed and used by the German mathematicians C. Runge and M.W. Kutta. Let's define our initial value problem as following,

$$\phi(k_\mu, \omega, r_i) = \phi_i, \quad \frac{\partial \phi(k_\mu, \omega, r_i)}{\partial r} = \dot{\phi}_i \tag{3.52}$$

For initial condition we want to start from horizon, or it is better to say near horizon, and calculate the Green funtion. In section 3.4 we learned that the solution near horizon has the form of,

$$\begin{aligned}
\phi(k_\mu, \omega, r) = & A(k_\mu, \omega)e^{-i\beta \ln(r-r_0)} [1 + a_1(r-r_0) + a_2(r-r_0)^2 + \dots] \\
& + B(k_\mu, \omega)e^{i\beta \ln(r-r_0)} [1 + a_1(r-r_0) + a_2(r-r_0)^2 + \dots]
\end{aligned} \tag{3.53}$$

Exactly in $r = r_0$ the fuction is not difined so one should start somewhere close to r_0 . We will start from point $r = r_0 + h$, lets call it start point, as our initial condition. So for initial condition we have,

$$\begin{aligned}
\phi_i(k_\mu, \omega, r_0 + h) &= A(k_\mu, \omega) e^{-i\beta \ln(h)} [1 + a_1(h) + a_2(h)^2 + \dots] \\
&\quad + B(k_\mu, \omega) e^{i\beta \ln(h)} [1 + a_1(h) + a_2(h)^2 + \dots] \\
\dot{\phi}_i(k_\mu, \omega, r_0 + h) &= -i\beta A(k_\mu, \omega) \frac{e^{-i\beta \ln(h)}}{h} [1 + a_1(h) + a_2(h)^2 + \dots] \\
&\quad + i\beta B(k_\mu, \omega) \frac{e^{i\beta \ln(h)}}{h} [1 + a_1(h) + a_2(h)^2 + \dots] \\
&\quad + A(k_\mu, \omega) e^{-i\beta \ln(h)} [a_1 + 2a_2(h) + \dots] \\
&\quad + B(k_\mu, \omega) e^{i\beta \ln(h)} [a_1 + 2a_2(h) + \dots]
\end{aligned} \tag{3.54}$$

We will use Runge-Kutta scheme to find the value of $\phi(k_\mu, \omega, r)$ in each point. We will start from start point and continue going forward in r direction untile we get to the aim point, let's say r_f . In each step we use forth order Runge-Kutta scheme to find the next point. If we have the derivative of n^{th} point then we are able to use the following formula to calculate the $(n+1)^{th}$ point,

$$\begin{aligned}
\phi_{n+1} &= \phi_n + \frac{k_1 + 2k_2 + 2k_3 + k_4}{6} \\
\dot{\phi}_{n+1} &= \dot{\phi}_n + \frac{l_1 + 2l_2 + 2l_3 + l_4}{6}
\end{aligned} \tag{3.55}$$

where,

$$\begin{aligned}
RH_{cons}(r, \phi, \dot{\phi}) &= \frac{\partial^2 \phi}{\partial r^2} \\
&= -\frac{1}{f} \left[\frac{d+1}{r} - \frac{(d-3)Q^2}{r^{2d-1}} - \frac{M}{r^{d+1}} \right] \dot{\phi} \\
&\quad - \left[\frac{R^4}{r^4 f^2} \left(k_t + \mu_q \left(1 - \frac{r_0^{d-2}}{r^{d-2}} \right) \right)^2 - \frac{R^4}{r^4 f} k_{\bar{x}} - \frac{R^2 m^2}{r^2 f} \right] \phi \\
l_1 &= h RH_{cons}(r_n, \phi_n, \dot{\phi}_n) \\
k_1 &= h \dot{\phi}_n \\
l_2 &= h RH_{cons}(r_n + 0.5h, \phi_n + 0.5k_1, \dot{\phi}_n + 0.5l_1) \\
k_2 &= h (\dot{\phi}_n + 0.5l_1) \\
l_3 &= h RH_{cons}(r_n + 0.5h, \phi_n + 0.5k_2, \dot{\phi}_n + 0.5l_2) \\
k_3 &= h (\dot{\phi}_n + 0.5l_2) \\
l_4 &= h RH_{cons}(r_n + h, \phi_n + k_3, \dot{\phi}_n + l_3) \\
k_4 &= h (\dot{\phi}_n + 0.5l_3)
\end{aligned} \tag{3.56}$$

This method is 4^{th} order Runge-Kutta method, say RK4, and its error is $\frac{1}{120}h^5\phi^5(\xi)$ which ξ is between r_n and r_{n+1} , and $\phi^5(\xi)$ is 5^{th} derivative of function ϕ .

3.5.1 Adaptive grid size

In Runge-Kutta scheme it is easy to change the grid size in each step of solution. In each step we can use a independent grid size of h , which means that this grid size do not depent on its last grid size and its following grid size. In prolem like this which in some rigions like near horizon the equations has fluctuating features for not loosing the details and get the appropriate solutions, one need to have a small grid size and close to boundary which it is a smooth function without any fluctuating behaviour small grid size slow down the computational process and make the code time expensive. So one need to find a way to deal with the grid size regard to feature of our function, make the grid size small when we have a fluctuations and/or make the grid size when we have a smooth function. We can change our varibale to new ones to make our function smooth in all regions to use the same grid size. Always this method is not working because it is hard to find an appropriate coordinates to change our function to a simple smooth function. The best way is that introduce a way to deal with grid

size directly. In each step it would be good to find an appropriate grid size independently and apply that grid size for that specific step. And we do the same process for each step, then we can control the error of our answer by changing the grid size into appropriate one.

First of all, we are looking at the error of each step. As we may know the error of 4th order Runge-Kutta is,

$$err = \frac{1}{120} h^5 \phi^5(\xi) \quad , \quad \phi^5(\xi) = \frac{\partial^5 \phi(\xi)}{\partial \xi^5} \quad , \quad r_i \leq \xi \leq r_i + h \quad (3.57)$$

so we can consider specific error for each step and then we are able to calculate step size. The problem is that we don't know exactly the value of $\phi^5(\xi)$. In maximum $\phi^5(\xi)$ is the maximum of 5th derivative of $\phi(\xi)$ between r_i and r_{i+1} , but even we don't know the maximum value of $\phi^5(\xi)$. So we need to define another constrain for h . We know that for this specific differential equations the solution near horizon is fluctuating so rapidly but as it going forward to infinity it gets smoother, by smoother we mean that its higher derivative gets smaller and smaller. So approximatly it is possible to find the maximum of 5th derivative by finding the 5th derivative of approximate solution near horizon. By using equation 3.19 and considering incoming wave for 5th derivative we gets,

$$\phi^5(r) = \frac{\partial^5 \phi(r)}{\partial r^5} = \frac{(10\beta^2 + 5\beta^4) + (24\beta + \beta^5)i}{(r - r_0)^5} e^{-i\beta \ln(r - r_0)} \quad (3.58)$$

If we start our solution from $r_0 + h_1$ then from equation it would be obvious that maximum of 5th derivative is $\sqrt{(10\beta^2 + 5\beta^4)^2 + (24\beta + \beta^5)^2} h_1^{-5}$. We can use this fomula and equation 3.57 to calculate the step size h_* near horizon. So for h we get:

$$h_* = h_1 \left[\frac{120 \text{ err}}{\sqrt{(10\beta^2 + 5\beta^4)^2 + (24\beta + \beta^5)^2}} \right]^{1/5} \quad (3.59)$$

and for all of our domain we can calculate the grid size h ,

$$h = \min(h_*, \left[\frac{120 \text{ err}}{\sqrt{(10\beta^2 + 5\beta^4)^2 + (24\beta + \beta^5)^2}} \right]^{1/5}, 1.0) \quad (3.60)$$

Chapter 4

Dirac Equation in Black Hole AdS

4.1 Introduction

In chapter 3 we learned how to deal with Scalar Field in Black Hole AdS geometry. Now, in the following chapter we go farther and will look at Dirac Equation and its solution in Black Hole AdS geometry, whose description gave rise to fermionic spin $1/2$ systems. First we will look at Lorentz Lie Algebra which is creating the basis for our discussion then try to derive Dirac equation of motion for Black Hole AdS and then we will learn more about the behaviour of the wave function solution near boundary and close to horizon, and for the next step we will try to solve the equation of motion completely, by applying appropriate numerical method to our problem. At the end we bring our results and discuss about them.

4.2 Lorentz Lie Algebra

Nature is Lorentz invariant so the realistic theory is a theory that obeys Lorentz invariance. We are just interested in theories that hold Lorentz transformation, and consequently obey Lorentz Lie Algebra. Fields can be in form of scalars, vectors, or in more complicated cases tensors. We learned how to deal with Lorentz transformation in scalar fields and as we have seen Klein-Gordon equation is scalar equation for describing the motion of waves with zero spin that is Lorentz invariant. So for studying systems with zero spins Klein-Gordon equation can be an appropriate choice. For spin $1/2$ systems, fermionic sys-

tems, Dirac Equation is our choice because as we will see it is hold after Lorents trasformation. We get back to this issue later. Now we have to focouse more one Lorentz trasformation for the vector field, because fermionic systems are discribed with vector fields. Assume that ϕ^μ 's are component of vector Φ , then for vectore Φ Lorentz transformation has the form of,

$$\phi^\mu \rightarrow \phi'^\mu(x') = \Lambda^\mu_\nu \phi^\nu(\Lambda^{-1}x) \quad (4.1)$$

note that x is a vector itself, it means that each component of vector $\phi^\mu(x)$ is a function of vector x . In 4-dimentional case $x = (x^0, x^1, x^2, x^3)$.

For creating a Lie Algebra we need to look at infinitesimal trasformation. For infinitestiamal Lorentz trasformation we have,

$$\Lambda^\mu_\nu = \delta^\mu_\nu + \omega^\mu_\nu \quad (4.2)$$

and for general cases we can define our Algebra,

$$[\mathcal{M}^{\mu\nu}, \mathcal{M}^{\rho\sigma}] = i[g^{\nu\rho} \mathcal{M}^{\mu\sigma} - g^{\mu\rho} \mathcal{M}^{\nu\sigma} + g^{\mu\sigma} \mathcal{M}^{\nu\rho} - g^{\nu\sigma} \mathcal{M}^{\mu\rho}] \quad (4.3)$$

then Lorentz trasformation can be expressed as exponential,

$$\Lambda = \exp(-i\theta_{\mu\nu} \mathcal{M}^{\mu\nu}) \quad (4.4)$$

where $\mathcal{M}^{\mu\nu}$'s are basis elements of Lorentz trasfomation group and $\theta_{\mu\nu}$'s are some numbers which tells what kind of Lorentz trasformation we are doing, let's say for each basis what value we assigned. For example we can say that for boost basis in x^1 direction we speed up the coordinate with v , or the coordiantes in (x, y) plain are rotated clockwise with ω .

It was the representation of Lorentz trasfomation in finite-dimention. Now, we are intrested to find the representation corresponding to spin 1/2. For that perpose we use Dirac Algebra. The Dirac Algebra says,

$$\{\gamma^\mu, \gamma^\nu\} \equiv \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu} \times \mathbf{1}_{n \times n} \quad (4.5)$$

and spin operator is defined by,

$$S^{\mu\nu} = \frac{i}{4} [\gamma^\mu, \gamma^\nu] \quad (4.6)$$

Having Dirac Algebra, equation 4.5, and definition of Spin operator, equation 4.6, it is easy to show that $S^{\mu\nu}$ obey the Lorentz Lie Algebra . For 4-dimentional Minkowski space, one form of respresentation of γ^j , and γ^0 is,

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix} \quad (4.7)$$

where σ^i 's are Pauli's sigma matrices, and by using the definition of Spin operators, we get,

$$S^{0i} = \frac{i}{4} [\gamma^0, \gamma^i] = -\frac{i}{2} \begin{pmatrix} \sigma^i & 0 \\ 0 & \sigma^i \end{pmatrix}, \quad S^{ij} = \frac{i}{4} [\gamma^i, \gamma^j] = \frac{1}{2} \varepsilon^{ijk} \begin{pmatrix} \sigma^k & 0 \\ 0 & \sigma^k \end{pmatrix} \quad (4.8)$$

4.2.1 Gamma matrices in n-dimensional space

Clifford Algebra, or Dirac Algebra, in general dimention nad Minkowski space-time has the form of,

$$[\gamma^\mu, \gamma^\nu] = \gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu = 2\eta^{\mu\nu} \mathbf{1} \quad (4.9)$$

and one specific representation of Gamma martices is,

$$\begin{aligned} \gamma^0 &= \mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1} \otimes \dots \\ \gamma^1 &= \sigma_1 \otimes \mathbf{1} \otimes \mathbf{1} \otimes \dots \\ \gamma^2 &= \sigma_2 \otimes \mathbf{1} \otimes \mathbf{1} \otimes \dots \\ \gamma^3 &= \sigma_3 \otimes \sigma_1 \otimes \mathbf{1} \otimes \dots \\ \gamma^4 &= \sigma_3 \otimes \sigma_2 \otimes \mathbf{1} \otimes \dots \\ \gamma^5 &= \sigma_3 \otimes \sigma_3 \otimes \sigma_1 \otimes \dots \\ \dots &= \dots \end{aligned} \quad (4.10)$$

Where σ_i is Pauli's matrices. now we are able to find gamma matrices for arbitrary dimention. Note that all of these matrices are Hermition matrices so to chage these matrices into Anti-Hermition, one can multiply the whole matric by i . Now we are able to creat spacelike or timelike gamma matrices.

4.2.2 Gamma matrices in Curved space-time

In this section we will focuse on d -dimensional curved space-time gamma matrices. From General Relativity, we know that in each point of p in manifold, there is sets of basises which in these basises the matric gets the form of Minkowski metric, lets call them Vielbein basis. It is easy to show that the law of trasfomation of basis for metrics has the following form,

$$\eta_{\alpha\beta} v_i^\alpha v_j^\beta = g_{ij} \quad (4.11)$$

where,

$$v_\mu^\alpha = \partial_\mu \xi^\alpha \quad (4.12)$$

If γ^α be the gamma matrix in vielbein coordinate basis and $\tilde{\gamma}^\mu$ be the gamma matrix in the original coordinate basis the relation between these two is,

$$\tilde{\gamma}^\mu = v_\mu^\alpha \gamma^\alpha \quad (4.13)$$

then one can check that the gamma matrices, $\tilde{\gamma}^\mu$, holds the Clifford Algebra in curved space-time

$$\{\tilde{\gamma}^\mu, \tilde{\gamma}^\nu\} = 2g^{\mu\nu} \quad (4.14)$$

So for any diagonal metric one can find the γ^μ matrices with,

$$\gamma^\mu = \sqrt{|g^{\mu\mu}|} \tilde{\gamma}^\mu \quad (4.15)$$

Lets define $\mathbf{sgn} \ g_{ij} = [- + + +]$. We are interested in AdS_5 space and Charged Black Hole AdS_5 . The metrics, respectively have forms of,

$$ds^2 = \frac{r^2}{L^2} (-dt^2 + dx^2 + dy^2 + dz^2) + \frac{L^2}{r^2} (dr^2) \quad (4.16)$$

and

$$ds^2 = \frac{r^2}{L^2} (-f(r)dt^2 + dx^2 + dy^2 + dz^2) + \frac{L^2}{r^2 f(r)} (dr^2) \quad (4.17)$$

where,

$$f(r) = 1 + \frac{Q^2}{r^{2d-2}} - \frac{M}{r^d} \quad (4.18)$$

and gamma matrices for simple AdS_5

$$\begin{aligned} \tilde{\gamma}^0 &= i\frac{L}{r} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \tilde{\gamma}^1 = i\frac{L}{r} \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix}, \quad \tilde{\gamma}^2 = i\frac{L}{r} \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \\ \tilde{\gamma}^3 &= i\frac{L}{r} \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix}, \quad \tilde{\gamma}^5 = \frac{r}{L} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \end{aligned} \quad (4.19)$$

and for Charged Blach Hole AdS_5 is,

$$\begin{aligned}\tilde{\gamma}^0 &= i \frac{L}{\sqrt{f(r)r}} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \tilde{\gamma}^1 = i \frac{L}{r} \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix}, \quad \tilde{\gamma}^2 = i \frac{L}{r} \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \\ \tilde{\gamma}^3 &= i \frac{L}{r} \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix}, \quad \tilde{\gamma}^5 = \frac{\sqrt{f(r)r}}{L} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}\end{aligned}\quad (4.20)$$

4.3 Dirac Equation for curved space-time

In this section we try to derive the theory of fermiones for curved space-time by considering the Lorentz invariance operators and fields, that were explained in detial on section 4.2. Scalar feild equation, Klain-Gordon, presents a great idea how to develope theory of fermions. So we are intrested in a theory which is looks like Klein-Gordon theory, but the vecor field version of that one. In section 4.2, we learnd how to do Lorentz transformation for spin 1/2 particles in curves space-time, now we apply those trasfomations to derive Lorentz invariance theory.

For getting Dirac equation in curved space-time we need to find covariant derivative of for the spinor field ψ , in spin space. Dirac equation has the form of,

$$[\tilde{\gamma}^\mu D_\mu - m] \psi = 0 \quad (4.21)$$

where D_μ is covariant derivative of for the spinor field ψ , and for case which one have electrical field Dirac equation get the form of,

$$[i\gamma^\mu (D_\mu - ieA_\mu) - m] \psi = 0 \quad (4.22)$$

The covariant derivative for spinor is,

$$D_\mu \psi = \left(\partial_\mu + \frac{i}{4} \omega_\mu^{\alpha\beta} \sigma_{\alpha\beta} \right) \psi \quad (4.23)$$

where,

$$\sigma_{\alpha\beta} = \frac{1}{2i} [\gamma_\alpha, \gamma_\beta] \quad (4.24)$$

and $\omega_\mu^{\alpha\beta}$ has the form of,

$$\omega_\mu^{\alpha\beta} = \frac{1}{2} \left(v_\nu^\alpha \partial_\mu g^{\nu\nu'} v_{\nu'}^\beta + v_\nu^\alpha g^{\sigma\sigma'} v_{\sigma'}^\beta \Gamma_{\sigma\mu}^\nu - (\alpha \leftrightarrow \beta) \right) \quad (4.25)$$

where,

$$\Gamma_{\mu\nu}^{\lambda} = \frac{1}{2} g^{\lambda\rho} (\partial_{\mu} g_{\nu\rho} + \partial_{\nu} g_{\rho\mu} - \partial_{\rho} g_{\mu\nu}) \quad (4.26)$$

For diagonal metrics, we can simplify $\omega_{\mu}^{\alpha\beta}$ to,

$$\omega_{\mu}^{\alpha\beta} = \frac{1}{2} \left[v_{\alpha}^{\alpha} v_{\beta}^{\beta} g^{\alpha\alpha} g^{\beta\beta} (\partial_{\beta} g_{\alpha\mu} - \partial_{\alpha} g_{\beta\mu}) \right] \quad (4.27)$$

this simplified ω is true for all kind of diagonal metrics. From above equation it is obvious that for diagonal metric ω is anti-symmetric. Now, lets do the calculations for Charged Black Hole in AdS_5 . v_{β}^{α} is,

$$\begin{aligned} v_j^i &= 0 & i \neq j, & & v_0^0 &= \frac{\sqrt{f}r}{L} \\ v_i^i &= \frac{r}{L} & i = 1, 2, 3, & & v_5^5 &= \frac{L}{\sqrt{f}r} \end{aligned} \quad (4.28)$$

For this spce the only non-zero terms for ω are,

$$\omega_0^{05} = -\omega_0^{50} = \partial_r \frac{r^2 f}{2L^2}, \quad \omega_i^{i5} = -\omega_i^{5i} = \frac{\sqrt{f}r}{L^2} \quad i = 1, 2, 3 \quad (4.29)$$

and,

$$\begin{aligned} \sigma^{50} &= \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} = -\sigma^{05}, \\ \sigma^{5i} &= \begin{pmatrix} 0 & -\sigma^i \\ -\sigma^i & 0 \end{pmatrix} = -\sigma^{i5} \quad i = 1, 2, 3 \end{aligned} \quad (4.30)$$

4.3.1 Diran Equation solution for simple AdS_5 space

Dirac equation has the form of,

$$[\tilde{\gamma}^{\mu} D_{\mu} - m] \psi = 0 \quad (4.31)$$

where,

$$D_{\mu} \psi = \left(\partial_{\mu} + \frac{i}{4} \omega_{\mu}^{\alpha\beta} \sigma_{\alpha\beta} \right) \psi \quad (4.32)$$

and,

$$\omega_0^{50} = -\omega_0^{05} = \frac{r}{L^2}, \quad \omega_i^{5i} = -\omega_i^{i5} = \frac{r}{L^2} \quad i = 1, 2, 3 \quad (4.33)$$

and,

$$\begin{aligned} \sigma^{50} &= \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} = -\sigma^{05}, \\ \sigma^{5i} &= \begin{pmatrix} 0 & -\sigma^i \\ -\sigma^i & 0 \end{pmatrix} = -\sigma^{i5} \quad i = 1, 2, 3 \end{aligned} \quad (4.34)$$

so we get,

$$\begin{aligned} 0 &= \left[\tilde{\gamma}^0 \left(\partial_t + i \frac{r}{2L^2} \sigma^{50} \right) + \tilde{\gamma}^1 \left(\partial_x + i \frac{r}{2L^2} \sigma^{51} \right) + \tilde{\gamma}^2 \left(\partial_y + i \frac{r}{2L^2} \sigma^{52} \right) \right. \\ &\quad \left. + \tilde{\gamma}^3 \left(\partial_z + i \frac{r}{2L^2} \sigma^{53} \right) + \tilde{\gamma}^5 \partial_r - m \right] \psi \end{aligned} \quad (4.35)$$

ψ is a four component vector which represnet ... ?!. Lets derive the equations for each component and try to solve them,

$$\begin{aligned} 0 &= \left[i \frac{L}{r} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \left[\partial_t + i \frac{r}{2L^2} \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \right] \right. \\ &\quad + i \frac{L}{r} \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix} \left[\partial_x - i \frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix} \right] \\ &\quad + i \frac{L}{r} \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix} \left[\partial_y - i \frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix} \right] \\ &\quad + i \frac{L}{r} \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix} \left[\partial_z - i \frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix} \right] \\ &\quad \left. + \frac{r}{L} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \partial_r - m \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \right] \psi \end{aligned} \quad (4.36)$$

For the first step lets assume that we have a plane wave in x , y , z , and t direction, then we try to solve the equations for r and at the end we are able to use fourier trasformation to sum up all plane waves to get the whole answer. $\tilde{\psi}(\omega, k_i, r) = \exp \{-i\omega t + ik_i x_i\} \psi(\omega, k_i, r)$, which is a for component vector has the form of,

$$\begin{aligned}
0 = & \left[i\frac{L}{r} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \left[-i\omega + i\frac{r}{2L^2} \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} \right] \right. \\
& + i\frac{L}{r} \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix} \left[ik_x - i\frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^1 \\ \sigma^1 & 0 \end{pmatrix} \right] \\
& + i\frac{L}{r} \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix} \left[ik_y - i\frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix} \right] \\
& + i\frac{L}{r} \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix} \left[ik_z - i\frac{r}{2L^2} \begin{pmatrix} 0 & \sigma^3 \\ \sigma^3 & 0 \end{pmatrix} \right] \\
& \left. + \frac{r}{L} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \partial_r - m \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \right] \psi(\omega, k_i, r) \quad (4.37)
\end{aligned}$$

and then,

$$\begin{aligned}
0 = & \left[\frac{L}{r} \begin{pmatrix} 0 & \omega - k_i \sigma^i \\ \omega + k_i \sigma^i & 0 \end{pmatrix} + \frac{1}{L} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \right] \psi(P, r) \\
& + \left[\begin{pmatrix} (\frac{r}{L} \partial_r - m)I & 0 \\ 0 & -(\frac{r}{L} \partial_r + m)I \end{pmatrix} \right] \psi(\omega, k_i, r) \quad (4.38)
\end{aligned}$$

Finally we get,

$$\begin{aligned}
0 = & \left[\frac{1}{L} + \frac{r}{L} \partial_r - m \right] L \Psi_1 + \frac{L}{r} (\omega I - k_i \sigma^i) \Psi_2 \\
0 = & \left[-\frac{1}{L} - \frac{r}{L} \partial_r - m \right] L \Psi_2 + \frac{L}{r} (\omega I + k_i \sigma^i) \Psi_1 \quad (4.39)
\end{aligned}$$

Ψ is two component vector and its differential equation is,

$$0 = \left[-\frac{1}{L} - \frac{r}{L} \partial_r - m \right] \left[\frac{1}{L} + \frac{r}{L} \partial_r - m \right] L \Psi_1 - \frac{L}{r} (\omega I - k_i \sigma^i) \frac{L}{r} (\omega I + k_i \sigma^i) \Psi_1 \quad (4.40)$$

$$0 = \left[m^2 - \left(\frac{1}{2L} + \frac{r}{L} \partial_r \right)^2 - \frac{L^2}{r^2} (\omega^2 - k_i^2) \right] \Psi_1 \quad (4.41)$$

To make our equation simpler we can define $\psi(\omega, k_i, r) = \frac{L}{r} \phi(\omega, k_i, r)$ and plug it into equation 4.42, then we get,

$$0 = \left[\frac{L^2}{r^2} \begin{pmatrix} 0 & \omega - k_i \sigma^i \\ \omega + k_i \sigma^i & 0 \end{pmatrix} - \frac{Lm}{r} \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} \right] \phi(\omega, k_i, r) \\ + \left[\begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \right] \frac{\partial \phi(\omega, k_i, r)}{\partial r} \quad (4.42)$$

4.3.2 General Note on Diagonal Mertric which depends on just one variable

In this part we try to simplify the Dirac operator, say $\tilde{\gamma}^\mu D_\mu$ for diagonal metrics which depends on just one variable, say r . Lets assume that the metric has the form of,

$$ds^2 = -f_0^2(r)dt^2 + f_1^2(r)dx^2 + f_2^2(r)dy^2 + f_3^2(r)dz^2 + f_5^2(r)dr^2 \quad (4.43)$$

which, $f_i^2(r)$ is a function of r . Now we want to simplify the Dirac operator into something easier to work with. Dirac equation in curved space has the form of,

$$[\tilde{\gamma}^\mu (D_\mu - m)] \psi = 0 \quad (4.44)$$

which,

$$D_\mu \psi = \left(\partial_\mu + \frac{i}{4} \omega_\mu^{\alpha\beta} \sigma_{\alpha\beta} \right) \psi \quad (4.45)$$

and,

$$\sigma_{\alpha\beta} = \frac{1}{2i} [\gamma_\alpha, \gamma_\beta] \quad (4.46)$$

and $\omega_\mu^{\alpha\beta}$ for diagonal metric has the form of,

$$\omega_\mu^{\alpha\beta} = \frac{1}{2} \left[v_\alpha^\alpha v_\beta^\beta g^{\alpha\alpha} g^{\beta\beta} (\partial_\beta g_{\alpha\mu} - \partial_\alpha g_{\beta\mu}) \right] \quad (4.47)$$

this simplified ω is true for all kind of diagonal metrics. From above equation it is obvious that for diagonal metric ω is anti-symmetric. In our case v_β^α is,

$$v_j^i = 0 \quad i \neq j, \quad v_i^i = f_i(r) \quad i = 0, 1, 2, 3, 5 \quad (4.48)$$

Gamma matrices in Minkowski space are,

$$\begin{aligned}\gamma^0 &= i \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma^1 = i \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix}, \quad \gamma^2 = i \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \\ \gamma^3 &= i \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix}, \quad \gamma^5 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}\end{aligned}\quad (4.49)$$

and Gamma matrices in Curved space are,

$$\begin{aligned}\tilde{\gamma}^0 &= i \frac{1}{f_0(r)} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \tilde{\gamma}^1 = i \frac{1}{f_1(r)} \begin{pmatrix} 0 & \sigma^1 \\ -\sigma^1 & 0 \end{pmatrix}, \quad \tilde{\gamma}^2 = i \frac{1}{f_2(r)} \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, \\ \tilde{\gamma}^3 &= i \frac{1}{f_3(r)} \begin{pmatrix} 0 & \sigma^3 \\ -\sigma^3 & 0 \end{pmatrix}, \quad \tilde{\gamma}^5 = \frac{1}{f_5(r)} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}\end{aligned}\quad (4.50)$$

and non zero terms of ω are,

$$\omega_i^{j5} = -\omega_i^{5i} = \frac{f'_i(r)}{f_5(r)} \quad (4.51)$$

Then the Dirac operator gets the form of,

$$\tilde{\gamma}^\mu D_\mu = \tilde{\gamma}^0 \left(\partial_t + \frac{i}{2} \frac{f'_0}{f_5} \sigma_{05} \right) + \tilde{\gamma}^i \left(\partial_i + \frac{i}{2} \frac{f'_i}{f_5} \sigma_{i5} \right) + \tilde{\gamma}^r \partial_r \quad (4.52)$$

then we choose our spinor vector, $\psi = \exp(-i\omega_t + ik_j x^j) (-gg^{55})^{-1/4} \phi(r)$, then by applying Dirac operator on ψ we get,

$$\begin{aligned}\tilde{\gamma}^\mu D_\mu \psi &= \begin{pmatrix} 0 & \frac{\omega}{f_0} - \frac{k_i \sigma^i}{f_i} \\ \frac{\omega}{f_0} + \frac{k_i \sigma^i}{f_i} & 0 \end{pmatrix} \frac{\exp\{-i\omega t + ik_j x^j\}}{\sqrt{f_0 f_1 f_2 f_3}} \phi(r) \\ &\quad - \frac{1}{f_5} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \frac{\exp\{-i\omega t + ik_j x^j\}}{\sqrt{f_0 f_1 f_2 f_3}} \frac{\partial \phi(r)}{\partial r}\end{aligned}\quad (4.53)$$

4.3.3 Dirac equation in Charged Field

Now assume that there is charge in our Dirac equation so the Dirac Operator changes to,

$$\tilde{\gamma}^\mu D_\mu = \tilde{\gamma}^\mu \left[\partial_\mu + \frac{i}{4} \omega_\mu^{\alpha\beta} \sigma_{\alpha\beta} - iq A_\mu \right] \quad (4.54)$$

Note that in case which we are interested the Dirac action depends on q only through,

$$\mu_q \equiv \mu q \quad (4.55)$$

and from before we have,

$$\sigma_{\alpha\beta} = \frac{1}{2i} [\gamma_\alpha, \gamma_\beta] \quad (4.56)$$

where,

$$\gamma_\mu = \left\{ -\gamma^0, \gamma^1, \gamma^2, \gamma^3, \gamma^5 \right\} \quad (4.57)$$

Note that these equation are correct in case of, **sgn**(- + + + +). In this section we choose this signs to derive our equations. And $\omega_\mu^{\alpha\beta}$ for diagonal metric, lets has say $ds^2 = -f_0(r)dt^2 + \sum_{i=1}^3 f_i(r)dx_i^2 + f_5(r)dr^2$, the form of,

$$\omega_\mu^{\alpha\beta} = \frac{1}{2} \left[v_\alpha^\alpha v_\beta^\beta g^{\alpha\alpha} g^{\beta\beta} (\partial_\beta g_{\alpha\mu} - \partial_\alpha g_{\beta\mu}) \right] \quad (4.58)$$

this simplified ω is true for all kind of diagonal metrics. From above equation it is obvious that for diagonal metric ω is anti-symmetric. In our case v_β^α is,

$$v_j^i = 0 \quad i \neq j, \quad v_i^i = f_i(r) \quad i = 0, 1, 2, 3, 5 \quad (4.59)$$

Gamma matrices in Minkowski space, with **sgn**(- + + + +), are

$$\gamma^0 = i \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma^i = i \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad \gamma^5 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \quad (4.60)$$

spinor has two component, lets say ψ_\pm , and in our study we prefer to use the spinot with following definition to make our equation simpler,

$$\psi_\pm = (-g^{rr})^{-1/4} \times \exp \{ -i\omega t + ik_i x^i \} \phi_\pm \quad (4.61)$$

Non-vanishing terms for $\omega_\mu^{\alpha\beta}$ are,

$$\omega_i^{55} = -\omega_i^{5i} = \frac{f'_i}{f_5}, \quad i = 0, 1, 2, 3 \quad (4.62)$$

and non-vanishing terms for $\sigma_{\alpha\beta}$ are,

$$\sigma_{05} = -\sigma_{50} = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \quad \sigma_{i5} = -\sigma_{5i} = \begin{pmatrix} 0 & -\sigma^i \\ -\sigma^i & 0 \end{pmatrix} \quad i = 1, 2, 3 \quad (4.63)$$

so in this case dirac operator get the form of,

$$\begin{aligned}\tilde{\gamma}^\mu D_\mu = & \frac{i}{f_0} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \left[\partial_t + \frac{i}{2} \frac{f'_0}{f_5} \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix} - iqA_t \right] + \frac{\partial_r}{f_5} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \\ & + \sum_{i=1}^3 \frac{i}{f_i} \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix} \left[\partial_i - \frac{i}{2} \frac{f'_i}{f_5} \begin{pmatrix} 0 & \sigma^i \\ \sigma^i & 0 \end{pmatrix} \right]\end{aligned}\quad (4.64)$$

after applying Dirac operator on spinor, equation 4.61, we get,

$$\begin{aligned}\tilde{\gamma}^\mu D_\mu \psi = & \begin{pmatrix} 0 & \frac{\omega + qA_t}{f_0} - \frac{k_i \sigma^i}{f_i} \\ \frac{\omega + qA_t}{f_0} + \frac{k_i \sigma^i}{f_i} & 0 \end{pmatrix} (-gg^{rr})^{-\frac{1}{4}} \times \exp\{-i\omega t + ik_i x^i\} \phi(r) \\ & + \frac{1}{f_5} \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} (-gg^{rr})^{-\frac{1}{4}} \times \exp\{-i\omega t + ik_i x^i\} \frac{\partial \phi(r)}{\partial r}\end{aligned}\quad (4.65)$$

Now we are able to write the Dirac equation with mass. After adding mass to our equation and cross $(-gg^{rr})^{-\frac{1}{4}} \times \exp\{-i\omega t + ik_i x^i\}$ from our spinor, we get,

$$0 = \begin{pmatrix} \frac{\partial_r}{f_5} - m & \frac{\omega + qA_t}{f_0} - \frac{k_i \sigma^i}{f_i} \\ \frac{\omega + qA_t}{f_0} + \frac{k_i \sigma^i}{f_i} & -\frac{\partial_r}{f_5} - m \end{pmatrix} \begin{pmatrix} \phi_+ \\ \phi_- \end{pmatrix}\quad (4.66)$$

lets define $\omega + qA_t = \omega + q(1 - (\frac{r_0}{r})^2) \equiv \Omega$ and solve the equation 4.66 for ϕ_\pm ,

$$\begin{aligned}0 &= \left(\frac{\partial_r}{f_5} - m \right) \phi_+ + \left(\frac{\Omega}{f_0} - \frac{k_i \sigma^i}{f_i} \right) \phi_- \\ 0 &= \left(-\frac{\partial_r}{f_5} - m \right) \phi_- + \left(\frac{\Omega}{f_0} + \frac{k_i \sigma^i}{f_i} \right) \phi_+\end{aligned}\quad (4.67)$$

we solved the equation in Chiral representation we can solve the equation for other representations, too. So in general one can prove that the equations has the form of,

$$\sqrt{\frac{g_{ii}}{g_{rr}}} (\partial_r \mp m \sqrt{g_{rr}}) \phi_\pm = \mp i K_\mu \gamma^\mu \phi_\mp\quad (4.68)$$

with,

$$K_\mu(r) = (-u(r), k_i), \quad u = \sqrt{\frac{g_{ii}}{-g_{tt}}} \left(\omega + \mu_q (1 - (\frac{r_0}{r})^2) \right)\quad (4.69)$$

4.4 solving the Dirac equation

In this section we try to find the analytical solution of Dirac equation near horizon and boundary of black hole in AdS space. The geometry is,

$$ds^2 = -\frac{r^2 f(r)}{L^2} dt^2 + \frac{r^2}{L^2} dx^2 + \frac{r^2}{L^2} dy^2 + \frac{r^2}{L^2} dz^2 + \frac{L^2}{r^2 f(r)} dr^2 \quad (4.70)$$

with,

$$f = 1 + \frac{Q^2}{r^{2d-2}} - \frac{M}{r^d} \quad (4.71)$$

for spinor $\psi_{\pm} = \frac{L^2}{r^2 f^{1/4}(r)} \times \exp \{-i\omega t + ik_i x^i\} \phi(r)_{\pm}$ and $\omega + q(1 - (\frac{r_0}{r})^2) \equiv \Omega$ we have,

$$\begin{aligned} 0 &= \left(\frac{\partial_r}{f_5} - m \right) \phi_+ + \left(\frac{\Omega}{f_0} - \frac{k_i \sigma^i}{f_i} \right) \phi_- \\ 0 &= \left(-\frac{\partial_r}{f_5} - m \right) \phi_- + \left(\frac{\Omega}{f_0} + \frac{k_i \sigma^i}{f_i} \right) \phi_+ \end{aligned} \quad (4.72)$$

for black hole geometry in AdS_5 one gets,

$$\begin{aligned} 0 &= \left(\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_+ + \left(\frac{L\Omega}{r\sqrt{f(r)}} - \frac{Lk_i \sigma^i}{r} \right) \phi_- \\ 0 &= \left(-\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_- + \left(\frac{L\Omega}{r\sqrt{f(r)}} + \frac{Lk_i \sigma^i}{r} \right) \phi_+ \end{aligned} \quad (4.73)$$

Near boundary, $r \rightarrow \infty$ up to second order answer has the form of,

$$\phi_+(r) = Ar^{mL} + Br^{-mL-1} + \dots, \quad \phi_-(r) = Cr^{-mL} + Dr^{mL-1} + \dots \quad (4.74)$$

with,

$$B = \frac{L^2 \Omega - L^2 k_i \sigma^i}{2m+1} C, \quad D = \frac{L^2 \Omega + L^2 k_i \sigma^i}{2m-1} A \quad (4.75)$$

and close to horizon, $r \rightarrow r_0$ where $f(r_0) = 0$, we get,

$$\begin{aligned}
0 &= \partial_r [(r - r_0) \partial_r \phi_+] + \frac{L^4 \Omega^2(r_0)}{\kappa^2 r_0^2 (r - r_0)} \phi_+ \\
0 &= (r - r_0) \partial_r \phi_+ + \frac{L^2 \Omega(r_0)}{r_0^4 \kappa} \phi_-
\end{aligned} \tag{4.76}$$

with,

$$\kappa = 4 \frac{r_0^6 - r_*^6}{r_0^7} \tag{4.77}$$

and the answer has the form of,

$$\begin{aligned}
\phi_+(r) &= A \exp\{i\beta \ln(r - r_0)\} + B \exp\{-i\beta \ln(r - r_0)\}, \\
\phi_-(r) &= -iA \exp\{i\beta \ln(r - r_0)\} + iB \exp\{-i\beta \ln(r - r_0)\}
\end{aligned} \tag{4.78}$$

with,

$$\beta = \frac{L^2 \omega}{\kappa r_0} \tag{4.79}$$

Because we are interested in retarded Green function so by applying the in-falling boundary condition at horizon we are set. Then in equation 4.78 we ignore the second term, $\exp\{-i\beta \ln(r - r_0)\}$ so we will use $A = 1$ and $B = 0$ to get an appropriate boundary condition.

4.4.1 Green function

If the relation between coefficient C (which is corresponded to expectation value) and A (which is corresponded to source) in equation 4.74 be $C = \mathcal{S}A$, then the Green function, say G_R , has the form of,

$$G_R = -i\mathcal{S}\gamma^0 \tag{4.80}$$

By having the boundary condition, equation 4.78, and then setting the in-falling boundary condition, say $B = 0$, one can find the Green function 4.80 by integrating equation 4.73 over r and finding the boundary correlation function and calculate the Green function directly.

Properties of Green function

Lets divide the spinors ϕ_+ and ϕ_- into their components, say ϕ_{++} , ϕ_{+-} , ϕ_{-+} , and ϕ_{--} , and without loosing the generality assume that $k_1 = k_2 = 0$ then we can rewrite equation 4.73 and get,

$$\begin{aligned}
 0 &= \left(\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_{++} + \left(\frac{L\Omega}{r\sqrt{f(r)}} - \frac{Lk_3}{r} \right) \phi_{-+} \\
 0 &= \left(\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_{+-} + \left(\frac{L\Omega}{r\sqrt{f(r)}} + \frac{Lk_3}{r} \right) \phi_{--} \\
 0 &= \left(-\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_{-+} + \left(\frac{L\Omega}{r\sqrt{f(r)}} + \frac{Lk_3}{r} \right) \phi_{++} \\
 0 &= \left(-\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_{--} + \left(\frac{L\Omega}{r\sqrt{f(r)}} - \frac{Lk_3}{r} \right) \phi_{+-} \quad (4.81)
 \end{aligned}$$

By changing k_3 into $-k_3$ in equation

....

4.4.2 Zero Temperature

When $r_* = r_0$ then we gets zero temprature. We defined the zero temperature in last chapter so here we just try to derive its formulation for Dirac equations. Note that in zero temprature only redshift factor $f(r)$, or $f(z)$ changing, lets define $z = r_0/r$. So we in r coordinate one have Dirac equations,

$$\begin{aligned}
 0 &= \left(\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_+ + \left(\frac{L\Omega}{r\sqrt{f(r)}} - \frac{Lk_i\sigma^i}{r} \right) \phi_- \\
 0 &= \left(-\frac{r\sqrt{f(r)}\partial_r}{L} - m \right) \phi_- + \left(\frac{L\Omega}{r\sqrt{f(r)}} + \frac{Lk_i\sigma^i}{r} \right) \phi_+ \quad (4.82)
 \end{aligned}$$

or in z coordinate,

$$\begin{aligned}
 0 &= \left(-\frac{\sqrt{f(z)}\partial_z}{zL} - m \right) \phi_+ + \left(\frac{zL\Omega}{r_0\sqrt{f(z)}} - \frac{zLk_i\sigma^i}{r_0} \right) \phi_- \\
 0 &= \left(-\frac{\sqrt{f(z)}\partial_z}{zL} - m \right) \phi_- + \left(\frac{zL\Omega}{r_0\sqrt{f(z)}} + \frac{zLk_i\sigma^i}{r_0} \right) \phi_+ \quad (4.83)
 \end{aligned}$$

with,

$$f = \frac{(r-r_0)^2(r+r_0)^2(r^2+2r_0^2)}{r^6} \quad (4.84)$$

or,

$$f = (1-z)^2(1+z)^2(1+2z^2) \quad (4.85)$$

so close to boundary, $r \rightarrow \infty$, answer has the form of,

$$\phi_+(r) = Ar^{mL} + Br^{-mL-1}, \quad \phi_-(r) = Cr^{-mL} + Dr^{mL-1} \quad (4.86)$$

or for z coordinate,

$$\phi_+(r) = A \left(\frac{r_0}{z} \right)^{mL} + B \left(\frac{r_0}{z} \right)^{-mL-1}, \quad \phi_-(r) = C \left(\frac{r_0}{z} \right)^{-mL} + D \left(\frac{r_0}{z} \right)^{mL-1} \quad (4.87)$$

with,

$$B = \frac{L^2\Omega - L^2k_i\sigma^i}{2m+1}C, \quad D = \frac{L^2\Omega + L^2k_i\sigma^i}{2m-1}A \quad (4.88)$$

and close to horizon, when $r \rightarrow r_0$, the differential equations gets the form of,

$$\begin{aligned} 0 &= \partial_r \left[(r-r_0)^2 \partial_r \phi_+ \right] + \frac{r_0^4 L^4 \Omega^2 (r_0)}{144 (r-r_0)^2} \phi_+ \\ 0 &= (r-r_0)^2 \partial_r \phi_+ + \frac{r_0^2 L^2 \Omega (r_0)}{12} \phi_- \end{aligned} \quad (4.89)$$

and the answer has the form of,

$$\begin{aligned} \phi_+(r) &= A \exp \left\{ \frac{i\theta}{(r-r_0)} \right\} + B \exp \left\{ -\frac{i\theta}{(r-r_0)} \right\}, \\ \phi_-(r) &= iA \exp \left\{ \frac{i\theta}{(r-r_0)} \right\} - iB \exp \left\{ -\frac{i\theta}{(r-r_0)} \right\} \end{aligned} \quad (4.90)$$

with,

$$\theta = \frac{L^2 \omega}{12} \quad (4.91)$$

4.4.3 Looking more carefully at zero temprature and field without mass

In this part we want to look at zero temprature and field with zero mass more carefully, So the Dirac euqation simplifies to,

$$\begin{aligned} 0 &= \left(\frac{r\sqrt{f(r)}\partial_r}{L} \right) \phi_+ + \left(\frac{L\Omega}{r\sqrt{f(r)}} - \frac{Lk_i\sigma^i}{r} \right) \phi_- \\ 0 &= \left(-\frac{r\sqrt{f(r)}\partial_r}{L} \right) \phi_- + \left(\frac{L\Omega}{r\sqrt{f(r)}} + \frac{Lk_i\sigma^i}{r} \right) \phi_+ \end{aligned} \quad (4.92)$$

with,

$$f = \frac{(r-r_0)^2(r+r_0)^2(r^2+2r_0^2)}{r^6} \quad (4.93)$$

now we want to look at equations of near horizon more carefully and derive more terms for our approximation, we assume that the outgoing boundary condition is zero so in general the answer has the form of,

$$\begin{aligned} \phi_+(r) &= A \exp \left\{ \frac{i\theta}{(r-r_0)} \right\} [1 + a_{1+}(r-r_0) + \dots], \\ \phi_-(r) &= iA \exp \left\{ \frac{i\theta}{(r-r_0)} \right\} [1 + a_{1-}(r-r_0) + \dots] \end{aligned} \quad (4.94)$$

with $\Omega \equiv \omega + \mu_q(1 - (\frac{r_0}{r})^2)$ and $\theta = \frac{L^2\omega}{12}$. Then we can calculate coefficients of a_{1+} and a_{1-} . By plugging equation 4.94 into Dirac equation , one gets,

$$\begin{aligned} -\frac{12\theta}{L} a_{1+} + L\omega a_{1-} &= -\frac{28\theta}{r_0 L} - 2L\mu_q + \frac{2Lk_3\sigma^3\sqrt{3}}{r_0} \\ L\omega a_{1+} - \frac{12\theta}{L} a_{1-} &= -\frac{28\theta}{r_0 L} - 2L\mu_q - \frac{2Lk_3\sigma^3\sqrt{3}}{r_0} \end{aligned} \quad (4.95)$$

and then,

$$\begin{aligned}
a_{1+} &= \frac{-\frac{12\theta}{L} \left(-\frac{28\theta}{r_0 L} - 2L\mu_q + \frac{2Lk_3\sigma^3\sqrt{3}}{r_0} \right) - L\omega \left(-\frac{28\theta}{r_0 L} - 2L\mu_q - \frac{2Lk\sqrt{3}}{r_0} \right)}{\frac{144\theta^2}{L^2} - L^2\omega^2} \\
a_{1-} &= \frac{-L\omega \left(-\frac{28\theta}{r_0 L} - 2L\mu_q + \frac{2Lk_3\sigma^3\sqrt{3}}{r_0} \right) - \frac{12\theta}{L} \left(-\frac{28\theta}{r_0 L} - 2L\mu_q - \frac{2Lk\sqrt{3}}{r_0} \right)}{\frac{144\theta^2}{L^2} - L^2\omega^2}
\end{aligned} \tag{4.96}$$

And near boundary the general answer has the form of,

$$\phi_+(r) = A + A_1 r^{-1} + A_2 r^{-2} + \dots, \quad \phi_-(r) = C + C_1 r^{-1} + C_2 r^{-2} + \dots \tag{4.97}$$

with,

$$\begin{aligned}
A_1 &= L^2(\omega + \mu_q - k_i \sigma^i)C, & C_1 &= -L^2(\omega + \mu_q + k_i \sigma^i)A \\
A_2 &= L^2 \frac{\omega + \mu_q}{2} C_1, & C_2 &= -L^2 \frac{\omega + \mu_q}{2} A_1
\end{aligned} \tag{4.98}$$

Chapter 5

Spin 3/2 particles

5.1 Introduction

5.2 Rarita-Schwinger Formalism

Spin 3/2 particles represented by Rarita-Schwinger field that is a tensor product between the first rank tensor and Dirac bispinor. Dirac bispinor can be denoted by $SU(2) \oplus SU(2)$ representation in the usual way,

$$\psi : (\frac{1}{2}, 0) \oplus (0, \frac{1}{2}) \quad (5.1)$$

A spin 1-field or Lorentz vector A_μ can be constructed as the direct product $(\frac{1}{2}, 0) \otimes (0, \frac{1}{2})$ by following,

$$A_\mu : (\frac{1}{2}, 0) \otimes (0, \frac{1}{2}) = (\frac{1}{2}, \frac{1}{2}) \quad (5.2)$$

The spin 3/2 vector spinor ψ_μ , according to Rarita-Schwinger can be written as,

$$\psi_\mu \sim A_\mu \otimes \psi : (\frac{1}{2}, \frac{1}{2}) \otimes \left((\frac{1}{2}, 0) \oplus (0, \frac{1}{2}) \right) \quad (5.3)$$

Rarita and Schwinger proposed an equation to describe free spin 3/2 particle as following,

$$(i\gamma^\mu \partial_\mu - m)\psi_V(x) = 0 \quad (5.4)$$

with two constrains,

$$\gamma^\mu \psi_\mu = 0 \quad \text{and} \quad \partial^\mu \psi_\mu = 0 \quad (5.5)$$

This constraints allow us to cancel the extra degree of freedom of our theory.

5.3 Equation of motion for 3/2 spin particles

A spin 3/2 field can be described by the Rarita-Schwinger Lagrangian given by the following formalism,

$$\mathcal{L}(A) = \bar{\psi}^\mu (i\partial_\alpha \Gamma_\mu^\alpha{}_\nu(A) - mB_{\mu\nu}(A)) \psi^\nu, \quad (5.6)$$

where $\bar{\psi}^\mu = (\psi^\mu)^\dagger \gamma^0$ is the adjoint vector spinor, m is the particle mass, ψ^μ is the vector spinor and the matrices $\Gamma_\mu^\alpha{}_\nu$ and $B_{\mu\nu}$ are given by,

$$\Gamma_\mu^\alpha{}_\nu(A) = g_{\mu\nu}\gamma^\alpha + A(\gamma_\mu g_\nu^\alpha + B\gamma_\mu\gamma^\alpha\gamma_\nu), \quad (5.7)$$

and

$$B_{\mu\nu}(A) = g_{\mu\nu} - C\gamma_\mu\gamma_\nu \quad (5.8)$$

with

$$B \equiv \frac{3}{2}A^2 + A + \frac{1}{2} \quad (5.9)$$

and

$$C \equiv 3A^2 + 3A + 1. \quad (5.10)$$

in case of Minkowski space and **Sgn**(-+++), γ^μ are Dirac matrices and A is an arbitrary parameter (except that $A \neq -1/2$). The parameter A has no physical meaning and the conventional choices made in the literature are $A = -1/3$, or $A = 0$, or $A = -1$. For $A = -1$ the Lagrangian reduces to:

$$\mathcal{L} = -\frac{i}{2} \bar{\psi}_\mu (\varepsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\mu \partial_\rho + m\sigma^{\mu\sigma}) \psi_\sigma \quad (5.11)$$

where $\varepsilon^{\mu\nu\rho\sigma}$ is the Levi-Civita symbol, $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$, and $\sigma^{\mu\sigma} = \frac{i}{2}[\gamma^\mu, \gamma^\sigma]$. By applying the Euler-Lagrangian equation to the Lagrangian, the equation of motion gets the form of,

$$(\varepsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\mu \partial_\rho + m\sigma) \psi_\sigma = 0 \quad (5.12)$$

while the above equation can be written in a simpler form plus supplementary conditions

$$(i\gamma^\mu \partial_\mu - m)\psi_\nu = 0 \quad (5.13)$$

with conditions,

$$\gamma^\nu \psi_\mu = 0 \quad (5.14)$$

$$\partial^\mu \psi_\mu = 0 \quad (5.15)$$

These requirements lead to the Rarita-Schwinger equation given [Rarita-Schwinger paper] by,

$$(\gamma_\mu \partial^\mu + M) \psi^\mu(X) = 0 \quad (5.16)$$

with condition,

$$\gamma_\mu \psi^\mu = 0 \quad (5.17)$$

The completely symmetric function $\Psi_{\alpha\beta\gamma}(x)$ can be built from $\psi^\mu(x)$. As it is seen all of the vector component of $\psi^\mu(x)$ satisfy Dirac-type equations, however they are not independent. they must fullfil the constraint given by equation 5.17. This constraint coming from the symmetry condition projects out to spin 1/2 component and the remaining components describe particles with spin 3/2.

Here we use the Einstein summation rule. The Dirac matrices, as we have seen in last chapter, γ_μ satisfy the anticommuter relation,

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2\delta_{\mu\nu} \quad (5.18)$$

5.4 RaritaSchwinger equation

In theoretical physics, the RaritaSchwinger equation is the relativistic field equation of spin-3/2 fermions. It is similar to the Dirac equation for spin-1/2 fermions. This equation was first introduced by William Rarita and Julian Schwinger in 1941. In modern notation it can be written as:

$$(\varepsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\nu \partial_\rho + m \sigma^{\mu\sigma}) \psi_\sigma = 0 \quad (5.19)$$

where $\varepsilon^{\mu\nu\rho\sigma}$ is the Levi-Civita symbol, γ_5 and γ_ν are Dirac matrices, m is the mass, $\sigma^{\mu\nu} \equiv i/2 [\gamma^\mu, \gamma^\nu]$, and ψ_σ is a vector-valued spinor with additional components compared to the four component spinor in the Dirac equation.

It corresponds to the $(\frac{1}{2}, \frac{1}{2}) \otimes ((\frac{1}{2}, 0) \oplus (0, \frac{1}{2}))$ representation of the Lorentz group, or rather, its $(1, \frac{1}{2}) \oplus (\frac{1}{2}, 1)$ part. This field equation can be derived as the EulerLagrange equation corresponding to the Rarita-Schwinger Lagrangian:

$$\mathcal{L} = -\frac{i}{2} \bar{\psi}_\mu (\epsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\nu \partial_\rho + m \sigma^{\mu\sigma}) \psi_\sigma \quad (5.20)$$

where the bar above ψ_μ denotes the Dirac adjoint.

This equation is useful for the wave function of composite objects such as the delta baryons Δ or for the hypothetical gravitino. So far, no elementary particle with spin 3/2 has been found experimentally.

The massless RaritaSchwinger equation has a gauge symmetry, under the gauge transformation of $\psi_\mu \rightarrow \psi_\mu + \partial_\mu \epsilon$, where ϵ is an arbitrary spinor field.

5.4.1 RaritaSchwinger equation in 4-dimentional Minkowskispac

In 4-dimentional Minkowski sapce, Dirac matrices are obeying the following anti-symmetric rule,

$$\{\gamma_\mu, \gamma_\nu\} = 2\eta_{\mu\nu} I_{4 \times 4} \quad (5.21)$$

ψ_σ has for component, $\sigma = 0, 1, 2, 3$, and each component has 4 element, lets write the equations for each component in Minkowski space and solve it. For Simplicity assume that $m = 0$ and we have a wave equation which is traveling in x_3 direction the wave has the form of,

$$\psi_\mu(X) = \exp\{-i\omega t + ik_3 x_3\} \phi_\mu(\omega, k_3) \quad (5.22)$$

lets define $\gamma_{ijk} \equiv \gamma_i \gamma_j \gamma_k$, and $\gamma_{ij} \equiv \gamma_i \gamma_j$, then for we get four equations,

$$\begin{aligned} -\gamma_{51} k_3 \phi_2 + \gamma_{52} k_3 \phi_1 &= 0 \quad \rightarrow \quad \phi_2 = \gamma_{12} \phi_1 \\ -\gamma_{52} \omega \phi_3 - \gamma_{52} k_3 \phi_0 + \gamma_{53} \omega \phi_2 + \gamma_{50} k_3 \phi_2 &= 0 \\ \gamma_{51} \omega \phi_3 - \gamma_{50} k_3 \phi_1 - \gamma_{53} \omega \phi_1 + \gamma_{51} k_3 \phi_0 &= 0 \end{aligned} \quad (5.23)$$

By choosing the **Sgn**(- + + +) we get,

$$\begin{aligned}
\phi_2 &= \gamma_{12} \phi_1 \\
\phi_3 &= -\frac{k_3}{\omega} \phi_1 \\
\gamma_1 \phi_0 &= (\gamma_0 + \gamma_1 + \gamma_3 \frac{\omega}{k_3}) \phi_1 \\
(\gamma_3 \omega + \gamma_0 k_3) \phi_1 &= 0
\end{aligned} \tag{5.24}$$

For holding the above equations one needs to get $k_3 = \pm \omega$, which based on Special relativity point of view this is correct. Lets assume that $k_3 = \omega$ and move to the next part. Lets choose Chiral representation for Gamma matrices and solve the equations. Once we have ϕ_1 , ϕ_2 , ϕ_3 , and ϕ_0 are not arbitrary any more. We have,

$$(\gamma_3 \omega + \gamma_0 k_3) \phi_1 = 0 \tag{5.25}$$

and we should find the eigenvector of this equation,

$$\gamma_0 \gamma_3 \omega \phi_1 = k_3 \phi_1 \tag{5.26}$$

and in Chiral representation we have,

$$\left[\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \right] \phi_1 = \phi_1 \tag{5.27}$$

so the eigenvectors for both $\omega = k_3$ and $\omega = -k_3$ are,

$$\phi_1 = \phi_{1a} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + \phi_{1b} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + \phi_{1c} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \phi_{1d} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \tag{5.28}$$

Now lets assume that we have some mass and compute the whole thing one more time. Our equation of motion has the form of,

$$(\epsilon^{\mu\nu\rho\sigma} \gamma_5 \gamma_\nu \partial_\rho + m_1 \sigma^{\mu\sigma} + im_2 g^{\mu\sigma}) \psi_\sigma = 0 \tag{5.29}$$

and we get,

$$\begin{aligned}
0 &= -\gamma_{51}k_3\phi_2 + \gamma_{52}k_3\phi_1 + m_1\gamma^{01}\phi_1 + m_1\gamma^{02}\phi_2 + m_1\gamma^{03}\phi_3 - m_2\phi_0 \\
0 &= -\gamma_{52}\omega\phi_3 - \gamma_{52}k_3\phi_0 + \gamma_{53}\omega\phi_2 + \gamma_{50}k_3\phi_2 + m_1\gamma^{10}\phi_0 + m_1\gamma^{12}\phi_2 + m_1\gamma^{13}\phi_3 + m_2\phi_1 \\
0 &= \gamma_{51}\omega\phi_3 - \gamma_{50}k_3\phi_1 - \gamma_{53}\omega\phi_1 + \gamma_{51}k_3\phi_0 + m_1\gamma^{20}\phi_0 + m_1\gamma^{21}\phi_1 + m_1\gamma^{23}\phi_3 + m_2\phi_2 \\
0 &= -\gamma_{51}\omega\phi_2 + \gamma_{52}\omega\phi_1 + m_1\gamma^{30}\phi_0 + m_1\gamma^{31}\phi_1 + m_1\gamma^{32}\phi_2 + m_2\phi_3
\end{aligned} \tag{5.30}$$

or,

$$\begin{aligned}
& -m_2 \begin{pmatrix} -\phi_0 \\ \phi_1 \\ \phi_2 \\ \phi_2 \end{pmatrix} = \\
& \begin{pmatrix} 0 & \gamma_{52}k_3 + m_1\gamma^{01} & m_1\gamma^{02} - \gamma_{51}k_3 & m_1\gamma^{03} \\ \gamma_{52}k_3 - m_1\gamma^{10} & 0 & \gamma_{53}\omega + \gamma_{50}k_3 + m_1\gamma^{12} & m_1\gamma^{13} - \gamma_{52} \\ -\gamma_{51}k_3 - m_1\gamma^{20} & m_1\gamma^{21} - \gamma_{50}k_3 - \gamma_{53}\omega & 0 & \gamma_{51}\omega + m_1\gamma^{23} \\ -m_1\gamma^{30} & m_1\gamma^{31} + \gamma_{52}\omega & m_1\gamma^{32} - \gamma_{51}\omega & 0 \end{pmatrix} \\
& \times \begin{pmatrix} -\phi_0 \\ \phi_1 \\ \phi_2 \\ \phi_2 \end{pmatrix}
\end{aligned} \tag{5.31}$$

5.5 Spin 3/2 in curved space

In the $d+1$ -dimensional curved space RaritaSchwinger action has the form of,

$$I = \int d^{d+1}x \sqrt{g} \bar{\psi}_\mu [\Gamma^{\mu\nu\sigma} D_\nu - m_1 \Gamma^{\mu\sigma} - m_2 g^{\mu\sigma}] \psi_\sigma \tag{5.32}$$

D_ν is the covariant derivative, and Γ_μ 's are curved space Dirac matrices so that $\gamma_\mu = e_\mu^a \gamma_a$ where γ_a is the Dirac matrix in Minkowski space with relation $\{\gamma_a, \gamma_b\} = 2\delta_{ab}$. $\Gamma^{\mu\nu\sigma} \equiv \Gamma^{[\mu} \Gamma^\nu \Gamma^{\sigma]}$ and $\Gamma^{\mu\sigma} \equiv \Gamma^{[\mu} \Gamma^{\sigma]}$. The action of 5.32 gets the equation of motion in the form of,

$$[\Gamma^{\mu\nu\sigma} D_\nu - m_1 \Gamma^{\mu\sigma} - m_2 g^{\mu\sigma}] \psi_\sigma = 0 \tag{5.33}$$

and for its conjugate one gets,

$$\bar{\psi}_\mu \left[\Gamma^{\mu\nu\sigma} \overleftarrow{D}_\nu + m_1 \Gamma^{\mu\sigma} + m_2 g^{\mu\sigma} \right] = 0 \tag{5.34}$$

and we can show that by defining $m_\pm = m_1 \pm m_2$, equation 5.33 can be simplified to,

$$\Gamma^\nu [D_\nu \psi_\mu - D_\mu \psi_\nu] + \frac{m_+}{d-1} \Gamma_\mu \Gamma^\nu \psi_\nu - m_- \psi_\mu = 0 \quad (5.35)$$

Chapter 6

NOT RELATED TO THE PROJECT

This section is not part of the project. Just self notes, keeping for future.

6.1 SELF NOTES FOR MY RECORD

6.1.1 Is the nature Lorentz Invariance ?

This is just I found in media, and I write here just to have it and in future read more about it. This is not part of the notes.

<http://physics.indiana.edu/kostelec/faq.html>

6.1.2 Black Rings in 5-dimension

The Neutral Black Ring in 5-dimension EMD theory has been given by [R. Emparan, JHEP. 03 (2004) 064]:

$$ds^2 = -\frac{F(y)}{F(x)} \left(dt - C(v, \lambda) R \frac{1+y}{F(y)} d\psi \right)^2 - \frac{R^2}{(x-y)^2} F(x) \left[-\frac{G(y)}{F(y)} d\psi^2 - \frac{dy^2}{G(y)} + \frac{dx^2}{G(x)} + \frac{G(x)}{F(x)} d\phi^2 \right] \quad (6.1)$$

where

$$F(\xi) = 1 + \lambda \xi \quad , \quad G(\xi) = (1 - \xi^2)(1 + \nu \xi)$$

$$G(\nu, \lambda) = \sqrt{\lambda(\lambda - \nu) \frac{1 + \lambda}{1 - \lambda}} \quad (6.2)$$

The parameter λ and ν are dimensionless and takes values in the range ($0 < \nu \leq \lambda < 1$), and to avoid the conical singularity also at $x = 1$, λ and ν must be related to each other via $\lambda = 2\nu/(1 + \nu^2)$. The coordinate ϕ and ψ are two cycles of the black ring, and x and y takes the range as $-1 \leq x \leq 1$ and $-\infty \leq y \leq -1$. The constant R has the dimensional of length and for thin large rings corresponds roughly to the radius of the ring circle [H. Elvang, Phys. Rev. D68 (2008) 124016]. The horizon is located at $y = y_h = -1/\nu$. The mass of the black ring is $M = 3\pi R^2 \lambda / 4[1 - \nu]$, and its angular momentum takes $J = \pi R^3 \sqrt{\lambda(\lambda - \nu)(1 + \lambda)}/[2(1 - \nu)^2]$. In addition, the spacetime contains three Killing coordinate t , ϕ , and ψ .

Bibliography