

Quick Intro to Probability

Data 3402- Lecture 6

Amir Farbin

Probability Theory




- Experiments
 - Observations
 - Measurements
- Theory
 - Hypothesis
 - Models
- Quantify Uncertainty

Probability Theory

- Probability Theory → Analyze frequency of “events”
- Probability p of event x happening
 - → Repeatable observations of event
 - → $p \sim$ fraction that x would be the outcome.
 - Example: Frequency of a disease in a population: 1 in 1000 → “Frequentist”
- What if you got a test → How do you interpret? You can’t repeat.
 - Accuracy of test
 - Frequency of disease
 - → Degree of belief → “Bayesian”

Basic Definitions

- Set S w subsets A and B
 - $P(S) = 1$
 - For all $\forall A : A \subset S \Rightarrow P(A) > 0$
 - $P(\bar{A}) = 1 - P(A)$
 - Bar means not in A .
 - $P(A \cup \bar{A}) = 1$
 - $P(\emptyset) = 0$
 - If there is no overlap between sets A and B
 - $A \cap B = \emptyset \Rightarrow P(A \cup B) = P(A) + P(B)$
 - \cup is overlap
 - \cap is union
 - $A \subset B \Rightarrow P(A) \leq P(B)$
 - \subset means subset
 - $P(A \cup B) = P(A) + P(B) - P(A \cap B)$



Conditional Probability

- $P(A|B) = \frac{P(A \cap B)}{P(B)}$
- Example:
 - Dice Roll (6-sided)
 - You are told rolled 3 or less
 - What is the Probability that you rolled a 1?
 - $P(1) = \frac{P(<3 \text{ in } n \text{ events})}{P(\text{Events})} = \frac{\frac{1}{3}}{\frac{3}{6}} = \frac{1}{3}.$
- If $A \cup B = \emptyset \Rightarrow$
 - $P(A, B) = P(A)P(B)$
 - $P(A|B) = \frac{P(A)P(B)}{P(B)} = P(A)$

Interpretation

- Relative Frequency:
 - A, B are outcomes of a repeatable experiment
 - $P(A) = \lim_{n \rightarrow \infty} \frac{\text{Times outcome is } A}{n}$
- Subjective Probability:
 - A, B are hypotheses (True/False) Statements.
 - $P(A)$ is degree of belief that A is true.

Bayes Theorem

- $P(A|B) = \frac{P(A \cap B)}{P(B)}$
- $P(B|A) = \frac{P(B \cap A)}{P(A)}$
- Since $P(A \cap B) = P(B \cap A)$
 - $\Rightarrow P(A|B)P(B) = P(B|A)P(A)$
 - $\Rightarrow P(A|B) = \frac{P(B|A)P(A)}{P(B)}$

Bayes Example

Recall $\Rightarrow P(A|B) = \frac{P(B|A)P(A)}{P(B)}$

- Example: Disease Testing
 - Prior Knowledge about Population
 - $A := \{\text{ sick, not sick} \}$
 - $P(\text{sick}) = 0.001$
 - $P(\text{not sick}) = 0.999$
 - Test
 - $B := \{+, -\}$
 - True Positive: $P(+ | \text{sick}) = 0.98$
 - False Negative: $P(- | \text{sick}) = 0.02$
 - False Positive: $P(+ | \text{not sick}) = 0.03$
 - True Negative: $P(- | \text{not sick}) = 0.97$
 - You get a positive result -> what is the probability that you are indeed sick?
 - $P(\text{sick} | +) = \frac{P(+|\text{sick})P(\text{sick})}{P(+|\text{sick})P(\text{sick}) + P(+|\text{not sick})P(\text{not sick})}$
 - $P(\text{sick} | +) = \frac{(0.98)(0.001)}{(0.98)(0.001) + (0.03)(0.999)} = 0.032$
- Why: because of the prior.
- So why should I ever believe a test?
 - Because $P(\text{ sick} | \text{ symptoms})$ is high.

Data

- Use students as example
- Data can be viewed a table
 - Rows are students (data points)
 - Columns are features
- The features are Random Variables
- Make a distribution

Instance	Name	Age	Major	GPA	...
1	XXX	XXX	XXX	XXX	
2	XXX	XXX	XXX	XXX	
3	XXX	XXX	XXX	XXX	
...	XXX	XXX	XXX	XXX	