

Online learning in repeated matrix games

Yoav Freund

February 24, 2006

Outline

Repeated Matrix Games

Outline

Repeated Matrix Games

The basic algorithm

Outline

Repeated Matrix Games

The basic algorithm

The basic analysis

Outline

Repeated Matrix Games

The basic algorithm

The basic analysis

Proof of minmax theorem

Zero sum games in matrix form

- ▶ Game between two players.

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ **Row** player chooses $i \in \{1, \dots, n\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ **Row** player chooses $i \in \{1, \dots, n\}$
- ▶ **Column** player chooses $j \in \{1, \dots, m\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$
- ▶ Game repeated many times.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**


Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .

Pure vs. mixed strategies


- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .
- ▶ **Column** player loses **$M(P, Q)$** .

Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r}b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$


Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

Q is a **column** vector. **P**^T is a row vector.

Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

Q is a **column** vector. **P^T** is a row vector.

$$\mathbf{M}(\mathbf{P}, \mathbf{Q}) = \mathbf{P}^T \mathbf{M} \mathbf{Q} = \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}(i) \mathbf{M}(i, j) \mathbf{Q}(j)$$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$
- ▶ $\eta > 0$ is the learning rate.

Main Theorem

- ▶ For **any** game matrix **M**.

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. $\mathbf{Q}_1, \dots, \mathbf{Q}_T$

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. **$\mathbf{Q}_1, \dots, \mathbf{Q}_T$**
- ▶ The sequence **$\mathbf{P}_1, \dots, \mathbf{P}_T$** produced by basic alg using **$\eta > 0$** satisfies

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_{\mathbf{P}} \left[\eta \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \text{RE}(\mathbf{P} \parallel \mathbf{P}_1) \right]$$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is

$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is

$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

- ▶ Where

$$\Delta_{T,n} = \sqrt{\frac{2 \ln n}{T}} + \frac{\ln n}{T} = O\left(\sqrt{\frac{\ln n}{T}}\right).$$

Main Lemma

On any iteration t

Main Lemma

On any iteration t

For any mixed strategy \tilde{P}

Main Lemma

On any iteration t

For any mixed strategy $\tilde{\mathbf{P}}$


$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$

Visual intuition

$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$

Visual intuition

$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$



Proof of Lemma (1)

$$\text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right)$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \end{aligned}$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\mathbf{P}_t(i)}{\mathbf{P}_{t+1}(i)} \end{aligned}$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\mathbf{P}_t(i)}{\mathbf{P}_{t+1}(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{Z_t}{e^{\eta \mathbf{M}(i, \mathbf{Q}_t)}} \end{aligned}$$

Proof of Lemma (2)

$$= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \end{aligned}$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \\ &= \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln (1 - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)) \end{aligned}$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \\ &= \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln (1 - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)) \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \end{aligned}$$

The minmax Theorem

John von Neumann, 1928.

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) \leq \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) \leq \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

In words: for **mixed** strategies, choosing second gives no advantage.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

$$= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\begin{aligned} \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} &\leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q} \\ &= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}} \\ &\leq \frac{1}{T} \sum_{t=1}^T \max_{\mathbf{Q}} \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \end{aligned}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t$$

by definition of \mathbf{Q}_t

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \overline{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \overline{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

but $\Delta_{T,n}$ can be set arbitrarily small.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.
- ▶ Is it realistic to assume that markets are at equilibrium?

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.
- ▶ Is it realistic to assume that markets are at equilibrium?
- ▶ If game is not zero sum (allows incentives to collaborate) and all players use learning then game converges to **correlated equilibrium**.