

Online learning in repeated matrix games

Yoav Freund

January 31, 2011

Outline

Repeated Games

Outline

Repeated Games

Fictitious play

Outline

Repeated Games

Fictitious play

Hannan Consistency

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Online strategies that converge to a correlated equilibrium

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Online strategies that converge to a correlated equilibrium

Repeated Zero-sum Games

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Online strategies that converge to a correlated equilibrium

Repeated Zero-sum Games

The basic algorithm

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Online strategies that converge to a correlated equilibrium

Repeated Zero-sum Games

The basic algorithm

The basic analysis

Outline

Repeated Games

Fictitious play

Hannan Consistency

Hannan set

Online strategies that converge to a correlated equilibrium

Repeated Zero-sum Games

The basic algorithm

The basic analysis

Proof of minmax theorem

Repeated games

- ▶ Game between two players.

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices ,**C**

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices, \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices, **C**
- ▶ **Row** player chooses $i \in \{1, \dots, n\}$
- ▶ **Column** player chooses $j \in \{1, \dots, m\}$
- ▶ Players Observe each other's action.

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices, \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Players Observe each other's action.
- ▶ Row player gains $(i, j) \in [0, 1]$

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Players Observe each other's action.
- ▶ Row player gains $(i, j) \in [0, 1]$
- ▶ Column player gains $\mathbf{C}(i, j) \in [0, 1]$

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Players Observe each other's action.
- ▶ Row player gains $(i, j) \in [0, 1]$
- ▶ Column player gains $\mathbf{C}(i, j) \in [0, 1]$
- ▶ Game repeated many times.

Repeated games

- ▶ Game between two players.
- ▶ Defined by two $n \times m$ matrices \mathbf{C}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Players Observe each other's action.
- ▶ Row player gains $(i, j) \in [0, 1]$
- ▶ Column player gains $\mathbf{C}(i, j) \in [0, 1]$
- ▶ Game repeated many times.
- ▶ Player choices can depend on the past.

Notions of Convergence

- ▶ 1 The players have diminishing regret (External or internal)

Notions of Convergence

- ▶ 1 The players have diminishing regret (External or internal)
- ▶ 2 The empirical distributions over the actions converge to a set of distributions over action k -tuples $\langle i_1, i_2, i_3, \dots, i_k \rangle$

Notions of Convergence

- ▶ 1 The players have diminishing regret (External or internal)
- ▶ 2 The empirical distributions over the actions converge to a set of distributions over action k -tuples $\langle i_1, i_2, i_3, \dots, i_k \rangle$
- ▶ The set of Nash Equilibria.

Notions of Convergence

- ▶ 1 The players have diminishing regret (External or internal)
- ▶ 2 The empirical distributions over the actions converge to a set of distributions over action k -tuples $\langle i_1, i_2, i_3, \dots, i_k \rangle$
- ▶ The set of Nash Equilibria.
- ▶ The set of Correlated equilibria.

Fictitious play

- **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.
- ▶ If **both** sides use fictitious play and the game is two-player zero sum game, then empirical distribution converges to the set of Nash Equilibria.

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.
- ▶ If **both** sides use fictitious play and the game is two-player zero sum game, then empirical distribution converges to the set of Nash Equilibria.
- ▶ If each player has only two actions, then convergence to Nash set also guaranteed.

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.
- ▶ If **both** sides use fictitious play and the game is two-player zero sum game, then empirical distribution converges to the set of Nash Equilibria.
- ▶ If each player has only two actions, then convergence to Nash set also guaranteed.
- ▶ Convergence to Nash set is not guaranteed in general.

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.
- ▶ If **both** sides use fictitious play and the game is two-player zero sum game, then empirical distribution converges to the set of Nash Equilibria.
- ▶ If each player has only two actions, then convergence to Nash set also guaranteed.
- ▶ Convergence to Nash set is not guaranteed in general.
- ▶ What if the other side does not follow fictitious play?

Fictitious play

- ▶ **Strategy:** Use best response to the empirical distribution of the actions of the opponent so far.
- ▶ Same as “follow the leader”.
- ▶ If **both** sides use fictitious play and the game is two-player zero sum game, then empirical distribution converges to the set of Nash Equilibria.
- ▶ If each player has only two actions, then convergence to Nash set also guaranteed.
- ▶ Convergence to Nash set is not guaranteed in general.
- ▶ What if the other side does not follow fictitious play?
- ▶ Conforming player can suffer non-diminishing regret.

Hannan's Algorithm

- ▶ *Approximation to Bayes risk in repeated play*,
Contributions to the theory of Games, 1957.

Hannan's Algorithm

- ▶ *Approximation to Bayes risk in repeated play*, Contributions to the theory of Games, 1957.
- ▶ 1957: IBM announces it will no longer be using vacuum tubes and releases its first computer that had 2000 transistors.

Hannan's Algorithm

- ▶ *Approximation to Bayes risk in repeated play*, Contributions to the theory of Games, 1957.
- ▶ 1957: IBM announces it will no longer be using vacuum tubes and releases its first computer that had 2000 transistors.
- ▶ Instead of using “follow the leader” use “follow the perturbed leader”, i.e. add a small amount of noise to the cumulative utility of each action, *then* pick the leader.

Hannan's Algorithm

- ▶ *Approximation to Bayes risk in repeated play*, Contributions to the theory of Games, 1957.
- ▶ 1957: IBM announces it will no longer be using vacuum tubes and releases its first computer that had 2000 transistors.
- ▶ Instead of using “follow the leader” use “follow the perturbed leader”, i.e. add a small amount of noise to the cumulative utility of each action, *then* pick the leader.
- ▶ **Hannan consistency:** Cumulative regret / Cumulative utility $\rightarrow 0$.

Hannan's Set

- ▶ Diminishing regret relative to playing a fixed pure strategy.

Hannan's Set

- ▶ Diminishing regret relative to playing a fixed pure strategy.
- ▶ Hannan's set contains all joint distributions over player's action where all players have no external regret.

Hannan's Set

- ▶ Diminishing regret relative to playing a fixed pure strategy.
- ▶ Hannan's set contains all joint distributions over player's action where all players have no external regret.
- ▶ Hannan's set contains the set of correlated equilibrium which contains the set of Nash Equilibria.

Reaching correlated equilibrium

- By all players minimizing internal regret.

Reaching correlated equilibrium

- ▶ By all players minimizing internal regret.
- ▶ By making a calibrated predictions of the opponent's next move and playing best response.

Zero sum games in matrix form

- ▶ Game between two players.

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ Row player chooses $i \in \{1, \dots, n\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$
- ▶ Game repeated many times.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**


Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .

Pure vs. mixed strategies


- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .
- ▶ **Column** player loses **$M(P, Q)$** .

Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$


Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

Q is a column vector. **P^T** is a row vector.

Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

Q is a column vector. **P^T** is a row vector.

$$\mathbf{M}(\mathbf{P}, \mathbf{Q}) = \mathbf{P}^T \mathbf{M} \mathbf{Q} = \sum_{i=1}^n \sum_{j=1}^m \mathbf{P}(i) \mathbf{M}(i, j) \mathbf{Q}(j)$$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$
- ▶ $\eta > 0$ is the learning rate.

Main Theorem

- ▶ For **any** game matrix **M**.

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. $\mathbf{Q}_1, \dots, \mathbf{Q}_T$

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. **$\mathbf{Q}_1, \dots, \mathbf{Q}_T$**
- ▶ The sequence **$\mathbf{P}_1, \dots, \mathbf{P}_T$** produced by basic alg using **$\eta > 0$** satisfies

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_{\mathbf{P}} \left[\eta \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \text{RE}(\mathbf{P} \parallel \mathbf{P}_1) \right]$$

Corollary

- Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is

$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is

$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

- ▶ Where

$$\Delta_{T,n} = \sqrt{\frac{2 \ln n}{T}} + \frac{\ln n}{T} = O\left(\sqrt{\frac{\ln n}{T}}\right).$$

Main Lemma

On any iteration t

Main Lemma

On any iteration t

For any mixed strategy $\tilde{\mathbf{P}}$

Main Lemma

On any iteration t

For any mixed strategy $\tilde{\mathbf{P}}$


$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$

Visual intuition

$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$

Visual intuition

$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)$$



Proof of Lemma (1)

$$\text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right)$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \end{aligned}$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\mathbf{P}_t(i)}{\mathbf{P}_{t+1}(i)} \end{aligned}$$

Proof of Lemma (1)

$$\begin{aligned} & \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1} \right) - \text{RE} \left(\tilde{\mathbf{P}} \parallel \mathbf{P}_t \right) \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_{t+1}(i)} - \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\tilde{\mathbf{P}}(i)}{\mathbf{P}_t(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{\mathbf{P}_t(i)}{\mathbf{P}_{t+1}(i)} \\ &= \sum_{i=1}^n \tilde{\mathbf{P}}(i) \ln \frac{Z_t}{e^{\eta \mathbf{M}(i, \mathbf{Q}_t)}} \end{aligned}$$

Proof of Lemma (2)

$$= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \end{aligned}$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \\ &= \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln (1 - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)) \end{aligned}$$

Proof of Lemma (2)

$$\begin{aligned} &= \eta \sum_{i=1}^n \tilde{\mathbf{P}}(i) \mathbf{M}(i, \mathbf{Q}_t) + \ln Z_t \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln \left[\sum_{i=1}^n \mathbf{P}_t(i) (1 - (1 - e^{-\eta}) \mathbf{M}(i, \mathbf{Q}_t)) \right] \\ &= \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + \ln (1 - (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)) \\ &\leq \eta \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}_t) + (1 - e^{-\eta}) \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \end{aligned}$$

The minmax Theorem

John von Neumann, 1928.

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) \leq \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) \leq \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

In words: for **mixed** strategies, choosing second gives no advantage.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Let $\bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t$ and $\bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Let $\bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t$ and $\bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

$$= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Let $\bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t$ and $\bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$

$$\begin{aligned} \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} &\leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q} \\ &= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}} \\ &\leq \frac{1}{T} \sum_{t=1}^T \max_{\mathbf{Q}} \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \end{aligned}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t$$

by definition of \mathbf{Q}_t

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

but $\Delta_{T,n}$ can be set arbitrarily small.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.
- ▶ Is it realistic to assume that markets are at equilibrium?

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.
- ▶ Is it realistic to assume that markets are at equilibrium?
- ▶ If game is not zero sum (allows incentives to collaborate) and all players use learning then game converges to **correlated equilibrium**.