

# Combining infinite sets of experts

Yoav Freund

January 24, 2006

# Outline

## Review

# Outline

Review

The Universal prediction machine

# Outline

Review

The Universal prediction machine

The biased coins set of experts

# Outline

Review

The Universal prediction machine

The biased coins set of experts

Bayes using Jeffrey's prior

- Laplace Approximation

- Choosing the optimal prior

- Kritchevski Trofimov Prediction Rule

- Laplace Rule of Succession

# Outline

Review

The Universal prediction machine

The biased coins set of experts

Bayes using Jeffrey's prior

- Laplace Approximation

- Choosing the optimal prior

- Kritchevski Trofimov Prediction Rule

- Laplace Rule of Succession

Shtarkov lower bound for finite horizon

# Outline

## Review

The Universal prediction machine

The biased coins set of experts

Bayes using Jeffrey's prior

- Laplace Approximation

- Choosing the optimal prior

- Kritchevski Trofimov Prediction Rule

- Laplace Rule of Succession

Shtarkov lower bound for finite horizon

Generalization to larger sets of distributions

# The online Bayes Algorithm

- Total loss of expert  $i$

$$L_i^t = - \sum_{s=1}^t \log p_i^s(c^s); \quad L_i^0 = 0$$



# The online Bayes Algorithm

- Total loss of expert  $i$

$$L_i^t = - \sum_{s=1}^t \log p_i^s(c^s); \quad L_i^0 = 0$$

- Weight of expert  $i$

$$w_i^t = w_i^1 e^{-L_i^{t-1}} = w_i^1 \prod_{s=1}^{t-1} p_i^s(c^s)$$

# The online Bayes Algorithm

- ▶ Total loss of expert  $i$

$$L_i^t = - \sum_{s=1}^t \log p_i^s(c^s); \quad L_i^0 = 0$$

- ▶ Weight of expert  $i$

$$w_i^t = w_i^1 e^{-L_i^{t-1}} = w_i^1 \prod_{s=1}^{t-1} p_i^s(c^s)$$

- ▶ Freedom to choose initial weights.

$$w_i^1 \geq 0, \sum_{i=1}^n w_i^1 = 1$$

# The online Bayes Algorithm

- Total loss of expert  $i$

$$L_i^t = - \sum_{s=1}^t \log p_i^s(c^s); \quad L_i^0 = 0$$

- Weight of expert  $i$

$$w_i^t = w_i^1 e^{-L_i^{t-1}} = w_i^1 \prod_{s=1}^{t-1} p_i^s(c^s)$$

- Freedom to choose initial weights.

$$w_i^1 \geq 0, \sum_{i=1}^N w_i^1 = 1$$

- Prediction of algorithm  $A$

$$\mathbf{p}_A^t = \frac{\sum_{i=1}^N w_i^t \mathbf{p}_i^t}{\sum_{i=1}^N w_i^t}$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t}$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t}$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t)$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\begin{aligned}\frac{W^{t+1}}{W^t} &= \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t) \\ -\log \frac{W^{t+1}}{W^t} &= -\log p_A^t(c^t)\end{aligned}$$



## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t)$$

$$-\log \frac{W^{t+1}}{W^t} = -\log p_A^t(c^t)$$

$$-\log \frac{W^{T+1}}{W^1} = -\sum_{t=1}^T \log p_A^t(c^t)$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t)$$

$$-\log \frac{W^{t+1}}{W^t} = -\log p_A^t(c^t)$$

$$-\log \frac{W^{T+1}}{W^1} = -\sum_{t=1}^T \log p_A^t(c^t) = L_A^T$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t)$$

$$-\log \frac{W^{t+1}}{W^t} = -\log p_A^t(c^t)$$

$$-\log W^{T+1} = -\log \frac{W^{T+1}}{W^1} = -\sum_{t=1}^T \log p_A^t(c^t) = L_A^T$$

## Cumulative loss vs. Final total weight

Total weight:  $W^t \doteq \sum_{i=1}^N w_i^t$

$$\frac{W^{t+1}}{W^t} = \frac{\sum_{i=1}^N w_i^t e^{\log p_i^t(c^t)}}{\sum_{i=1}^N w_i^t} = \frac{\sum_{i=1}^N w_i^t p_i^t(c^t)}{\sum_{i=1}^N w_i^t} = p_A^t(c^t)$$

$$-\log \frac{W^{t+1}}{W^t} = -\log p_A^t(c^t)$$

$$-\log W^{T+1} = -\log \frac{W^{T+1}}{W^1} = -\sum_{t=1}^T \log p_A^t(c^t) = L_A^T$$

**EQUALITY** not bound!

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$L_A^T = -\log W^{T+1}$$

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$L_A^T = -\log W^{T+1}$$

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$L_A^T = -\log W^{T+1} = -\log \sum_{i=1}^N w_i^{T+1}$$



## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$\begin{aligned} L_A^T &= -\log W^{T+1} = -\log \sum_{i=1}^N w_i^{T+1} \\ &= -\log \sum_{i=1}^N w_i^1 e^{-L_i^T} \end{aligned}$$

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$\begin{aligned} L_A^T &= -\log W^{T+1} = -\log \sum_{i=1}^N w_i^{T+1} \\ &= -\log \sum_{i=1}^N w_i^1 e^{-L_i^T} \leq -\log \max_i \left( w_i^1 e^{-L_i^T} \right) \end{aligned}$$

## Simple Bound

- ▶ Use non-uniform initial weights  $\sum_i w_i^1 = 1$
- ▶ Total Weight is at least the weight of the best expert.

$$\begin{aligned} L_A^T &= -\log W^{T+1} = -\log \sum_{i=1}^N w_i^{T+1} \\ &= -\log \sum_{i=1}^N w_i^1 e^{-L_i^T} \leq -\log \max_i \left( w_i^1 e^{-L_i^T} \right) \\ &= \min_i \left( L_i^T - \log w_i^1 \right) \end{aligned}$$

## Standardizing online prediction algorithms

- Fix a universal Turing machine  $U$ .

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .



## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .
  - ▶ To ensure  $p$  has a finite description. Restrict to rational numbers  $n/m$

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .
  - ▶ To ensure  $p$  has a finite description. Restrict to rational numbers  $n/m$
- ▶ Any online prediction algorithm can be represented as code  $\vec{b}(E)$  for  $U$ . The code length is  $|\vec{b}(E)|$ .

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .
  - ▶ To ensure  $p$  has a finite description. Restrict to rational numbers  $n/m$
- ▶ Any online prediction algorithm can be represented as code  $\vec{b}(E)$  for  $U$ . The code length is  $|\vec{b}(E)|$ .
- ▶ Most sequences do not correspond to valid prediction algorithms.

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .
  - ▶ To ensure  $p$  has a finite description. Restrict to rational numbers  $n/m$
- ▶ Any online prediction algorithm can be represented as code  $\vec{b}(E)$  for  $U$ . The code length is  $|\vec{b}(E)|$ .
- ▶ Most sequences do not correspond to valid prediction algorithms.
- ▶  $V(\vec{b}, \vec{X}, t) = 1$  if the program  $\vec{b}$ , given  $\vec{X}$  as input, halts within  $t$  steps and outputs a well-formed prediction. Otherwise  $V(\vec{b}, \vec{X}, t) = 0$

## Standardizing online prediction algorithms

- ▶ Fix a universal Turing machine  $U$ .
- ▶ An online prediction algorithm  $E$  is a program that
  - ▶ given as input The past  $\vec{X} \in \{0, 1\}^t$
  - ▶ runs finite time and outputs
  - ▶ A prediction for the next bit  $p(\vec{X}) \in [0, 1]$ .
  - ▶ To ensure  $p$  has a finite description. Restrict to rational numbers  $n/m$
- ▶ Any online prediction algorithm can be represented as code  $\vec{b}(E)$  for  $U$ . The code length is  $|\vec{b}(E)|$ .
- ▶ Most sequences do not correspond to valid prediction algorithms.
- ▶  $V(\vec{b}, \vec{X}, t) = 1$  if the program  $\vec{b}$ , given  $\vec{X}$  as input, halts within  $t$  steps and outputs a well-formed prediction. Otherwise  $V(\vec{b}, \vec{X}, t) = 0$
- ▶  $V(\vec{b}, \vec{X}, t)$  is computable (recursively enumerable).

# A universal prediction machine

- ▶ Assign to the code  $\vec{b}$  the initial weight  $w_{\vec{b}}^1 = 2^{-|\vec{b}| - \log_2 |\vec{b}|}$ .

## A universal prediction machine

- ▶ Assign to the code  $\vec{b}$  the initial weight  $w_{\vec{b}}^1 = 2^{-|\vec{b}| - \log_2 |\vec{b}|}$ .
- ▶ The total initial weight over all finite binary sequences is one.

# A universal prediction machine

- ▶ Assign to the code  $\vec{b}$  the initial weight  $w_{\vec{b}}^1 = 2^{-|\vec{b}| - \log_2 |\vec{b}|}$ .
- ▶ The total initial weight over all finite binary sequences is one.
- ▶ Run the Bayes algorithm over “all” prediction algorithms.



# A universal prediction machine

- ▶ Assign to the code  $\vec{b}$  the initial weight  $w_{\vec{b}}^1 = 2^{-|\vec{b}| - \log_2 |\vec{b}|}$ .
- ▶ The total initial weight over all finite binary sequences is one.
- ▶ Run the Bayes algorithm over “all” prediction algorithms.
- ▶ **technical details:** On iteration  $t$ ,  $|\vec{X}| = t$ . Use the predictions of programs  $\vec{b}$  such that  $|\vec{b}| \leq t$  and for which  $V(\vec{b}, \vec{X}, 2^t) = 1$ . Assign the remaining mass the prediction  $1/2$  (insuring a loss of 1)

## Performance of the universal prediction algorithm

- Using  $L_A \leq \min_i (L_i - \log w_i^1)$

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$
- ▶ When  $t \leq |\vec{b}(E)|$  the algorithm is not used and thus its loss is 1

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$
- ▶ When  $t \leq |\vec{b}(E)|$  the algorithm is not used and thus its loss is 1
- ▶ We get that the loss of the Universal algorithm is at most  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)| + L_E$

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$
- ▶ When  $t \leq |\vec{b}(E)|$  the algorithm is not used and thus its loss is 1
- ▶ We get that the loss of the Universal algorithm is at most  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)| + L_E$
- ▶ More careful analysis can reduce  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)|$  to  $|\vec{b}(E)|$

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$
- ▶ When  $t \leq |\vec{b}(E)|$  the algorithm is not used and thus its loss is 1
- ▶ We get that the loss of the Universal algorithm is at most  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)| + L_E$
- ▶ More careful analysis can reduce  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)|$  to  $|\vec{b}(E)|$
- ▶ Code length is arbitrarily close to the Kolmogorov Complexity of the sequence.

## Performance of the universal prediction algorithm

- ▶ Using  $L_A \leq \min_i (L_i - \log w_i^1)$
- ▶ Assume  $E$  is a prediction algorithm which generates the  $t$ th prediction in time smaller than  $2^t$
- ▶ When  $t \leq |\vec{b}(E)|$  the algorithm is not used and thus its loss is 1
- ▶ We get that the loss of the Universal algorithm is at most  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)| + L_E$
- ▶ More careful analysis can reduce  $2|\vec{b}(E)| + \log_2 |\vec{b}(E)|$  to  $|\vec{b}(E)|$
- ▶ Code length is arbitrarily close to the Kolmogorov Complexity of the sequence.
- ▶ Ridiculously bad running time.



## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression

## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression
- ▶ Suppose we have  $K$  copies of each expert.

## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression
- ▶ Suppose we have  $K$  copies of each expert.
- ▶ Two part code has to point to one of the  $KN$  experts

$$L_A \leq \log NK + \min_i L_i^T = \log NK + \min_i L_i^T$$

## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression
- ▶ Suppose we have  $K$  copies of each expert.
- ▶ Two part code has to point to one of the  $KN$  experts  
 $L_A \leq \log NK + \min_i L_i^T = \log NK + \min_i L_i^T$
- ▶ If we use Bayes predictor + arithmetic coding we get:

$$L_A = -\log W^{T+1} \leq \log K \max_i \frac{1}{NK} e^{-L_i^T} = \log N + \min_i L_i^T$$

## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression
- ▶ Suppose we have  $K$  copies of each expert.
- ▶ Two part code has to point to one of the  $KN$  experts  
 $L_A \leq \log NK + \min_i L_i^T = \log NK + \min_i L_i^T$
- ▶ If we use Bayes predictor + arithmetic coding we get:

$$L_A = -\log W^{T+1} \leq \log K \max_i \frac{1}{NK} e^{-L_i^T} = \log N + \min_i L_i^T$$

- ▶ We don't pay a penalty for copies.

## Bayes coding is better than two part codes

- ▶ Simple bound as good as bound for two part codes (MDL) but enables online compression
- ▶ Suppose we have  $K$  copies of each expert.
- ▶ Two part code has to point to one of the  $KN$  experts  
 $L_A \leq \log NK + \min_i L_i^T = \log NK + \min_i L_i^T$
- ▶ If we use Bayes predictor + arithmetic coding we get:

$$L_A = -\log W^{T+1} \leq \log K \max_i \frac{1}{NK} e^{-L_i^T} = \log N + \min_i L_i^T$$

- ▶ We don't pay a penalty for copies.
- ▶ More generally, the regret is smaller if many of the experts perform well.

## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .

## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .
- ▶ Set of experts is **uncountably infinite**.



## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .
- ▶ Set of experts is **uncountably infinite**.
- ▶ Only countably many experts can be assigned non-zero weight.

## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .
- ▶ Set of experts is **uncountably infinite**.
- ▶ Only countably many experts can be assigned non-zero weight.
- ▶ Instead, we assign the experts a **Density Measure**.

## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .
- ▶ Set of experts is **uncountably infinite**.
- ▶ Only countably many experts can be assigned non-zero weight.
- ▶ Instead, we assign the experts a **Density Measure**.
- ▶  $L_A \leq \min_i (L_i - \log w_i^1)$  is meaningless.

## The biased coins set of experts

- ▶ Each expert corresponds to a biased coin, predicts with a fixed  $\theta \in [0, 1]$ .
- ▶ Set of experts is **uncountably infinite**.
- ▶ Only countably many experts can be assigned non-zero weight.
- ▶ Instead, we assign the experts a **Density Measure**.
- ▶  $L_A \leq \min_i (L_i - \log w_i^1)$  is meaningless.
- ▶ Can we still get a meaningful bound?

## Bayes Algorithm for biased coins

- Replace the initial weight by a density measure

$$w(\theta) = w^1(\theta), \int_0^1 w(\theta) d\theta = 1$$

## Bayes Algorithm for biased coins

- ▶ Replace the initial weight by a density measure  
 $w(\theta) = w^1(\theta), \int_0^1 w(\theta) d\theta = 1$
- ▶ Relationship between final total weight and total log loss remains unchanged:

$$L_A = \ln \int_0^1 w(\theta) e^{-L_\theta^{T+1}} d\theta$$

## Bayes Algorithm for biased coins

- ▶ Replace the initial weight by a density measure  
 $w(\theta) = w^1(\theta), \int_0^1 w(\theta) d\theta = 1$
- ▶ Relationship between final total weight and total log loss remains unchanged:

$$L_A = \ln \int_0^1 w(\theta) e^{-L_\theta^{T+1}} d\theta$$

- ▶ We need a new **lower bound** on the final total weight


## Main Idea

If  $w^t(\theta)$  is large then  $w^t(\theta + \epsilon)$  is also large.



## Main Idea

If  $w^t(\theta)$  is large then  $w^t(\theta + \epsilon)$  is also large.



## Expanding the exponent around the peak

- For log loss the best  $\theta$  is empirical distribution of the seq.

$$\hat{\theta} = \frac{\#\{x^t = 1; 1 \leq t \leq T\}}{T}$$

## Expanding the exponent around the peak

- For log loss the best  $\theta$  is empirical distribution of the seq.

$$\hat{\theta} = \frac{\#\{x^t = 1; 1 \leq t \leq T\}}{T}$$

- The total loss scales with  $T$

$$L_{\theta} = T \cdot (\hat{\theta} \ell(\theta, 1) + (1 - \hat{\theta}) \ell(\theta, 0)) \doteq T \cdot g(\hat{\theta}, \theta)$$

## Expanding the exponent around the peak

- For log loss the best  $\theta$  is empirical distribution of the seq.

$$\hat{\theta} = \frac{\#\{x^t = 1; 1 \leq t \leq T\}}{T}$$

- The total loss scales with  $T$

$$L_{\theta} = T \cdot (\hat{\theta} \ell(\theta, 1) + (1 - \hat{\theta}) \ell(\theta, 0)) \doteq T \cdot g(\hat{\theta}, \theta)$$

## Expanding the exponent around the peak

- For log loss the best  $\theta$  is empirical distribution of the seq.

$$\hat{\theta} = \frac{\#\{x^t = 1; 1 \leq t \leq T\}}{T}$$

- The total loss scales with  $T$

$$L_{\theta} = T \cdot (\hat{\theta} \ell(\theta, 1) + (1 - \hat{\theta}) \ell(\theta, 0)) \doteq T \cdot g(\hat{\theta}, \theta)$$

$$L_A - L_{\min} \leq \ln \int_0^1 w(\theta) e^{-L_{\theta}} d\theta - \ln e^{L_{\min}}$$

## Expanding the exponent around the peak

- For log loss the best  $\theta$  is empirical distribution of the seq.

$$\hat{\theta} = \frac{\#\{x^t = 1; 1 \leq t \leq T\}}{T}$$

- The total loss scales with  $T$

$$L_{\theta} = T \cdot (\hat{\theta} \ell(\theta, 1) + (1 - \hat{\theta}) \ell(\theta, 0)) \doteq T \cdot g(\hat{\theta}, \theta)$$

$$L_A - L_{\min} \leq \ln \int_0^1 w(\theta) e^{-L_{\theta}} d\theta - \ln e^{L_{\min}}$$

$$= \ln \int_0^1 w(\theta) e^{-(L_{\theta} - L_{\min})} d\theta$$

$$\text{pause} = \ln \int_0^1 w(\theta) e^{T(g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))} d\theta$$

## Laplace approximation (idea)

- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .

## Laplace approximation (idea)

- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .
- ▶ First and second terms in the expansion are zero.



## Laplace approximation (idea)

- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .
- ▶ First and second terms in the expansion are zero.
- ▶ Third term gives a quadratic expression in the exponent

## Laplace approximation (idea)


- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .
- ▶ First and second terms in the expansion are zero.
- ▶ Third term gives a quadratic expression in the exponent
- ▶  $\Rightarrow$  a gaussian approximation of the posterior.

## Laplace approximation (idea)

- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .
- ▶ First and second terms in the expansion are zero.
- ▶ Third term gives a quadratic expression in the exponent
- ▶  $\Rightarrow$  a gaussian approximation of the posterior.

## Laplace approximation (idea)

- ▶ Taylor expansion of  $g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta})$  around  $\theta = \hat{\theta}$ .
- ▶ First and second terms in the expansion are zero.
- ▶ Third term gives a quadratic expression in the exponent
- ▶  $\Rightarrow$  a gaussian approximation of the posterior.



## Laplace Approximation (details)

$$\int_0^1 w(\theta) e^{T(g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))} d\theta$$

# Laplace Approximation (details)

$$\begin{aligned} & \int_0^1 w(\theta) e^{T(g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))} d\theta \\ &= w(\hat{\theta}) \sqrt{\frac{-2\pi}{T \left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} (g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))}} + O(T^{-3/2}) \end{aligned}$$

- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## Choosing the optimal prior

- Choose  $w(\theta)$  to maximize the worst-case final total weight

$$\min_{\hat{\theta}} w(\hat{\theta}) \sqrt{\frac{-2\pi}{T \left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}}} (g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))}$$

## Choosing the optimal prior

- Choose  $w(\theta)$  to maximize the worst-case final total weight

$$\min_{\hat{\theta}} w(\hat{\theta}) \sqrt{\frac{-2\pi}{T \left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} (g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))}}$$

- Make bound equal for all  $\hat{\theta} \in [0, 1]$  by choosing

$$w^*(\hat{\theta}) = \frac{1}{Z} \sqrt{\frac{\left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} (g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))}{-2\pi}},$$

where  $Z$  is the normalization factor:

$$Z = \sqrt{\frac{1}{2\pi}} \int_0^1 \sqrt{\left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} (g(\hat{\theta}, \hat{\theta}) - g(\hat{\theta}, \theta))} d\hat{\theta}$$



- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## The bound for the optimal prior

- Plugging in we get

$$\begin{aligned} L_A - L_{\min} &\leq \ln \int_0^1 w^*(\theta) e^{T(g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}))} d\theta \\ &= \ln \left( \sqrt{\frac{2\pi Z}{T}} + O(T^{-3/2}) \right) \\ &= \frac{1}{2} \ln \frac{T}{2\pi} - \frac{1}{2} \ln Z + O(1/T) . \end{aligned}$$

- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## Solving for log-loss

- The exponent in the integral is

$$g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}) = \hat{\theta} \ln \frac{\hat{\theta}}{\theta} + (1 - \hat{\theta}) \ln \frac{1 - \hat{\theta}}{1 - \theta} = D_{KL}(\hat{\theta} || \theta)$$

- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## Solving for log-loss

- ▶ The exponent in the integral is

$$g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}) = \hat{\theta} \ln \frac{\hat{\theta}}{\theta} + (1 - \hat{\theta}) \ln \frac{1 - \hat{\theta}}{1 - \theta} = D_{KL}(\hat{\theta} || \theta)$$

- ▶ The second derivative

$$\left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} D_{KL}(\hat{\theta} || \theta) = \hat{\theta}(1 - \hat{\theta})$$

Is called the **empirical Fisher information**

- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## Solving for log-loss

- ▶ The exponent in the integral is

$$g(\hat{\theta}, \theta) - g(\hat{\theta}, \hat{\theta}) = \hat{\theta} \ln \frac{\hat{\theta}}{\theta} + (1 - \hat{\theta}) \ln \frac{1 - \hat{\theta}}{1 - \theta} = D_{KL}(\hat{\theta} || \theta)$$

- ▶ The second derivative

$$\left. \frac{d^2}{d\theta^2} \right|_{\theta=\hat{\theta}} D_{KL}(\hat{\theta} || \theta) = \hat{\theta}(1 - \hat{\theta})$$

Is called the **empirical Fisher information**

- ▶ The optimal prior:

$$w^*(\hat{\theta}) = \frac{1}{\pi \sqrt{\hat{\theta}(1 - \hat{\theta})}}$$

Known in general as **Jeffrey's prior**. And, in this case, the **Dirichlet-(1/2, 1/2) prior**.

- └ Bayes using Jeffrey's prior
- └ Choosing the optimal prior

## The cumulative log loss of Bayes using Jeffrey's prior



$$L_A - L_{\min} \leq \frac{1}{2} \ln(T + 1) + \frac{1}{2} \ln \frac{\pi}{2} + O(1/T)$$

## But what is the prediction rule?

- ▶ As luck would have it the Dirichlet prior is the **conjugate prior** for the Binomial distribution.

## But what is the prediction rule?

- ▶ As luck would have it the Dirichlet prior is the **conjugate prior** for the Binomial distribution.
- ▶ Observed  $t$  bits,  $n$  of which were  $1$ . The posterior is:

$$\frac{1}{Z \sqrt{\theta(1-\theta)}} \theta^n (1-\theta)^{t-n} = \frac{1}{Z} \theta^{n-1/2} (1-\theta)^{t-n-1/2}$$

## But what is the prediction rule?

- ▶ As luck would have it the Dirichlet prior is the **conjugate prior** for the Binomial distribution.
- ▶ Observed  $t$  bits,  $n$  of which were  $1$ . The posterior is:

$$\frac{1}{Z \sqrt{\theta(1-\theta)}} \theta^n (1-\theta)^{t-n} = \frac{1}{Z} \theta^{n-1/2} (1-\theta)^{t-n-1/2}$$

- ▶ The posterior average is:

$$\frac{\int_0^1 \theta^{n+1/2} (1-\theta)^{t-n-1/2} d\theta}{\int_0^1 \theta^{n-1/2} (1-\theta)^{t-n-1/2} d\theta} = \frac{n+1/2}{t+1}$$



## But what is the prediction rule?

- ▶ As luck would have it the Dirichlet prior is the **conjugate prior** for the Binomial distribution.
- ▶ Observed  $t$  bits,  $n$  of which were  $1$ . The posterior is:

$$\frac{1}{Z \sqrt{\theta(1-\theta)}} \theta^n (1-\theta)^{t-n} = \frac{1}{Z} \theta^{n-1/2} (1-\theta)^{t-n-1/2}$$

- ▶ The posterior average is:

$$\frac{\int_0^1 \theta^{n+1/2} (1-\theta)^{t-n-1/2} d\theta}{\int_0^1 \theta^{n-1/2} (1-\theta)^{t-n-1/2} d\theta} = \frac{n+1/2}{t+1}$$

- ▶ This is called the Trichevsky Trofimov prediction rule.

- └ Bayes using Jeffrey's prior
- └ Laplace Rule of Succession

## Laplace Rule of Succession

- ▶ Laplace suggested using the uniform prior, which is also a conjugate prior.

## Laplace Rule of Succession

- ▶ Laplace suggested using the uniform prior, which is also a conjugate prior.
- ▶ In this case the posterior average is:

$$\frac{\int_0^1 \theta^{n+1} (1 - \theta)^{t-n} d\theta}{\int_0^1 \theta^n (1 - \theta)^{t-n} d\theta} = \frac{n+1}{t+2}$$

## Laplace Rule of Succession

- ▶ Laplace suggested using the uniform prior, which is also a conjugate prior.
- ▶ In this case the posterior average is:

$$\frac{\int_0^1 \theta^{n+1} (1 - \theta)^{t-n} d\theta}{\int_0^1 \theta^n (1 - \theta)^{t-n} d\theta} = \frac{n+1}{t+2}$$

- ▶ The bound on the cumulative log loss is worse:

$$L_A - L_{\min} = \ln T + O(1)$$

## Laplace Rule of Succession

- ▶ Laplace suggested using the uniform prior, which is also a conjugate prior.
- ▶ In this case the posterior average is:

$$\frac{\int_0^1 \theta^{n+1} (1 - \theta)^{t-n} d\theta}{\int_0^1 \theta^n (1 - \theta)^{t-n} d\theta} = \frac{n+1}{t+2}$$

- ▶ The bound on the cumulative log loss is worse:

$$L_A - L_{\min} = \ln T + O(1)$$

- ▶ Suffers larger regret when  $\hat{\theta}$  is far from  $1/2$

## Shtarkov Lower bound

- What is the **optimal** prediction when  $T$  is known in advance?

## Shtarkov Lower bound

- ▶ What is the **optimal** prediction when  $T$  is known in advance?



$$L_*^T - \min_{\theta} L_{\theta}^T \geq \frac{1}{2} \ln(T+1) + \frac{1}{2} \ln \frac{\pi}{2} - O\left(\frac{1}{\sqrt{T}}\right)$$

## Multinomial Distributions

- For a distribution over  $k$  elements (Multinomial) [Xie and Barron]



## Multinomial Distributions

- ▶ For a distribution over  $k$  elements (Multinomial) [Xie and Barron]
- ▶ Use the add 1/2 rule (KT).

$$p(i) = \frac{n_i + 1/2}{t + k/2}$$

## Multinomial Distributions

- ▶ For a distribution over  $k$  elements (Multinomial) [Xie and Barron]
- ▶ Use the add 1/2 rule (KT).

$$p(i) = \frac{n_i + 1/2}{t + k/2}$$

- ▶ Bound is

$$L_A - L_{\min} \leq \frac{k-1}{2} \ln T + C + o(1)$$

## Multinomial Distributions

- ▶ For a distribution over  $k$  elements (Multinomial) [Xie and Barron]
- ▶ Use the add 1/2 rule (KT).

$$p(i) = \frac{n_i + 1/2}{t + k/2}$$

- ▶ Bound is

$$L_A - L_{\min} \leq \frac{k-1}{2} \ln T + C + o(1)$$

- ▶ The constant  $C$  is optimal.

## Exponential Distributions

- ▶ For any set of distributions from the exponential family defined by  $k$  parameters (Some technical conditions on closure of set??) [Rissanen??]

## Exponential Distributions

- ▶ For any set of distributions from the exponential family defined by  $k$  parameters (Some technical conditions on closure of set??) [Rissanen??]
- ▶ Use Bayes Algorithm with Jeffrey's prior:

$$w^*(\hat{\theta}) = \frac{1}{Z} \frac{1}{\sqrt{|\mathbf{H}(D_{KL}(\hat{\theta}||\theta))|_{\theta=\hat{\theta}}|}}$$

$\mathbf{H}$  denotes the Hessian.

## Exponential Distributions

- ▶ For any set of distributions from the exponential family defined by  $k$  parameters (Some technical conditions on closure of set??) [Rissanen??]
- ▶ Use Bayes Algorithm with Jeffrey's prior:

$$w^*(\hat{\theta}) = \frac{1}{Z} \frac{1}{\sqrt{|\mathbf{H}(D_{KL}(\hat{\theta}||\theta))|_{\theta=\hat{\theta}}|}}$$

$\mathbf{H}$  denotes the Hessian.



$$L_A - L_{\min} \leq \frac{k-1}{2} \ln T - \ln Z + o(1)$$

## General Distributions

- ▶ Characterize distribution family by metric entropy.

## General Distributions

- ▶ Characterize distribution family by metric entropy.
- ▶ Fixed parameter set usually corresponds to polynomial metrix entropy

$$N(1/\epsilon) = O\left(\frac{1}{\epsilon^d}\right)$$

$d$  is the number of parameters.



## General Distributions

- ▶ Characterize distribution family by metric entropy.
- ▶ Fixed parameter set usually corresponds to polynomial metrix entropy

$$N(1/\epsilon) = O\left(\frac{1}{\epsilon^d}\right)$$

$d$  is the number of parameters.

- ▶ [Haussler and Opper] show that the coefficient in front of  $\ln T$  is optimal for distribution families where the metric entropy is up to

$$N(1/\epsilon) = O\left(e^{\epsilon^{-\alpha}}\right)$$

For all  $\alpha \leq 5/2$ .

## next Class

- ▶ Variable-length markov models - a set of distributions with increasing number of parameters.

## next Class

- ▶ Variable-length markov models - a set of distributions with increasing number of parameters.
- ▶ The context algorithm: An efficient implementation of the Bayes algorithm which achieves close-to-optimal worst case bounds.