Masked observations

1 Defining equations

Lemma 1.1. *For every natural number Q and natural number t,* $\mathrm{centergate}(Q, t) = \mathrm{not}(\mathrm{center}(t))\,\mathrm{and}\,\mathrm{rightdefect}(Q, t, 1)$.

Proof. The goal follows from $\mathrm{centergatedefinition}(Q, t)$. □

Lemma 1.2. *For every natural number Q and natural number t,* $\mathrm{alignedbackground}(Q, t) = \mathrm{wittshl}(\mathrm{wittrow}(t), Q)$.

Proof. The goal follows from $\mathrm{backgrounddefinition}(Q, t)$. □

Lemma 1.3. *For every natural number Q and natural number t,* $\mathrm{aligneddefect}(Q, t) = \mathrm{wittxor}(\mathrm{wittrow}(t + Q), \mathrm{alignedbackground}(Q, t))$.

Proof. The goal follows from $\mathrm{aligneddefectdefinition}(Q, t)$. □

Lemma 1.4. *For every natural number L and natural number H,* $\mathrm{carryledger}(L, H, 0) = 0$.

Proof. The goal follows from $\mathrm{carryledgerdefinition}(L, H)$. □

Lemma 1.5. *For every natural number L and natural number H and natural number t,* $\mathrm{carryledger}(L, H, \mathrm{succ}(t)) = \mathrm{wittxor}(\mathrm{carryledger}(L, H, t), \mathrm{wittxor}(\mathrm{wittcarry}(\mathrm{page}(L, H, t)), \mathrm{na}$

Proof. The goal follows from $\mathrm{carryledgerdefinition2}(L, H, t)$. □

Lemma 1.6. *For every natural number n,* $\mathrm{centerflipmask}(n) = \mathrm{wittshl}(7, n)$.

Proof. The goal follows from $\mathrm{centerflipmaskdefinition}(n)$. □

Lemma 1.7. *For every natural number $start$,* $\mathrm{centersign}(start, 0) = 1$.

Proof. The goal follows from $\mathrm{centersigndefinition}(start)$. □

Lemma 1.8. *For every natural number $start$ and natural number t,* $\mathrm{centersign}(start, \mathrm{succ}(t)) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(start))) \cdot \mathrm{centersign}(start + 1, t)$.

Proof. The goal follows from $\mathrm{centersigndefinition2}(start, t)$. □

Lemma 1.9. *For every natural number b and natural number d,* $\mathrm{defectstep}(b, d) = \mathrm{wittxor}(\mathrm{wittphi}(d), \mathrm{wittshl}(\mathrm{crosscarry}(b, d), 1))$.

Proof. The goal follows from $\mathrm{defectstepdefinition}(b, d)$. □

Lemma 1.10. *For every natural number Q and natural number d and natural number t,* diagonalphase(Q, d, t) = bitvalue(ldiag$(d, t{+}Q)$)$-$bitvalue(ldiag(d, t)).

Proof. The goal follows from diagonalphasedefinition(Q, d, t). □

Lemma 1.11. *For every natural number Q and natural number t,* observersourceflux$(Q, t, 0)$ = 0.

Proof. The goal follows from observersourcefluxdefinition(Q, t). □

Lemma 1.12. *For every natural number Q and natural number t and natural number $t1$,* observersourceflux$(Q, t, \text{succ}(t1))$ = wittxor(wittlinear(observersourceflux$(Q, t, t1)$), wittshl($t1), t1 + 1)$).

Proof. The goal follows from observersourcefluxdefinition2$(Q, t, t1)$. □

Lemma 1.13. *For every natural number L and natural number H,* pageboundarycarry$(L, H, 0)$ = false.

Proof. The goal follows from pageboundarycarrydefinition(L, H). □

Lemma 1.14. *For every natural number L and natural number H and natural number t,* pageboundarycarry$(L, H, \text{succ}(t))$ = rowbit(page$(L, H, t + 1), 0)$ and rowbit(page$(L, H, t), L - 1)$.

Proof. The goal follows from pageboundarycarrydefinition2(L, H, t). □

Lemma 1.15. *For every natural number L and natural number H,* pagefold$(L, H, 0)$ = 0.

Proof. The goal follows from pagefolddefinition(L, H). □

Lemma 1.16. *For every natural number L and natural number H and natural number t,* pagefold$(L, H, \text{succ}(t))$ = wittxor(pagefold(L, H, t), page(L, H, t)).

Proof. The goal follows from pagefolddefinition2(L, H, t). □

Lemma 1.17. *For every natural number Q and natural number t and natural number k,* pathflux$(Q, t, k, 0) = 0$.

Proof. The goal follows from pathfluxdefinition(Q, t, k). □

Lemma 1.18. *For every natural number d,* qordersearch$(d, 0) = 0$.

Proof. The goal follows from qordersearchdefinition(d). □

Lemma 1.19. *For every natural number t and natural number k,* $\mathrm{rayflux}(t, k, 0) = 0$.

Proof. The goal follows from $\mathrm{rayfluxdefinition}(t, k)$. □

Lemma 1.20. *For every truth value a and truth value g and integer $outgoing$ and integer $gate$,* $\mathrm{recoverphase}(a, g, outgoing, gate) = (1 - 2 \cdot \mathrm{bitvalue}(g)) \cdot (outgoing - (1 - 2 \cdot \mathrm{bitvalue}(a)) \cdot gate)$.

Proof. The goal follows from $\mathrm{recoverphasedefinition}(a, g, outgoing, gate)$. □

Lemma 1.21. *For every natural number t,* $\mathrm{rightimport}(t) = \mathrm{not}(\mathrm{rightcell}(t, 1))\,\mathrm{and}\,\mathrm{rightcell}(t, 2)$.

Proof. The goal follows from $\mathrm{rightimportdefinition}(t)$. □

Lemma 1.22. *For every natural number Q and natural number t and natural number k,* $\mathrm{rightphase}(Q, t, k) = \mathrm{bitvalue}(\mathrm{rightcell}(t{+}Q, k)){-}\mathrm{bitvalue}(\mathrm{rightcell}(t, k))$.

Proof. The goal follows from $\mathrm{rightphasedefinition}(Q, t, k)$. □

Lemma 1.23. *For every truth value a and truth value g and integer $incoming$ and integer $gate$,* $\mathrm{transferphase}(a, g, incoming, gate) = (1 - 2 \cdot \mathrm{bitvalue}(g)) \cdot incoming + (1 - 2 \cdot \mathrm{bitvalue}(a)) \cdot gate$.

Proof. The goal follows from $\mathrm{transferphasedefinition}(a, g, incoming, gate)$. □

Lemma 1.24. *For every natural number n,* $\mathrm{twopow}(\mathrm{succ}(n)) = \mathrm{twopow}(n){+}\,\mathrm{twopow}(n)$.

Proof. The goal follows from $\mathrm{twopowdefinition2}(n)$. □

Lemma 1.25. *For every natural number y,* $\mathrm{wittlinear}(y) = \mathrm{wittxor}(\mathrm{wittxor}(y, \mathrm{wittshl}(y, 1)), \mathrm{wittshl}(y, 2$

Proof. The goal follows from $\mathrm{philineardefinition}(y)$. □

Lemma 1.26. *For every natural number $h1$ and natural number m,* $\mathrm{wittlinpow}(\mathrm{succ}(m), h1) = \mathrm{wittlinear}(\mathrm{wittlinpow}(m, h1))$.

Proof. The goal follows from $\mathrm{philinearpowdefinition2}(h1, m)$. □

Lemma 1.27. *For every natural number n,* $\mathrm{wittrow}(\mathrm{succ}(n)) = \mathrm{wittphi}(\mathrm{wittrow}(n))$.

Proof. The goal follows from $\mathrm{rowdefinition2}(n)$. □

Lemma 1.28. *For every natural number Q and natural number t and natural number i,* $\mathrm{alignedphase}(Q, t, i) = \mathrm{pairphase}(\mathrm{alignedbackground}(Q, t), \mathrm{aligneddefect}(Q, t), i)$.

Proof. The goal follows from alignedphasedefinition(Q, t, i). □

Lemma 1.29. *For every natural number h and natural number A,* blockendpoint$(h, A) =$ rowbit$(A, 0)$ and rowbit$(A, h - 1)$.

Proof. The goal follows from blockendpointdefinition(h, A). □

Lemma 1.30. *For every natural number t,* center$(t) =$ cell$(t, 0)$.

Proof. The goal follows from centerdefinition(t). □

Lemma 1.31. *For every natural number Q and natural number d and natural number t and natural number p,* diagonalflux$(Q, d, t, p) =$ rayflux$(t + Q, t + Q - d, t + Q - d - p) -$ rayflux$(t, t - d, t - d - p)$.

Proof. The goal follows from diagonalfluxdefinition(Q, d, t, p). □

Lemma 1.32. *For every natural number Q and natural number t and natural number k,* erasureevent$(Q, t, k) =$ (leftdefect$(Q, t, k+1) =$ true $\wedge$ leftdefect$(Q, t+1, k) =$ false $\wedge$ gatecode$(Q, t, k) \neq 0 \wedge$collisioncode$(Q, t, k) = 0 \wedge$(leftcolour$(Q, t+1, k) = 0 \vee$ leftcolour$(Q, t + 1, k) = 3)$).

Proof. The goal follows from erasureeventdefinition(Q, t, k). □

Lemma 1.33. ldiag$(0, 0) =$ true.

Proof. The goal follows from ldiagdefinition. □

Lemma 1.34. *For every natural number n,* ldiag$(\text{succ}(n), 0) =$ false.

Proof. The goal follows from ldiagdefinition2(n). □

Lemma 1.35. *For every natural number s,* ldiag$(0, \text{succ}(s)) =$ true.

Proof. The goal follows from ldiagdefinition3(s). □

Lemma 1.36. *For every natural number s,* ldiag$(\text{succ}(0), \text{succ}(s)) =$ true.

Proof. The goal follows from ldiagdefinition4(s). □

Lemma 1.37. *For every natural number s and natural number $n1$,* ldiag$(\text{succ}(\text{succ}(n1)), \text{succ}(s)) =$ bxor(ldiag$(n1, s)$, ldiag$(\text{succ}(n1), s)$ or ldiag$(\text{succ}(\text{succ}(n1)), s)$).

Proof. The goal follows from ldiagdefinition5$(s, n1)$. □

Lemma 1.38. *For every natural number Q and natural number t and natural number k,* $\mathrm{leadingadvance}(Q, t, k) = (2 \leq k \wedge \mathrm{advancingatdepth}(Q, t, k) \wedge \mathrm{rightmostleft}(Q, t, k))$.

Proof. The goal follows from $\mathrm{leadingadvancedefinition}(Q, t, k)$. □

Lemma 1.39. *For every truth value b,* $\mathrm{naturalbit}(b) = \mathrm{naturalbit}(b)$.

Proof. The goal follows from $\mathrm{naturalbitdefinition}(b)$. □

Lemma 1.40. *For every natural number t,* $\mathrm{observerinput}(t) = \mathrm{not}(\mathrm{center}(t))$ and $\mathrm{rightcell}(t, 1)$.

Proof. The goal follows from $\mathrm{observerinputdefinition}(t)$. □

Lemma 1.41. *For every natural number Q and natural number t,* $\mathrm{observerqrank}(Q, t) = t + Q - 2 \cdot \mathrm{qorder}(\mathrm{observerdefect}(Q, t))$.

Proof. The goal follows from $\mathrm{observerqrankdefinition}(Q, t)$. □

Lemma 1.42. *For every natural number b and natural number d and natural number i,* $\mathrm{pairphase}(b, d, i) = \mathrm{bitvalue}(\mathrm{rowbit}(\mathrm{wittxor}(b, d), i)) - \mathrm{bitvalue}(\mathrm{rowbit}(b, i))$.

Proof. The goal follows from $\mathrm{pairphasedefinition}(b, d, i)$. □

Lemma 1.43. *For every natural number a and natural number i,* $\mathrm{rowbit}(a, i) = \mathrm{rowbit}(a, i)$.

Proof. The goal follows from $\mathrm{rowbitdefinition}(a, i)$. □

Lemma 1.44. *For every natural number $h1$,* $\mathrm{wittwin}(h1, 0) = 0$.

Proof. The goal follows from $\mathrm{winwittdefinition}(h1)$. □

Lemma 1.45. *For every natural number $h1$ and natural number s,* $\mathrm{wittwin}(h1, \mathrm{succ}(s)) = \mathrm{wittxor}(\mathrm{wittlinear}(\mathrm{wittwin}(h1, s)), \mathrm{wittshl}(\mathrm{wittcarry}(\mathrm{wittrow}(h1 + s)), 1))$.

Proof. The goal follows from $\mathrm{winwittdefinition2}(h1, s)$. □

Lemma 1.46. *For every natural number n and natural number j and truth value b,* $\mathrm{bitcarryeq}(n, j, b) = (\mathrm{wittbit}(\mathrm{wittcarry}(n), j) = \mathrm{bton}(b))$.

Proof. The goal follows from $\mathrm{bitcarryeqdefinition}(n, j, b)$. □

Lemma 1.47. *For every natural number a and natural number b and natural number j and truth value c,* $\mathrm{bitcrosscarryeq}(a, b, j, c) = (\mathrm{wittbit}(\mathrm{crosscarry}(a, b), j) = \mathrm{bton}(c))$.

Proof. The goal follows from bitcrosscarryeqdefinition(a, b, j, c). □

Lemma 1.48. *For every natural number a and natural number b and natural number $j1$ and natural number $j2$ and truth value c,* bitcrosscarryxoreq$(a, b, j1, j2, c) =$ (bxor(rowbit(crosscarry$(a, b), j1)$, rowbit(crosscarry$(a, b), j2)) = c$).

Proof. The goal follows from bitcrosscarryxoreqdefinition$(a, b, j1, j2, c)$. □

Lemma 1.49. *For every natural number k and natural number y and natural number j and truth value b,* bitfrobdefecteq$(k, y, j, b) =$ (wittbit(frobdefect$(k, y), j) =$ bton(b)).

Proof. The goal follows from bitfrobdefecteqdefinition(k, y, j, b). □

Lemma 1.50. *For every natural number $t0$ and natural number m and natural number j and truth value b,* bitwinwitteq$(t0, m, j, b) =$ (wittbit(wittwin$(t0, m), j) =$ bton(b)).

Proof. The goal follows from bitwinwitteqdefinition$(t0, m, j, b)$. □

Lemma 1.51. *For every natural number k and natural number y,* frobdefect$(k, y) =$ wittxor(wittcarry(wittscale$(k, y))$, wittscale$(k,$ wittcarry$(y)))$.

Proof. The goal follows from frobdefectdefinition(k, y). □

Lemma 1.52. *For every natural number $t0$,* maskcount$(t0, 0) =$ false.

Proof. The goal follows from maskcountdefinition$(t0)$. □

Lemma 1.53. *For every natural number t and natural number k,* maskparity$(t, k) =$ maskcount$(t,$ twopow$(k))$.

Proof. The goal follows from maskparitydefinition(t, k). □

Lemma 1.54. *For every natural number t,* rcenter$(t) =$ rdiag(t, t).

Proof. The goal follows from rcenterdefinition(t). □

Lemma 1.55. rdiag$(0, 0) =$ true.

Proof. The goal follows from rdiagdefinition. □

Lemma 1.56. *For every natural number n,* rdiag(succ$(n), 0) =$ false.

Proof. The goal follows from rdiagdefinition2(n). □

Lemma 1.57. *For every natural number s,* rdiag$(0,$ succ$(s)) =$ true.

Proof. The goal follows from rdiagdefinition3(s). □

Lemma 1.58. *For every natural number s,* rdiag(succ(0), succ(s)) = not(rdiag(succ(0), s)).

Proof. The goal follows from rdiagdefinition4(s). □

Lemma 1.59. *For every natural number s and natural number $n1$,* rdiag(succ(succ($n1$)), succ(s)) = bxor(rdiag(succ(succ($n1$)), s), rdiag(succ($n1$), s) or rdiag($n1$, s)).

Proof. The goal follows from rdiagdefinition5(s, $n1$). □

Lemma 1.60. *For every natural number t and natural number k,* response(t, k) = window(t, twopow(k), 0).

Proof. The goal follows from responsedefinition(t, k). □

Lemma 1.61. *For every truth value a and truth value b and truth value c,* rule(a, b, c) = bxor(a, b or c).

Proof. The goal follows from ruledefinition(a, b, c). □

Lemma 1.62. twopow(0) = 1.

Proof. The goal follows from twopowdefinition. □

Lemma 1.63. *For every natural number $h1$ and integer $h2$,* window($h1$, 0, $h2$) = false.

Proof. The goal follows from window60definition($h1$, $h2$). □

Lemma 1.64. *For every natural number y and natural number j,* wittbitbool(y, j) = rowbit(y, j).

Proof. The goal follows from wittbitbooldefinition(y, j). □

Lemma 1.65. *For every natural number t,* wittcenter(t) = wittbit(wittrow(t), t).

Proof. The goal follows from rowcenterdefinition(t). □

Lemma 1.66. *For every natural number $h1$,* wittlinpow(0, $h1$) = $h1$.

Proof. The goal follows from philinearpowdefinition($h1$). □

Lemma 1.67. *For every natural number t and natural number k,* wittresp(t, k) = rowbit(wittwin(t, twopow(k)), t + twopow(k)).

Proof. The goal follows from wittrespbooldefinition(t, k). □

Lemma 1.68. wittrow$(0) = 1$.

Proof. The goal follows from rowdefinition. □

Lemma 1.69. *For every natural number k and natural number y,* wittscale$(k, y) =$ wittxor(wittxor$(y,$ wittshl$(y,$ twopow$(k))),$ wittshl$(y,$ twopow$(k)+$twopow$(k)))$.

Proof. The goal follows from phiscaledefinition(k, y). □

Lemma 1.70. *For every natural number t,* ledgerdiagonal$(t) =$ rowbit(wittwin$(0, t), t)$.

Proof. The goal follows from ledgerdiagonaldefinition(t). □

Lemma 1.71. *For every natural number t and natural number j,* ledgerbit$(t, j) =$ rowbit(wittwin$(0, t), j)$.

Proof. The goal follows from ledgerbitdefinition(t, j). □

Lemma 1.72. *For every natural number t,* edgediagonal$(t) =$ rowbit(edgeledger$(t), t)$.

Proof. The goal follows from edgediagonaldefinition(t). □

Lemma 1.73. *For every natural number t,* interiordiagonal$(t) =$ rowbit(interiorledger$(t), t)$.

Proof. The goal follows from interiordiagonaldefinition(t). □

Lemma 1.74. *For every natural number t,* interior$(t) =$ wittxor(wittcarry(wittrow$(t))$, edgesource$(t))$.

Proof. The goal follows from interiordefinition(t). □

2 Masked observations

Theorem 2.1. *For every natural number t,* whitecount$(t, 0) = 0$.

Proof. The goal follows from whitecountzero(t). □

Theorem 2.2. *For every natural number t and natural number n, if* center$(t) =$ true*, then* whitecount$(t, n + 1) =$ whitecount$(t + 1, n)$.

Proof. The goal follows from whitecountblackstep(t, n). □

Theorem 2.3. *For every natural number t and natural number n, if* center$(t) =$ false*, then* whitecount$(t, n + 1) =$ whitecount$(t + 1, n) + 1$.

Proof. The goal follows from whitecountwhitestep(t, n). □

Theorem 2.4. *For every natural number $t1$ and natural number $t2$ and natural number n, if* centerwindowagree$(t1, t2, n)$*, then* leftprefixagree$(t1, t2, n)$ *if and only if* whitetraceagree$(t1, t2, n)$.

Proof. The goal follows from prefixagreementiffwhitetrace$(t1, t2, n)$. □

Theorem 2.5. *For every natural number T and natural number P and natural number t and natural number n, if $0 < P$, then if* centerperiodic(T, P)*, then if $T \leq t$, then there exists a natural number a such that there exists a natural number b such that $a < b$ and $b \leq$* twopow(whitecount$(t, n)) \wedge$ leftprefixagree$(t + a \cdot P, t + b \cdot P, n)$.

Proof. The goal follows from compressedprefixrepeat(T, P, t, n). □

Theorem 2.6. *For every natural number T and natural number P and natural number t and natural number n and natural number M, if $0 < P$, then if* centerperiodic(T, P)*, then if $T \leq t$, then if* prefixseparated(t, P, n, M)*, then $M \leq$* twopow(whitecount(t, n)).

Proof. The goal follows from separatedprefixbound(T, P, t, n, M). □

Theorem 2.7. *For every natural number T and natural number P and natural number t and natural number n, if $0 < P$, then if* centerperiodic(T, P)*, then if $T \leq t$, then $n < t +$* twopow(whitecount$(t, n)) \cdot P$.

Proof. The goal follows from edgeinformationbound(T, P, t, n). □

Theorem 2.8. *For every natural number n and natural number t and natural number w and natural number P, if $n < (w+1) \cdot P$, then $n < t +$*twopow$(w) \cdot P$.

Proof. The goal follows from informationboundaudit(n, t, w, P). □

3 Central collision recurrence

Theorem 3.1. *For every natural number Q and natural number t and natural number r, if* leftadjacentdefect$(Q, t) =$ true*, then if* center$(t + r) =$ true*, then if* center$(t + r + Q) =$ true*, then* nearcollision$(Q, t, r + 1, 2 \cdot (r + 1))$.

Proof. The goal follows from unitdefectforcesnearcollision(Q, t, r). □

Theorem 3.2. *For every natural number T and natural number P and natural number m and natural number t, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq t$, then if* leftadjacentdefect$(m \cdot P, t) =$ true*, then* nearcollision$(m \cdot P, t, P, 2 \cdot P)$.

Proof. The goal follows from periodicunitdefectforcesnearcollision(T, P, m, t). □

Theorem 3.3. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq N$, then there exists a natural number t such that $N \leq t$ and $t+1 \leq 2 \cdot N + m \cdot P + P \wedge$* centralcollision$(m \cdot P, t, 2 \cdot P)$.

Proof. The goal follows from periodiccentralcollisionwindow(T, P, m, N). □

Theorem 3.4. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then there exists a natural number t such that $N \leq t$ and* centralcollision$(m \cdot P, t, 2 \cdot P)$.

Proof. The goal follows from periodiccentralcollisions(T, P, m, N). □

Theorem 3.5. *For every natural number T and natural number P and natural number m, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then* recurrentcentralcollision$(m \cdot P, 2 \cdot P)$.

Proof. The goal follows from periodicrecurrentcentralcolumn(T, P, m). □

Theorem 3.6. *For every natural number Q and natural number b and natural number r, if $0 < r$, then if* centerwindowagree$(b, b + Q, r + 1)$*, then if* center$(b) =$ true*, then if* leftadjacentdefect$(Q, b+r) =$ true*, then there exists a natural number s such that $2 \cdot s \leq r+1$ and $s \leq r \wedge$*leftcollision$(Q, b+s, r+1)$.

Proof. The goal follows from blackwindowforcesleftcollision(Q, b, r). □

Theorem 3.7. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq N$, then there exists a natural number t such that $N \leq t$ and $t + 1 \leq 2 \cdot N + m \cdot P + 2 \cdot P \wedge$* leftcollision$(m \cdot P, t, P)$.

Proof. The goal follows from periodicleftcollisionwindow(T, P, m, N). □

Theorem 3.8. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then there exists a natural number t such that $N \leq t$ and* leftcollision$(m \cdot P, t, P)$.

Proof. The goal follows from periodicleftcollisions(T, P, m, N). □

Theorem 3.9. *For every natural number T and natural number P and natural number m, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then* recurrentleftcollision$(m \cdot P, P)$.

Proof. The goal follows from periodicrecurrentleftcolumn(T, P, m). □

Theorem 3.10. *For every natural number Q and natural number t and natural number k,* collisioncode$(Q, t, k) \neq 0$ *if and only if* collisionbit$(Q, t, k) =$ true.

Proof. The goal follows from collisioncodenonzero(Q, t, k). □

Theorem 3.11. *For every natural number Q and natural number t and natural number k,* collisioncode$(Q, t, k) \leq 6$.

Proof. The goal follows from collisioncodebound(Q, t, k). □

Theorem 3.12. *For every natural number Q and natural number t and natural number k, if $0 < k$, then* collisioncode$(Q, t, k) = 1 \leftrightarrow$ leftcolour$(Q, t, k) =$ $1 \wedge$leftcolour$(Q, t, k-1) = 2$ *and* (collisioncode$(Q, t, k) = 2 \leftrightarrow$ leftcolour$(Q, t, k) =$ $2 \wedge$leftcolour$(Q, t, k-1) = 1) \wedge$(collisioncode$(Q, t, k) = 3 \leftrightarrow$ leftcolour$(Q, t, k) =$ $1 \wedge$leftcolour$(Q, t, k-1) = 3) \wedge$(collisioncode$(Q, t, k) = 4 \leftrightarrow$ leftcolour$(Q, t, k) =$ $3 \wedge$leftcolour$(Q, t, k-1) = 1) \wedge$(collisioncode$(Q, t, k) = 5 \leftrightarrow$ leftcolour$(Q, t, k) =$ $2 \wedge$leftcolour$(Q, t, k-1) = 3) \wedge$(collisioncode$(Q, t, k) = 6 \leftrightarrow$ leftcolour$(Q, t, k) =$ $3 \wedge$ leftcolour$(Q, t, k-1) = 2)$.

Proof. The goal follows from collisioncodecolours(Q, t, k). □

Theorem 3.13. *For every natural number Q and natural number t and natural number k,* advancingatdepth(Q, t, k) *if and only if* collisioncode$(Q, t, k) =$ 3 *or* collisioncode$(Q, t, k) = 5$.

Proof. The goal follows from advancingcollisioncodes(Q, t, k). □

Theorem 3.14. *For every natural number Q and natural number t and natural number k, if* collisioncode$(Q, t, k) \neq 0$*, then* leftdefect$(Q, t+1, k) =$ leftdefect$(Q, t, k+1)$.

Proof. The goal follows from collisiontransmitsleft(Q, t, k). □

Theorem 3.15. *For every natural number Q and natural number t and natural number k, if* collisioncode$(Q, t, k) \neq 0$*, then* leftgatedifference$(Q, t, k) =$ false.

Proof. The goal follows from collisiongatesagree(Q, t, k). □

Theorem 3.16. *For every natural number Q and natural number t and natural number k, if* $2 \leq k$, *then* $\text{collisioncode}(Q, t, k) = 1 \to \text{leftdefect}(Q, t+1, k-1) = \text{leftcell}(t, k-2)$ *and* $(\text{collisioncode}(Q, t, k) = 2 \to \text{leftdefect}(Q, t+1, k-1) = \text{leftcell}(t+Q, k-2)) \wedge (\text{collisioncode}(Q, t, k) = 3 \to \text{leftdefect}(Q, t+1, k-1) = \text{true}) \wedge (\text{collisioncode}(Q, t, k) = 4 \to \text{leftdefect}(Q, t+1, k-1) = \text{not}(\text{leftcell}(t+Q, k-2))) \wedge (\text{collisioncode}(Q, t, k) = 5 \to \text{leftdefect}(Q, t+1, k-1) = \text{true}) \wedge (\text{collisioncode}(Q, t, k) = 6 \to \text{leftdefect}(Q, t+1, k-1) = \text{not}(\text{leftcell}(t, k-2)))$.

Proof. The goal follows from $\text{collisionrighttransporttable}(Q, t, k)$. □

Theorem 3.17. *For every natural number Q and natural number t and natural number k, if* $0 < k$, *then* $\text{collisioncode}(Q, t, k) = 3 \to \text{leftcolour}(Q, t+1, k-1) = 2$ *and if* $\text{collisioncode}(Q, t, k) = 5$, *then* $\text{leftcolour}(Q, t+1, k-1) = 1$.

Proof. The goal follows from $\text{advancingrightcolourtable}(Q, t, k)$. □

Theorem 3.18. *For every natural number Q and natural number t and natural number k,* $\text{leftdefect}(Q, t+1, k) = \text{not}(\text{leftdefect}(Q, t, k+1))$ *if and only if* $\text{leftgatedifference}(Q, t, k) = \text{true}$.

Proof. The goal follows from $\text{gatedifferenceiffflip}(Q, t, k)$. □

Theorem 3.19. *For every natural number Q and natural number t and natural number k, if* $\text{leftdefect}(Q, t, k+1) = \text{true}$, *then* $\text{leftdefect}(Q, t+1, k) = \text{false}$ *if and only if* $\text{leftgatedifference}(Q, t, k) = \text{true}$.

Proof. The goal follows from $\text{incomingdefecterasediff}(Q, t, k)$. □

Theorem 3.20. *For every natural number Q and natural number t and natural number k, if* $\text{leftdefect}(Q, t, k+1) = \text{true}$, *then if* $\text{leftdefect}(Q, t+1, k) = \text{false}$, *then* $\text{collisioncode}(Q, t, k) = 0$.

Proof. The goal follows from $\text{erasurehasnocollision}(Q, t, k)$. □

Theorem 3.21. *For every natural number Q and natural number k, if* $0 < k$, *then* $\text{leftcolour}(Q, 0, k) = 0$ *or* $\text{leftcolour}(Q, 0, k) = 2$.

Proof. The goal follows from $\text{seedleftcolours}(Q, k)$. □

Theorem 3.22. *For every natural number Q and natural number k, if* $2 \leq k$, *then* $\text{collisioncode}(Q, 0, k) = 0$.

Proof. The goal follows from $\text{seedleftcollisionzero}(Q, k)$. □

Theorem 3.23. collisioncode$(1, 1, 2) = 5$.

Proof. The goal follows from seedadvancingcollisionbirth. □

Theorem 3.24. leftdefect$(3, 0, 3) = \text{true}$ *and* leftdefect$(3, 1, 2) = \text{false} \wedge$ leftgatedifference$(3, 0, 2) = \text{true}$.

Proof. The goal follows from seedgateerasure. □

Theorem 3.25. *For every natural number Q and natural number t and natural number k, if* leftdefect$(Q, t, k) = \text{true}$*, then if* center$(t) = \text{true}$*, then if* center$(t + Q) = \text{true}$*, then* advancingleftcone(Q, t, k).

Proof. The goal follows from leftdefectblackcone(Q, t, k). □

Theorem 3.26. *For every natural number Q and natural number b and natural number r, if $0 < r$, then if* centerwindowagree$(b, b + Q, r + 1)$*, then if* center$(b) = \text{true}$*, then if* leftadjacentdefect$(Q, b+r) = \text{true}$*, then there exists a natural number s such that $2 \cdot s \leq r+1$ and $s \leq r \wedge$*advancingleftcollision$(Q, b+s, r + 1)$.

Proof. The goal follows from blackwindowforcesadvancingcollision(Q, b, r). □

Theorem 3.27. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq N$, then there exists a natural number t such that $N \leq t$ and $t + 1 \leq 2 \cdot N + m \cdot P + 2 \cdot P \wedge$* advancingleftcollision$(m \cdot P, t, P)$.

Proof. The goal follows from periodicadvancingleftcollisionwindow(T, P, m, N). □

Theorem 3.28. *For every natural number T and natural number P and natural number m, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then* recurrentadvancingleftcollision$(m \cdot P, P)$.

Proof. The goal follows from periodicrecurrentadvancingleftcolumn(T, P, m). □

Theorem 3.29. *For every natural number Q and natural number t and natural number k,* leftdefect$(Q, t, k) = \text{false}$ *if and only if* leftcolour$(Q, t, k) = 0$ *or* leftcolour$(Q, t, k) = 3$.

Proof. The goal follows from leftdefectfalsecolours(Q, t, k). □

Theorem 3.30. *For every natural number Q and natural number t and natural number k,* $\mathrm{leftdefect}(Q, t, k) = \mathrm{true}$ *if and only if* $\mathrm{leftcolour}(Q, t, k) = 1$ *or* $\mathrm{leftcolour}(Q, t, k) = 2$.

Proof. The goal follows from $\mathrm{leftdefecttruecolours}(Q, t, k)$. □

Theorem 3.31. *For every natural number Q and natural number t and natural number k, if $t < k$, then* $\mathrm{leftcolour}(Q, t, k) = 0$ *or* $\mathrm{leftcolour}(Q, t, k) = 2$.

Proof. The goal follows from $\mathrm{leftcoloursoutsideseedcone}(Q, t, k)$. □

Theorem 3.32. *For every natural number Q and natural number t and natural number k, if* $\mathrm{collisioncode}(Q, t, k) \neq 0$, *then* $k \leq t + 1$.

Proof. The goal follows from $\mathrm{collisioncodesupport}(Q, t, k)$. □

Theorem 3.33. *For every natural number Q and natural number t and natural number k,* $\mathrm{gatecode}(Q, t, k) \neq 0$ *if and only if* $\mathrm{leftgatedifference}(Q, t, k) = \mathrm{true}$.

Proof. The goal follows from $\mathrm{gatecodenonzero}(Q, t, k)$. □

Theorem 3.34. *For every natural number Q and natural number t and natural number k,* $\mathrm{gatecode}(Q, t, k) \leq 6$.

Proof. The goal follows from $\mathrm{gatecodebound}(Q, t, k)$. □

Theorem 3.35. *For every natural number Q and natural number t and natural number k, if $0 < k$, then* $\mathrm{gatecode}(Q, t, k) = 1 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 1 \wedge \mathrm{leftcolour}(Q, t, k - 1) = 0$ *and* $(\mathrm{gatecode}(Q, t, k) = 2 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 0 \wedge \mathrm{leftcolour}(Q, t, k-1) = 1) \wedge (\mathrm{gatecode}(Q, t, k) = 3 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 1 \wedge \mathrm{leftcolour}(Q, t, k-1) = 1) \wedge (\mathrm{gatecode}(Q, t, k) = 4 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 2 \wedge \mathrm{leftcolour}(Q, t, k-1) = 0) \wedge (\mathrm{gatecode}(Q, t, k) = 5 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 0 \wedge \mathrm{leftcolour}(Q, t, k-1) = 2) \wedge (\mathrm{gatecode}(Q, t, k) = 6 \leftrightarrow \mathrm{leftcolour}(Q, t, k) = 2 \wedge \mathrm{leftcolour}(Q, t, k - 1) = 2)$.

Proof. The goal follows from $\mathrm{gatecodecolours}(Q, t, k)$. □

Theorem 3.36. *For every natural number Q and natural number t and natural number k, if* $\mathrm{leftdefect}(Q, t, k + 1) = \mathrm{true}$, *then if* $\mathrm{leftdefect}(Q, t + 1, k) = \mathrm{false}$, *then* $\mathrm{erasureevent}(Q, t, k)$.

Proof. The goal follows from $\mathrm{erasureeventofstep}(Q, t, k)$. □

Theorem 3.37. *For every natural number Q and natural number t and natural number k, if* erasureevent(Q, t, k), *then* leftcolour$(Q, t + 1, k) = 3$ *if and only if* gatecode$(Q, t, k) \leq 3$ *and* leftcolour$(Q, t, k + 1) = 2$ *or* $4 \leq$ gatecode(Q, t, k) *and* leftcolour$(Q, t, k + 1) = 1$.

Proof. The goal follows from erasureblackphaseiff(Q, t, k). □

Theorem 3.38. *For every natural number Q and natural number t and natural number k, if $t+1 < k$, then if* erasureevent(Q, t, k), *then* leftcolour$(Q, t+1, k) = 0$.

Proof. The goal follows from erasureoutsideseedconewhite(Q, t, k). □

Theorem 3.39. *For every natural number Q and natural number t and natural number k, if $2 \leq k$, then if* advancingatdepth(Q, t, k), *then if* leftdefect$(Q, t+k, 0) =$ false, *then there exists a natural number j such that* firstpatherasure(Q, t, k, j).

Proof. The goal follows from advancingsourcefirsterasure(Q, t, k). □

Theorem 3.40. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P), *then if $T \leq N$, then there exists a natural number u such that there exists a natural number d such that $N \leq u$ and $u + 1 \leq 2 \cdot N + m \cdot P + 3 \cdot P \wedge d + 2 \leq P \wedge$* erasureevent$(m \cdot P, u, d)$.

Proof. The goal follows from periodicerasurewindow(T, P, m, N). □

Theorem 3.41. *For every natural number T and natural number P and natural number m, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P), *then* recurrentsourceerasure$(m \cdot P, P)$.

Proof. The goal follows from periodicrecurrentsourceerasure(T, P, m). □

Theorem 3.42. noperiod *if and only if* norecurrentcentralcollision.

Proof. The goal follows from noperiodiffnorecurrentcentralcollision. □

Theorem 3.43. norowcenterperiod *if and only if* norecurrentcentralcollision.

Proof. The goal follows from norowcenterperiodiffnorecurrentcentralcollision. □

Theorem 3.44. problemone *if and only if* norecurrentcentralcollision.

Proof. The goal follows from problemoneiffnorecurrentcentralcollision. □

Theorem 3.45. *If* norecurrentcentralcollision*, then* noperiod*.*

Proof. Assume h. The goal follows from noperiodofnorecurrentcentralcollision(h). □

Theorem 3.46. *For every natural number T and natural number P, if $0 < P$, then if* norecurrentcentralcollision*, then not* centerperiodic(T, P)*.*

Proof. The goal follows from notcenterperiodicofnorecurrentcentralcollision(T, P). □

Theorem 3.47. *If* norecurrentcentralcollision*, then* problemone*.*

Proof. Assume h. The goal follows from noteventuallyperiodicofnorecurrentcentralcollision(h). □

Theorem 3.48. *For every natural number b and natural number d and natural number i,* pairphase(wittphi(b), wittxor(wittphi(d), wittshl(crosscarry$(b, d), 1)), i) =$ bitvalue(rowbit(wittphi(wittxor$(b, d)), i)) -$ bitvalue(rowbit(wittphi$(b), i)$)*.*

Proof. The goal follows from pairphasestep(b, d, i). □

Theorem 3.49. *For every natural number Q and natural number t and natural number k,* alignedphase$(Q, t, t + k + Q) =$ leftphase(Q, t, k)*.*

Proof. The goal follows from alignedphaseleft(Q, t, k). □

Theorem 3.50. *For every natural number Q and natural number t and natural number k,* leftphase$(Q, t, k) = 0$ *if and only if* leftdefect$(Q, t, k) =$ false*.*

Proof. The goal follows from phasezeroiff(Q, t, k). □

Theorem 3.51. *For every natural number Q and natural number t and natural number k,* leftphase$(Q, t, k) = -1$ *or* leftphase$(Q, t, k) = 1$ *if and only if* leftdefect$(Q, t, k) =$ true*.*

Proof. The goal follows from phaseunitiff(Q, t, k). □

Theorem 3.52. *For every natural number Q and natural number t and natural number k,* leftphase$(Q, t+1, k) =$ leftphase$(Q, t, k+1)+$leftphaseflux(Q, t, k)*.*

Proof. The goal follows from leftphasebalance(Q, t, k). □

Theorem 3.53. *For every natural number Q and natural number t and natural number k, if* $\text{collisioncode}(Q, t, k) \neq 0$*, then* $\text{leftphaseflux}(Q, t, k) = -2 \cdot \text{leftphase}(Q, t, k + 1)$.

Proof. The goal follows from $\text{collisionphaseflip}(Q, t, k)$. □

Theorem 3.54. *For every natural number Q and natural number t and natural number k, if $0 < k$, then if* $\text{advancingatdepth}(Q, t, k)$*, then* $\text{leftphase}(Q, t + 1, k - 1) = -\text{leftphase}(Q, t, k)$.

Proof. The goal follows from $\text{advancingphaseflip}(Q, t, k)$. □

Theorem 3.55. *For every natural number Q and natural number t and natural number k,* $\text{gatecode}(Q, t, k) = 1 \to \text{leftgatephase}(Q, t, k) = -1$ *and* $(\text{gatecode}(Q, t, k) = 2 \to \text{leftgatephase}(Q, t, k) = -1) \wedge (\text{gatecode}(Q, t, k) = 3 \to \text{leftgatephase}(Q, t, k) = -1) \wedge (\text{gatecode}(Q, t, k) = 4 \to \text{leftgatephase}(Q, t, k) = 1) \wedge (\text{gatecode}(Q, t, k) = 5 \to \text{leftgatephase}(Q, t, k) = 1) \wedge (\text{gatecode}(Q, t, k) = 6 \to \text{leftgatephase}(Q, t, k) = 1)$.

Proof. The goal follows from $\text{gatephasecases}(Q, t, k)$. □

Theorem 3.56. *For every natural number Q and natural number t and natural number k, if* $\text{erasureevent}(Q, t, k)$*, then* $\text{leftphase}(Q, t + 1, k) = 0$ *and* $\text{leftphaseflux}(Q, t, k) = -\text{leftphase}(Q, t, k + 1) \wedge (\text{leftphase}(Q, t, k + 1) = -1 \vee \text{leftphase}(Q, t, k + 1) = 1)$.

Proof. The goal follows from $\text{erasurephasebalance}(Q, t, k)$. □

Theorem 3.57. *For every natural number Q and natural number t and natural number k, if* $\text{erasureevent}(Q, t, k)$*, then* $\text{leftcolour}(Q, t + 1, k) = 3 \leftrightarrow \text{leftgatephase}(Q, t, k) \cdot \text{leftphase}(Q, t, k + 1) = -1$ *and* $\text{leftcolour}(Q, t + 1, k) = 0 \leftrightarrow \text{leftgatephase}(Q, t, k) \cdot \text{leftphase}(Q, t, k + 1) = 1$.

Proof. The goal follows from $\text{erasureresetproduct}(Q, t, k)$. □

Theorem 3.58. *For every natural number Q and natural number t and natural number k and natural number n,* $\text{pathphase}(Q, t, k, n) = \text{leftphase}(Q, t, k) + \text{pathflux}(Q, t, k, n)$.

Proof. The goal follows from $\text{pathphasebalance}(Q, t, k, n)$. □

Theorem 3.59. *For every natural number Q and natural number t, if $0 < Q$, then* $\text{leftphase}(Q, t, t + Q) = 1$.

Proof. The goal follows from $\text{seedleftphase}(Q, t)$. □

Theorem 3.60. *For every natural number Q and natural number t, if $0 < Q$, then* $\mathrm{leftphase}(Q, t + (t + Q), 0) = 1 + \mathrm{pathflux}(Q, t, t + Q, t + Q)$.

Proof. The goal follows from $\mathrm{seedcenterphasebalance}(Q, t)$. □

Theorem 3.61. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$, *then if $T \leq N$, then* $\mathrm{pathflux}(m \cdot P, N, N + m \cdot P, N + m \cdot P) = -1$.

Proof. The goal follows from $\mathrm{periodicseedflux}(T, P, m, N)$. □

Theorem 3.62. $\mathrm{leftphase}(3, 0, 3) = 1$ *and* $\mathrm{leftphase}(3, 1, 2) = 0 \wedge \mathrm{leftphaseflux}(3, 0, 2) = -1 \wedge \mathrm{leftgatephase}(3, 0, 2) = 1$.

Proof. The goal follows from seedphasecancellation. □

Theorem 3.63. *For every natural number Q and natural number t and natural number k and natural number s, if* $\mathrm{rightmostleft}(Q, t, k)$, *then if $s < k$, then if* $\mathrm{centercleanthrough}(Q, t, s)$, *then* $\mathrm{rightmostleft}(Q, t + s, k - s)$.

Proof. The goal follows from $\mathrm{rightmostleftprofile}(Q, t, k, s)$. □

Theorem 3.64. *For every natural number Q and natural number t and natural number k, if* $\mathrm{leadingadvance}(Q, t, k)$, *then if* $\mathrm{centercleanthrough}(Q, t, k)$, *then* $\mathrm{firstpatherasure}(Q, t, k, k - 2)$ *and* $\mathrm{erasureevent}(Q, t + (k - 1), 0) \wedge (\mathrm{gatecode}(Q, t + (k - 1), 0) = 2 \vee \mathrm{gatecode}(Q, t + (k - 1), 0) = 5) \wedge \mathrm{center}(t + (k - 1)) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{leadingfirsterasure}(Q, t, k)$. □

Theorem 3.65. *For every natural number Q and natural number b and natural number r, if $0 < r$, then if* $\mathrm{centerwindowagree}(b, b + Q, r + 1)$, *then if* $\mathrm{center}(b) = \mathrm{true}$, *then if* $\mathrm{leftadjacentdefect}(Q, b + r) = \mathrm{true}$, *then there exists a natural number s such that there exists a natural number k such that $2 \cdot s \leq r + 1$ and $s \leq r \wedge k + s \leq r + 1 \wedge \mathrm{leadingadvance}(Q, b + s, k)$.*

Proof. The goal follows from $\mathrm{blackwindowforcesleadingsource}(Q, b, r)$. □

Theorem 3.66. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$, *then if $T \leq N$, then there exists a natural number t such that there exists a natural number k such that $N \leq t$ and $t + 1 \leq 2 \cdot N + m \cdot P + 2 \cdot P \wedge k \leq P \wedge \mathrm{leadingadvance}(m \cdot P, t, k)$.*

Proof. The goal follows from periodicleadingsourcewindow(T, P, m, N). □

Theorem 3.67. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq N$, then there exists a natural number t such that there exists a natural number k such that $N \leq t$ and $t+1 \leq 2 \cdot N + m \cdot P + 2 \cdot P \wedge k \leq P \wedge$*leadingadvance$(m \cdot P, t, k) \wedge$firstpatherasure$(m \cdot P, t, k, k-2) \wedge (\text{gatecode}(m \cdot P, t + (k-1), 0) = 2 \vee \text{gatecode}(m \cdot P, t + (k-1), 0) = 5)$.

Proof. The goal follows from periodicleadingerasurewindow(T, P, m, N). □

Theorem 3.68. *For every natural number T and natural number P and natural number m, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then* recurrentleadingerasure$(m \cdot P, P)$.

Proof. The goal follows from periodicrecurrentleadingerasure(T, P, m). □

Theorem 3.69. *For every natural number Q and natural number R and natural number t and natural number k,* leftphase$(Q+R, t, k) =$ leftphase$(Q, t, k) +$ leftphase$(R, t + Q, k)$.

Proof. The goal follows from leftphaselagadd(Q, R, t, k). □

Theorem 3.70. *For every natural number Q and natural number R and natural number t and natural number k and natural number n,* pathflux$(Q + R, t, k, n) =$ pathflux$(Q, t, k, n) +$ pathflux$(R, t + Q, k, n)$.

Proof. The goal follows from pathfluxlagadd(Q, R, t, k, n). □

Theorem 3.71. *For every natural number Q and natural number t and natural number k, if* leadingadvance(Q, t, k)*, then if* centercleanthrough(Q, t, k)*, then* leftcolour$(Q, t+k, 0) = 3 \leftrightarrow$ leftgatephase$(Q, t+(k-1), 0) \cdot$(leftphase$(Q, t, k) +$ pathflux$(Q, t, k, k-1)) = -1$ *and* leftcolour$(Q, t+k, 0) = 0 \leftrightarrow$ leftgatephase$(Q, t + (k-1), 0) \cdot (\text{leftphase}(Q, t, k) + \text{pathflux}(Q, t, k, k-1)) = 1$.

Proof. The goal follows from leadingphasereset(Q, t, k). □

Theorem 3.72. *For every natural number Q and natural number t, if* leftdefect$(Q, t, 0) =$ false*, then if* leftdefect$(Q, t + 1, 0) =$ false*, then* erasureevent$(Q, t, 0)$ *if and only if* gatecode$(Q, t, 0) = 2$ *or* gatecode$(Q, t, 0) = 5$.

Proof. The goal follows from centererasureiffgate(Q, t). □

Theorem 3.73. *For every natural number Q and natural number b and natural number n and natural number i and natural number j and natural number k, if* commonwhiteblock(Q, b, n)*, then if* $i \leq j$*, then if* $j \leq k$*, then if* $k \leq n$*, then if* gatecode$(Q, b + i, 0) = 2$ *or* gatecode$(Q, b + i, 0) = 5$*, then if* gatecode$(Q, b + k, 0) = 2$ *or* gatecode$(Q, b + k, 0) = 5$*, then* gatecode$(Q, b + j, 0) =$ gatecode$(Q, b + i, 0)$.

Proof. The goal follows from whiteblockgateconvex(Q, b, n, i, j, k). □

Theorem 3.74. *For every natural number Q and natural number b and natural number n and natural number i and natural number j and natural number k, if* commonwhiteblock(Q, b, n)*, then if* centercleanthrough$(Q, b, n+1)$*, then if* $i \leq j$*, then if* $j \leq k$*, then if* $k \leq n$*, then if* erasureevent$(Q, b+i, 0)$*, then if* erasureevent$(Q, b + k, 0)$*, then* erasureevent$(Q, b + j, 0)$ *and* gatecode$(Q, b + j, 0) =$ gatecode$(Q, b + i, 0)$.

Proof. The goal follows from whiteblockerasureconvex(Q, b, n, i, j, k). □

Theorem 3.75. leadingadvance$(4, 5, 2)$ *and* centercleanthrough$(4, 5, 2) \wedge$ gatecode$(4, 6, 0) = 5$.

Proof. The goal follows from seedleadinggatefive. □

Theorem 3.76. leadingadvance$(7, 7, 4)$ *and* centercleanthrough$(7, 7, 4) \wedge$ gatecode$(7, 10, 0) = 2$.

Proof. The goal follows from seedleadinggatetwo. □

Theorem 3.77. leadingadvance$(11, 4, 4)$ *and* centercleanthrough$(11, 4, 4) \wedge$ gatecode$(11, 7, 0) = 5 \wedge$ leftcolour$(11, 8, 0) = 3$.

Proof. The goal follows from seedleadingblackreset. □

Theorem 3.78. *For every truth value incomingBit and truth value laterBit and integer incoming and integer gate,* recoverphase$(incomingBit, laterBit,$ transferphase$(incomingBi$ *incoming*.

Proof. The goal follows from transferrecover$(incomingBit, laterBit, incoming, gate)$. □

Theorem 3.79. *For every truth value incomingBit and truth value laterBit and integer incoming and integer gate,* transferphase$(incomingBit, laterBit, incoming, gate) =$ 0 *if and only if* $incoming =$ recoverphase$(incomingBit, laterBit, 0, gate)$.

Proof. The goal follows from transferkernel($incomingBit, laterBit, incoming, gate$). □

Theorem 3.80. *For every natural number Q and natural number t and natural number k,* leftphase($Q, t{+}1, k$) = transferphase(leftcell($t, k{+}1$), latergate(Q, t, k), leftphase($Q, t, k{+}1$), leftgatephase(Q, t, k)).

Proof. The goal follows from phasetransfer(Q, t, k). □

Theorem 3.81. *For every natural number Q and natural number t and natural number k,* leftphase($Q, t, k{+}1$) = recoverphase(leftcell($t, k{+}1$), latergate(Q, t, k), leftphase($Q, t{+}1, k$), leftgatephase(Q, t, k)).

Proof. The goal follows from phaserecover(Q, t, k). □

Theorem 3.82. *For every natural number Q and natural number t and natural number k, if* erasureevent(Q, t, k)*, then not* leftgatephase(Q, t, k) = 0 *and* leftphase($Q, t, k{+}1$) = recoverphase(leftcell($t, k{+}1$), latergate(Q, t, k), 0, leftgatephase(Q, t, k)).

Proof. The goal follows from erasurerecovery(Q, t, k). □

Theorem 3.83. *For every natural number T and natural number P and natural number m and natural number N and natural number g, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $g = 2$ or $g = 5$, then not not there exists a natural number t such that $N \leq t$ and* erasureevent($m \cdot P, t, 0$) $\wedge$ gatecode($m \cdot P, t, 0$) = g.

Proof. The goal follows from periodicerasureorientation(T, P, m, N, g). □

Theorem 3.84. *For every natural number Q and natural number t and natural number k, if* leadingadvance(Q, t, k)*, then if* centercleanthrough(Q, t, k)*, then there exists a natural number s such that $s + 1 < k$ and* center($t + s$) = true.

Proof. The goal follows from leadingblackbeforearrival(Q, t, k). □

Theorem 3.85. *For every natural number Q and natural number b and natural number r, if $0 < r$, then if* centerwindowagree($b, b + Q, r + 1$)*, then if* center(b) = true*, then if* whitecentergap(b, r)*, then if* leftadjacentdefect($Q, b{+}r$) = true*, then there exists a natural number k such that $k \leq r + 1$ and* leadingadvance(Q, b, k).

Proof. The goal follows from blackgapforcesinitialleading(Q, b, r). □

Theorem 3.86. *For every natural number Q and natural number t, if* leadingadvance$(Q, t, 2)$, *then if* centercleanthrough$(Q, t, 2)$, *then* center$(t) =$ true.

Proof. The goal follows from leadingdepthtwoblack(Q, t). □

Theorem 3.87. *For every natural number Q and natural number b and natural number r and natural number g, if $0 < r$, then if* centerwindowagree$(b, b+Q, r+1)$, *then if* center$(b) =$ true, *then if* whitecentergap(b, r), *then if* erasureevent$(Q, b+r, 0)$, *then if* gatecode$(Q, b+r, 0) = g$, *then there exists a natural number k such that $k \leq r+1$ and* leadingadvance$(Q, b, k) \wedge$ firstpatherasure$(Q, b, k, k-2) \wedge$ gatecode$(Q, b+(k-1), 0) = g$.

Proof. The goal follows from blackgaperasureorigin(Q, b, r, g). □

Theorem 3.88. *For every natural number T and natural number P and natural number m and natural number N and natural number g, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P), *then if $g = 2$ or $g = 5$, then not not there exists a natural number b such that there exists a natural number k such that $N \leq b$ and $k \leq P \wedge$* center$(b) =$ true $\wedge$ whitecentergap$(b, k-1) \wedge$ leadingadvance$(m \cdot P, b, k) \wedge$ firstpatherasure$(m \cdot P, b, k, k-2) \wedge$ gatecode$(m \cdot P, b+(k-1), 0) = g$.

Proof. The goal follows from periodicblackerasureorigin(T, P, m, N, g). □

Theorem 3.89. *For every natural number t,* rightcell$(t+1, 1) =$ bxor(center(t), bxor(rightcell$(t, 1)$, right

Proof. The goal follows from rightimportstep(t). □

Theorem 3.90. *For every natural number Q and natural number t, if* center$(t+Q) =$ center(t), *then* rightdefect$(Q, t+1, 1) =$ bxor(rightdefect$(Q, t, 1)$, rightimportdifference(Q, t)).

Proof. The goal follows from rightdefectimportbalance(Q, t). □

Theorem 3.91. *For every natural number Q and natural number t, if* center$(t) =$ true, *then if* center$(t+Q) =$ true, *then if* rightdefect$(Q, t, 1) =$ true, *then if* rightdefect$(Q, t+1, 1) =$ true, *then* rightphase$(Q, t+1, 1) = -$ rightphase$(Q, t, 1)$.

Proof. The goal follows from blackresetsurvival(Q, t). □

Theorem 3.92. *For every natural number Q and natural number t, if* center$(t) =$ true, *then if* center$(t+Q) =$ true, *then if* rightdefect$(Q, t, 1) =$ false, *then if* rightdefect$(Q, t+1, 1) =$ true, *then* rightcell$(t, 1) =$ false *and* rightcell$(t+Q, 1) =$ false $\wedge$ rightdefect$(Q, t, 2) =$ true $\wedge$ rightphase$(Q, t+1, 1) = -$ rightphase$(Q, t, 2)$.

Proof. The goal follows from blackresetimport(Q, t). □

Theorem 3.93. *For every natural number T and natural number P and natural number m and natural number N and truth value $before$, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$*, then not not there exists a natural number t such that $N \leq t$ and* $\mathrm{rightimport}(t) = before \wedge \mathrm{rightimport}(t + m \cdot P) = \mathrm{not}(before) \wedge \mathrm{rightdefect}(m \cdot P, t + 1, 1) = \mathrm{not}(\mathrm{rightdefect}(m \cdot P, t, 1))$.

Proof. The goal follows from $\mathrm{periodicrightimportorientation}(T, P, m, N, before)$. □

Theorem 3.94. *For every natural number Q and natural number t, if* $\mathrm{center}(t + Q) = \mathrm{center}(t)$*, then if* $\mathrm{rightdefect}(Q, t, 1) = \mathrm{false}$*, then if* $\mathrm{rightdefect}(Q, t + 1, 1) = \mathrm{true}$*, then* $\mathrm{rightcell}(t, 1) = \mathrm{false}$ *and* $\mathrm{rightcell}(t+Q, 1) = \mathrm{false} \wedge \mathrm{rightdefect}(Q, t, 2) = \mathrm{true} \wedge \mathrm{rightimportdifference}(Q, t) = \mathrm{true} \wedge \mathrm{rightphase}(Q, t + 1, 1) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(t))) \cdot \mathrm{rightphase}(Q, t, 2)$.

Proof. The goal follows from $\mathrm{rightheadbirth}(Q, t)$. □

Theorem 3.95. *For every natural number Q and natural number $start$ and natural number $finish$, if* $\mathrm{headimportpath}(Q, start, finish)$*, then if* $\mathrm{centercleanthrough}(Q, start, finish - start)$*, then* $\mathrm{rightphase}(Q, finish, 1) = \mathrm{centersign}(start, finish - start) \cdot \mathrm{rightphase}(Q, start, 2)$.

Proof. The goal follows from $\mathrm{headimportpathphase}(Q, start, finish)$. □

Theorem 3.96. *For every natural number Q and natural number t, if* $\mathrm{leftdefect}(Q, t, 0) = \mathrm{false}$*, then if* $\mathrm{leftdefect}(Q, t + 1, 0) = \mathrm{false}$*, then if* $\mathrm{center}(t) = \mathrm{false}$*, then* $\mathrm{leftphase}(Q, t, 1) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(t + 1))) \cdot \mathrm{rightphase}(Q, t, 1)$.

Proof. The goal follows from $\mathrm{whitecenterphasematching}(Q, t)$. □

Theorem 3.97. *For every natural number T and natural number P and natural number m and natural number N and natural number g, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$*, then if $g = 2$ or $g = 5$, then not not there exists a natural number $start$ such that there exists a natural number b such that there exists a natural number k such that $N \leq start$ and $N \leq b \wedge k \leq P \wedge \mathrm{center}(b) = \mathrm{true} \wedge \mathrm{whitecentergap}(b, k - 1) \wedge \mathrm{leadingadvance}(m \cdot P, b, k) \wedge \mathrm{firstpatherasure}(m \cdot P, b, k, k - 2) \wedge \mathrm{gatecode}(m \cdot P, b + (k - 1), 0) = g \wedge \mathrm{headimportpath}(m \cdot P, start, b + (k - 1)) \wedge \mathrm{leftphase}(m \cdot P, b, k) + \mathrm{pathflux}(m \cdot P, b, k, k - 1) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(b + k))) \cdot \mathrm{centersign}(start, b + (k - 1) - start) \cdot \mathrm{rightphase}(m \cdot P, start, 2)$.

Proof. The goal follows from $\mathrm{periodiccoupledcancellation}(T, P, m, N, g)$. □

Theorem 3.98. *For every natural number Q and natural number b and natural number k, if* $\mathrm{leadingadvance}(Q, b, k)$*, then if* $\mathrm{centercleanthrough}(Q, b, k)$*, then* $\mathrm{leftphase}(Q, b, k) + \mathrm{pathflux}(Q, b, k, k-1) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(b + k))) \cdot \mathrm{leftgatephase}(Q, b + (k-1), 0)$.

Proof. The goal follows from $\mathrm{leadingarrivalphaseeqgatephase}(Q, b, k)$. □

Theorem 3.99. *For every natural number Q and natural number $start$ and natural number $finish$, if* $\mathrm{headimportpath}(Q, start, finish)$*, then if* $\mathrm{centercleanthrough}(Q, start, finish - start)$*, then if* $\mathrm{center}(finish) = \mathrm{false}$*, then* $\mathrm{leftgatephase}(Q, finish, 0) = \mathrm{centersign}(start, finish - start) \cdot \mathrm{rightphase}(Q, start, 2)$.

Proof. The goal follows from $\mathrm{headimportpathgatephase}(Q, start, finish)$. □

Theorem 3.100. *For every natural number Q and natural number $start$ and natural number $finish$, if* $\mathrm{headimportpath}(Q, start, finish)$*, then if* $\mathrm{centercleanthrough}(Q, start, finish - start)$*, then if* $\mathrm{center}(finish) = \mathrm{false}$*, then not* $\mathrm{leftgatephase}(Q, finish, 0) = -(\mathrm{centersign}(start, finish - start) \cdot \mathrm{rightphase}(Q, start, 2))$.

Proof. The goal follows from $\mathrm{headimportpathrejectsnegativegate}(Q, start, finish)$. □

Theorem 3.101. *For every natural number T and natural number P and natural number m and natural number N and natural number g, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$*, then if $g = 2$ or $g = 5$, then not not there exists a natural number start such that there exists a natural number b such that there exists a natural number k such that $N \leq start$ and $N \leq b \wedge k \leq P \wedge \mathrm{center}(b) = \mathrm{true} \wedge \mathrm{whitecentergap}(b, k-1) \wedge \mathrm{leadingadvance}(m \cdot P, b, k) \wedge \mathrm{firstpatherasure}(m \cdot P, b, k, k-2) \wedge \mathrm{gatecode}(m \cdot P, b + (k-1), 0) = g \wedge \mathrm{headimportpath}(m \cdot P, start, b + (k-1)) \wedge \mathrm{leftgatephase}(m \cdot P, b + (k-1), 0) = \mathrm{centersign}(start, b + (k-1) - start) \cdot \mathrm{rightphase}(m \cdot P, start, 2)$.

Proof. The goal follows from $\mathrm{periodiccoupledgatephasesign}(T, P, m, N, g)$. □

Theorem 3.102. *For every natural number T and natural number P and natural number m and natural number N, if $0 < P$, then if $0 < m$, then if* $\mathrm{centerperiodic}(T, P)$*, then if $T \leq N$, then* $\mathrm{pathflux}(m \cdot P + m \cdot P, N, N + (m \cdot P + m \cdot P), N + (m \cdot P + m \cdot P)) = -1$ *and* $\mathrm{pathflux}(m \cdot P, N, N + (m \cdot P + m \cdot P), N + (m \cdot P + m \cdot P)) = 0 \wedge \mathrm{pathflux}(m \cdot P, N + m \cdot P, N + (m \cdot P + m \cdot P), N + (m \cdot P + m \cdot P)) = -1$.

Proof. The goal follows from periodicdoubledseedfluxsplit(T, P, m, N). □

Theorem 3.103. leadingadvance$(11, 4, 4)$ *and* centercleanthrough$(11, 4, 4) \wedge$ headimportpath$(11, 6, 7) \wedge$centercleanthrough$(11, 6, 1) \wedge$leftgatephase$(11, 7, 0) = 1 \wedge$ centersign$(6, 1) = 1 \wedge$ rightphase$(11, 6, 2) = 1$.

Proof. The goal follows from seedcoupledgatesign. □

Theorem 3.104. *For every natural number Q and natural number t and natural number j, if* rightdefect$(Q, t+1, j+1) =$ true*, then* rightdefect$(Q, t, j) =$ true *or* rightdefect$(Q, t, j + 1) =$ true $\vee$ rightdefect$(Q, t, j + 2) =$ true.

Proof. The goal follows from rightdefecthasancestor(Q, t, j). □

Theorem 3.105. *For every natural number Q and natural number t and natural number j and natural number T, if $1 \leq j$, then if $T \leq t$, then if* rightdefect$(Q, t, j) =$ true*, then if* centercleanthrough$(Q, T, t - T)$*, then* rightlineage(Q, t, j, T).

Proof. The goal follows from rightlineageofcleancenter(Q, t, j, T). □

Theorem 3.106. *For every natural number Q and natural number t and natural number k and natural number T, if $1 \leq k$, then if $T \leq t$, then if* leftdefect$(Q, t, k) =$ true*, then if* centercleanthrough$(Q, T, t - T)$*, then* leftlineage(Q, t, k, T).

Proof. The goal follows from leftlineageofcleancenter(Q, t, k, T). □

Theorem 3.107. *For every natural number Q and natural number t and natural number j and natural number T, if $T \leq t$, then if* rightlineage(Q, t, j, T)*, then there exists a natural number i such that $1 \leq i$ and $i \leq j + (t - T) \wedge j \leq i + (t - T) \wedge$* rightdefect$(Q, T, i) =$ true.

Proof. The goal follows from rightlineageatstart(Q, t, j, T). □

Theorem 3.108. *For every natural number Q and natural number t and natural number k and natural number T, if $T \leq t$, then if* leftlineage(Q, t, k, T)*, then there exists a natural number i such that $1 \leq i$ and $i \leq k + (t - T) \wedge k \leq i + (t - T) \wedge$* leftdefect$(Q, T, i) =$ true.

Proof. The goal follows from leftlineageatstart(Q, t, k, T). □

Theorem 3.109. *For every natural number Q and natural number t and natural number j, if* $\mathrm{rightlineage}(Q, t, j, 0)$*, then there exists a natural number i such that $1 \leq i$ and $i \leq Q \wedge i \leq j + t \wedge j \leq i + t \wedge \mathrm{rightdefect}(Q, 0, i) = \mathrm{true} \wedge \mathrm{rightcell}(Q, i) = \mathrm{true} \wedge \mathrm{rightphase}(Q, 0, i) = 1$.*

Proof. The goal follows from $\mathrm{rightlineageseed}(Q, t, j)$. □

Theorem 3.110. *For every natural number Q and natural number t and natural number k, if* $\mathrm{leftlineage}(Q, t, k, 0)$*, then there exists a natural number i such that $1 \leq i$ and $i \leq Q \wedge i \leq k + t \wedge k \leq i + t \wedge \mathrm{leftdefect}(Q, 0, i) = \mathrm{true} \wedge \mathrm{leftcell}(Q, i) = \mathrm{true} \wedge \mathrm{leftphase}(Q, 0, i) = 1$.*

Proof. The goal follows from $\mathrm{leftlineageseed}(Q, t, k)$. □

Theorem 3.111. *For every natural number Q and natural number t, if $0 < Q$, then there exists a natural number y such that $y \leq t + Q$ and* $\mathrm{rightfront}(Q, t, y)$*.*

Proof. The goal follows from $\mathrm{existsrightfront}(Q, t)$. □

Theorem 3.112. *For every natural number Q and natural number t and natural number y and natural number z, if* $\mathrm{leftdefect}(Q, t, 0) = \mathrm{false}$*, then if* $\mathrm{rightfront}(Q, t, y)$*, then if* $\mathrm{rightfront}(Q, t + 1, z)$*, then $y \leq z + 1$.*

Proof. The goal follows from $\mathrm{rightfrontlower}(Q, t, y, z)$. □

Theorem 3.113. *For every natural number Q and natural number t and natural number y, if* $\mathrm{leftdefect}(Q, t, 0) = \mathrm{false}$*, then if* $\mathrm{rightfront}(Q, t, y)$*, then if $2 \leq y$, then* $\mathrm{rightdefect}(Q, t + 1, y - 1) = \mathrm{not}(\mathrm{rightcell}(t, y - 1))$*.*

Proof. The goal follows from $\mathrm{rightfrontdescent}(Q, t, y)$. □

Theorem 3.114. *For every natural number Q and natural number t and natural number y and natural number z, if* $\mathrm{leftdefect}(Q, t, 0) = \mathrm{false}$*, then if* $\mathrm{rightfront}(Q, t, y)$*, then if $2 \leq y$, then if* $\mathrm{rightfront}(Q, t + 1, z)$*, then if $z + 1 \leq y$, then $z = y - 1$ and* $\mathrm{rightcell}(t, y - 1) = \mathrm{false}$*.*

Proof. The goal follows from $\mathrm{rightfrontlockstep}(Q, t, y, z)$. □

Theorem 3.115. *For every natural number Q and natural number b and natural number k and natural number s, if* $\mathrm{rightmostleft}(Q, b, k)$*, then if* $\mathrm{centercleanthrough}(Q, b, k)$*, then if $s+1 < k$, then if* $\mathrm{rightdefect}(Q, b+s, 1) = \mathrm{true}$*, then* $\mathrm{center}(b + s) = \mathrm{true}$*.*

Proof. The goal follows from $\mathrm{leadingheadblackbeforearrival}(Q, b, k, s)$. □

Theorem 3.116. *For every natural number Q and natural number b and natural number k and natural number start, if $3 \leq k$, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k-1)$*, then if* headimportpath$(Q, start, b + (k-1))$*, then $start = b + (k-2)$.*

Proof. The goal follows from coupledimportbornatarrival$(Q, b, k, start)$. □

Theorem 3.117. *For every natural number Q and natural number b and natural number k and natural number s, if $0 < Q$, then if $3 \leq k$, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k-1)$*, then if $1 \leq s$, then if $s+1 < k$, then there exists a natural number y such that* rightdefect$(Q, b+s, 1) =$ false *and* rightfront$(Q, b+s, y) \wedge 2 \leq y \wedge y \leq k - s$.

Proof. The goal follows from coupledrightfrontmirrorat(Q, b, k, s). □

Theorem 3.118. *For every natural number Q and natural number b and natural number k and natural number s, if $0 < Q$, then if $3 \leq k$, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k-1)$*, then if $1 \leq s$, then if $s + 2 < k$, then if* rightfront$(Q, b+s, k-s)$*, then* rightcell$(b+s, k-s-1) =$ false *and* rightfront$(Q, b+(s+1), k-(s+1))$.

Proof. The goal follows from coupledwhitestaircaseat(Q, b, k, s). □

Theorem 3.119. *For every natural number T and natural number P and natural number m and natural number N and natural number g, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $g = 2$ or $g = 5$, then not not there exists a natural number start such that there exists a natural number b such that there exists a natural number k such that there exists a natural number y such that $N \leq start$ and $N \leq b \wedge T \leq b \wedge T \leq start \wedge 2 \leq k \wedge k \leq P \wedge$* center$(b) =$ true $\wedge$ whitecentergap$(b, k-1) \wedge$ leadingadvance$(m \cdot P, b, k) \wedge$ gatecode$(m \cdot P, b+(k-1), 0) = g \wedge$ headimportpath$(m \cdot P, start, b+(k-1)) \wedge (3 \leq k \to start = b+(k-2)) \wedge (3 \leq k \to$ rightfront$(m \cdot P, b+1, y) \wedge 2 \leq y \wedge y \leq k-1) \wedge$ rightlineage$(m \cdot P, start, 2, T) \wedge$ leftlineage$(m \cdot P, b, k, T)$.

Proof. The goal follows from periodiccoupledimportancestry(T, P, m, N, g). □

Theorem 3.120. leadingadvance$(14, 5, 3)$ *and* centercleanthrough$(14, 5, 3) \wedge$ whitecentergap$(5, 2) \wedge$center$(5) =$ true $\wedge$ rightfront$(14, 6, 2) \wedge$headimportpath$(14, 6, 7)$.

Proof. The goal follows from seedcoupledimportatarrival. □

Theorem 3.121. *For every natural number t and natural number k and natural number n,* $\mathrm{bitvalue}(\mathrm{raybit}(t,k,n)) = \mathrm{bitvalue}(\mathrm{raybit}(t,k,0)) + \mathrm{rayflux}(t,k,n)$.

Proof. The goal follows from $\mathrm{raybitbalance}(t,k,n)$. □

Theorem 3.122. *For every natural number Q and natural number d and natural number t and natural number p, if $d+p \leq t$, then* $\mathrm{leftphase}(Q+Q, t+(t-d-p), p) = \mathrm{diagonalphase}(Q,d,t) + \mathrm{diagonalflux}(Q,d,t,p)$.

Proof. The goal follows from $\mathrm{diagonalraybalance}(Q,d,t,p)$. □

Theorem 3.123. *For every natural number Q and natural number R and natural number d and natural number t and natural number p,* $\mathrm{diagonalflux}(Q+R,d,t,p) = \mathrm{diagonalflux}(Q,d,t,p) + \mathrm{diagonalflux}(R,d,t+Q,p)$.

Proof. The goal follows from $\mathrm{diagonalfluxlagadd}(Q,R,d,t,p)$. □

Theorem 3.124. *For every natural number Q and natural number z and natural number t, if* $\mathrm{zerodiagonal}(z)$*, then if* $\mathrm{doublesat}(Q,z)$*, then if* $\mathrm{settledby}(z+1,t)$*, then* $\mathrm{diagonalphase}(Q,z+1,t) = -1 \vee \mathrm{diagonalphase}(Q,z+1,t) = 1$ *and* $\mathrm{diagonalphase}(Q,z+1,t+Q) = -\mathrm{diagonalphase}(Q,z+1,t) \wedge \mathrm{diagonalphase}(Q+Q,z+1,t) = 0$.

Proof. The goal follows from $\mathrm{diagonalphasedoubles}(Q,z,t)$. □

Theorem 3.125. *For every natural number Q and natural number z and natural number t, if* $\mathrm{zerodiagonal}(z)$*, then if* $\mathrm{settledby}(z+2,t)$*, then* $\mathrm{diagonalphase}(Q,z+3,t+1) = -\mathrm{diagonalphase}(Q,z+1,t)$.

Proof. The goal follows from $\mathrm{diagonalresetphase}(Q,z,t)$. □

Theorem 3.126. *For every natural number Q and natural number z and natural number t, if* $\mathrm{zerodiagonal}(z)$*, then if* $\mathrm{settledby}(z+2,t)$*, then if* $z+3 \leq t$*, then* $\mathrm{diagonalflux}(Q,z+3,t+1,1) = \mathrm{diagonalflux}(Q,z+1,t,1) + 2 \cdot \mathrm{diagonalphase}(Q,z+1,t)$.

Proof. The goal follows from $\mathrm{diagonalresetfluxjump}(Q,z,t)$. □

Theorem 3.127. *For every natural number T and natural number P and natural number m and natural number d and natural number t, if* $\mathrm{centerperiodic}(T,P)$*, then if* $d+1 \leq t$*, then if* $T \leq t+(t-d-1)$*, then if* $\mathrm{center}(t+(t-d-1)) = \mathrm{false}$*, then* $\mathrm{diagonalphase}(m \cdot P, d, t) + \mathrm{diagonalflux}(m \cdot P, d, t, 1) = (1 - 2 \cdot \mathrm{bitvalue}(\mathrm{center}(t+(t-d-1)+1))) \cdot \mathrm{rightphase}(m \cdot P + m \cdot P, t+(t-d-1), 1)$.

Proof. The goal follows from periodicdiagonalwhiteheadbalance(T, P, m, d, t). □

Theorem 3.128. *For every natural number T and natural number P and natural number m and natural number d and natural number t and natural number start, if* centerperiodic(T, P)*, then if* $d + 1 \leq t$*, then if* $T \leq start$*, then if* headimportpath$(m \cdot P + m \cdot P, start, t + (t - d - 1))$*, then if* center$(t + (t - d - 1)) =$ false*, then* diagonalphase$(m \cdot P, d, t) +$ diagonalflux$(m \cdot P, d, t, 1) = (1 - 2 \cdot$ bitvalue(center$(t + (t - d - 1) + 1))) \cdot ($centersign$(start, t + (t - d - 1) - start) \cdot$ rightphase$(m \cdot P + m \cdot P, start, 2))$.

Proof. The goal follows from periodicdiagonalimportbalance$(T, P, m, d, t, start)$. □

Theorem 3.129. *For every natural number Q and natural number b and natural number k and natural number start and natural number q, if* $0 < Q$*, then if* $3 \leq k$*, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k-1)$*, then if* headimportpath$(Q, start, b+(k-1))$*, then if* $1 \leq q$*, then if* $q + 1 < k$*, then* rightfront$(Q, b + 1, k - 1)$ *and* rightcell$(b + 1, q) =$ false $\wedge$ rightcell$(b + 1 + Q, q) =$ false*.*

Proof. The goal follows from coupledrightfrontexact$(Q, b, k, start, q)$. □

Theorem 3.130. *For every natural number Q and natural number b and natural number k and natural number start and natural number i, if* $0 < Q$*, then if* $3 \leq k$*, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k - 1)$*, then if* headimportpath$(Q, start, b + (k - 1))$*, then if* $1 \leq i$*, then if* $i + 1 < k$*, then* rightfront$(Q, b + i, k - i)$ *and* rightcell$(b + i, k - i - 1) =$ false $\wedge$ rightcell$(b + i + Q, k - i - 1) =$ false*.*

Proof. The goal follows from coupledforcedwhitestaircase$(Q, b, k, start, i)$. □

Theorem 3.131. *For every natural number Q and natural number b and natural number start, if* leadingadvance$(Q, b, 2)$*, then if* centercleanthrough$(Q, b, 2)$*, then if* headimportpath$(Q, start, b+1)$*, then* center$(b) =$ true *and* center$(b+1) =$ false $\wedge$($start < b \wedge$ rightdefect$(Q, b, 1) =$ true $\wedge$ rightimportdifference$(Q, b) =$ false $\wedge$ leftphase$(Q, b, 2) = (1-2 \cdot$ bitvalue(center$(b+2))) \cdot$ rightphase$(Q, b, 1) \vee$ $start = b \wedge$ rightdefect$(Q, b, 1) =$ false $\wedge$ rightcell$(b, 1) =$ false $\wedge$ rightcell$(b + Q, 1) =$ false $\wedge$ rightdefect$(Q, b, 2) =$ true $\wedge$ rightimportdifference$(Q, b) =$ true $\wedge$ leftphase$(Q, b, 2) = (1-2 \cdot$ bitvalue(center$(b+2))) \cdot$ rightphase$(Q, b, 2))$.

Proof. The goal follows from depthtwocancellationcases$(Q, b, start)$. □

Theorem 3.132. *For every natural number Q and natural number b and natural number k and natural number start, if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* whitecentergap$(b, k-1)$*, then if* headimportpath$(Q, start, b+(k-1))$*, then start* $= b+(k-2)$ *or* $k = 2$ *and start* $< b \wedge$ rightdefect$(Q, b, 1) =$ true.

Proof. The goal follows from coupledimportbirthordepthtwosurvival$(Q, b, k, start)$. □

Theorem 3.133. *For every natural number Q and natural number b and natural number k and natural number start and natural number q, if* $0 < Q$*, then if* $3 \leq k$*, then if* leadingadvance(Q, b, k)*, then if* centercleanthrough(Q, b, k)*, then if* center$(b) =$ true*, then if* whitecentergap$(b, k-1)$*, then if* headimportpath$(Q, start, b+(k-1))$*, then if* $q+2 < k$*, then* rightcell$(b, q) =$ true *and* rightcell$(b+Q, q) =$ true.

Proof. The goal follows from coupledblackrootprefix$(Q, b, k, start, q)$. □

Theorem 3.134. leadingadvance$(4, 5, 2)$ *and* centercleanthrough$(4, 5, 2) \wedge$ headimportpath$(4, 2, 6) \wedge$ rightdefect$(4, 5, 1) =$ true $\wedge$ rightimportdifference$(4, 5) =$ false.

Proof. The goal follows from seeddepthtwosurvival. □

Theorem 3.135. *For every natural number Q and natural number t,* alignedbackground$(Q, t+1) =$ wittphi(alignedbackground(Q, t)).

Proof. The goal follows from alignedbackgroundstep(Q, t). □

Theorem 3.136. *For every natural number Q and natural number t,* aligneddefect$(Q, t+1) =$ defectstep(alignedbackground(Q, t), aligneddefect(Q, t)).

Proof. The goal follows from aligneddefectstep(Q, t). □

Theorem 3.137. *For every natural number Q and natural number t,* rowbit(alignedbackground$(Q, t), 2 \cdot t + Q) =$ true.

Proof. The goal follows from backgroundtopbit(Q, t). □

Theorem 3.138. *For every natural number Q and natural number t, if* $0 < Q$*, then* rowbit(aligneddefect$(Q, t), 2 \cdot (t + Q)) =$ true.

Proof. The goal follows from aligneddefecttopbit(Q, t). □

Theorem 3.139. *For every natural number Q and natural number t, if* $0 < Q$, *then* $\mathrm{twopow}(2 \cdot (t + Q)) \leq \mathrm{aligneddefect}(Q, t)$ *and* $\mathrm{aligneddefect}(Q, t) < \mathrm{twopow}(2 \cdot (t + Q) + 1)$.

Proof. The goal follows from $\mathrm{aligneddefectmagnitude}(Q, t)$. □

Theorem 3.140. *For every natural number Q and natural number t and natural number j, if* $2 \cdot (t+Q) < j$, *then* $\mathrm{rowbit}(\mathrm{aligneddefect}(Q, t), j) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{aligneddefectabovetop}(Q, t, j)$. □

Theorem 3.141. *For every natural number Q and natural number t and natural number j, if* $j < Q$, *then* $\mathrm{rowbit}(\mathrm{aligneddefect}(Q, t), j) = \mathrm{rowbit}(\mathrm{wittrow}(t+ Q), j)$.

Proof. The goal follows from $\mathrm{aligneddefectlowprefix}(Q, t, j)$. □

Theorem 3.142. *For every natural number b and natural number d,* $\mathrm{defectstep}(b, d) = \mathrm{wittxor}(\mathrm{wittlinear}(d), \mathrm{wittshl}(\mathrm{wittxor}(\mathrm{wittcarry}(d), \mathrm{crosscarry}(b, d)), 1))$.

Proof. The goal follows from $\mathrm{defectsteplinear}(b, d)$. □

Theorem 3.143. *For every natural number b and natural number d,* $\mathrm{rowbit}(\mathrm{defectstep}(b, d), 0) = \mathrm{rowbit}(d, 0)$.

Proof. The goal follows from $\mathrm{defectsteplowbit}(b, d)$. □

Theorem 3.144. *For every natural number b and natural number d,* $\mathrm{vala}(\mathrm{defectstep}(b, d)) = \mathrm{valb}(\mathrm{wittxor}(\mathrm{wittcarry}(d), \mathrm{crosscarry}(b, d)))$ *and* $\mathrm{valb}(\mathrm{defectstep}(b, d)) = \mathrm{bxor}(\mathrm{vala}(\mathrm{wittxor}(\mathrm{wittcarry}(d$

Proof. The goal follows from $\mathrm{defectstepvalcoordinates}(b, d)$. □

Theorem 3.145. *For every natural number j,* $\mathrm{vala}(\mathrm{wittshl}(1, j + 3)) = \mathrm{vala}(\mathrm{wittshl}(1, j))$ *and* $\mathrm{valb}(\mathrm{wittshl}(1, j+3)) = \mathrm{valb}(\mathrm{wittshl}(1, j)) \wedge (\mathrm{vala}(\mathrm{wittshl}(1, j)) = \mathrm{true} \vee \mathrm{valb}(\mathrm{wittshl}(1, j)) = \mathrm{true})$.

Proof. The goal follows from $\mathrm{edgevalcoordinates}(j)$. □

Theorem 3.146. *For every natural number t and natural number k,* $\mathrm{aligneddefect}(\mathrm{twopow}(k), t) = \mathrm{wittxor}(\mathrm{wittwin}(t, \mathrm{twopow}(k)), \mathrm{wittxor}(\mathrm{wittrow}(t), \mathrm{wittshl}(\mathrm{wittrow}(t), \mathrm{twopow}(k)+ \mathrm{twopow}(k))))$.

Proof. The goal follows from $\mathrm{aligneddefectwindow}(t, k)$. □

Theorem 3.147. *For every natural number t and natural number k, if* $t <$ twopow(k)*, then* rowbit(aligneddefect(twopow$(k), t), 0)$ = true *and* rowbit(wittwin$(t,$ twopow$(k)), 0) =$ false $\wedge$ rowbit(aligneddefect(twopow$(k), t), 2\cdot(t+$twopow$(k))) =$ true $\wedge$ rowbit(wittwin$(t,$ twopow$(k)), 2\cdot$ $(t +$ twopow$(k))) =$ false $\wedge$ vala(wittwin$(t,$ twopow$(k))) =$ vala(wittrow$(t +$ twopow$(k)))\wedge$valb(wittwin$(t,$ twopow$(k))) =$ valb(wittrow$(t+$twopow$(k)))\wedge$ rowbit(aligneddefect(twopow$(k), t), t+$twopow$(k)) =$ rowbit(wittwin$(t,$ twopow$(k)), t+$ twopow$(k))$.

Proof. The goal follows from dyadicwindowreadings(t, k). □

Theorem 3.148. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if* $T \leq$ t*, then* rowbit(wittphi(aligneddefect$(m\cdot P, t)), t+m\cdot P+1) =$ rowbit(crosscarry(alignedbackground$(m\cdot$ $P, t),$ aligneddefect$(m \cdot P, t)), t + m \cdot P)$.

Proof. The goal follows from periodiccentercrosscarry(T, P, m, t). □

Theorem 3.149. aligneddefect$(1, 0) = 5$ *and* vala(aligneddefect$(1, 0)) =$ false $\wedge$ valb(aligneddefect$(1, 0)) =$ true $\wedge$ aligneddefect$(1, 2) = 93\wedge$vala(aligneddefect$(1, 2)) =$ false $\wedge$ valb(aligneddefect$(1, 2)) =$ false $\wedge$ rowbit(aligneddefect$(1, 2), 3) =$ true $\wedge$ aligneddefect$(4, 0) =$ $385\wedge$vala(aligneddefect$(4, 0)) =$ false $\wedge$ valb(aligneddefect$(4, 0)) =$ false $\wedge$ rowbit(aligneddefect$(4, 0), 4) =$ false.

Proof. The goal follows from seedvalcancellation. □

Theorem 3.150. *For every natural number d and natural number n and natural number j, if $j < n$ or $n+2 < j$, then* rowbit(wittxor$(d,$ centerflipmask$(n)), n) =$ not(rowbit$(d, n))$ *and* rowbit(wittxor$(d,$ centerflipmask$(n)), j) =$ rowbit$(d, j)\wedge$ vala(wittxor$(d,$ centerflipmask$(n))) =$ vala$(d)\wedge$valb(wittxor$(d,$ centerflipmask$(n))) =$ valb(d).

Proof. The goal follows from centerflipmaskinvisible(d, n, j). □

Theorem 3.151. *For every natural number t and natural number j,* rowbit(observercode$(t), j) =$ leftcell(t, j).

Proof. The goal follows from observercodereadout(t, j). □

Theorem 3.152. *For every natural number Q and natural number t,* observerdefect$(Q, t) =$ wittxor(observercode$(t + Q),$ observercode$(t))$.

Proof. The goal follows from observerdefectalignment(Q, t). □

Theorem 3.153. *For every natural number Q and natural number t and natural number j,* $\mathrm{rowbit}(\mathrm{observerdefect}(Q,t),j) = \mathrm{rowbit}(\mathrm{aligneddefect}(Q,t),t+Q+j)$.

Proof. The goal follows from $\mathrm{observerdefectreadout}(Q,t,j)$. □

Theorem 3.154. *For every natural number t,* $\mathrm{vala}(\mathrm{observercode}(t+1)) = \mathrm{bxor}(\mathrm{center}(t),\mathrm{bxor}(\mathrm{vala}(\mathrm{wittcarry}(\mathrm{observercode}(t))),\mathrm{observerinput}(t)))$ *and* $\mathrm{valb}(\mathrm{observercode}(t+1)) = \mathrm{bxor}(\mathrm{center}(t),\mathrm{valb}(\mathrm{wittcarry}(\mathrm{observercode}(t))))$.

Proof. The goal follows from $\mathrm{observervalstep}(t)$. □

Theorem 3.155. *For every natural number T and natural number P and natural number m and natural number t, if* $\mathrm{centerperiodic}(T,P)$*, then if* $T \leq t$*, then* $\mathrm{vala}(\mathrm{observerdefect}(m\cdot P,t+1)) = \mathrm{bxor}(\mathrm{vala}(\mathrm{wittxor}(\mathrm{wittcarry}(\mathrm{observercode}(t+m\cdot P)),\mathrm{wittcarry}(\mathrm{observercode}(t)))),\mathrm{leftdefect}(m\cdot P,t,1))$ *and* $\mathrm{valb}(\mathrm{observerdefect}(m\cdot P,t+1)) = \mathrm{valb}(\mathrm{wittxor}(\mathrm{wittcarry}(\mathrm{observercode}(t+m\cdot P)),\mathrm{wittcarry}(\mathrm{observercode}(t))))$.

Proof. The goal follows from $\mathrm{periodicobservervalstep}(T,P,m,t)$. □

Theorem 3.156. *For every natural number T and natural number P and natural number m and natural number t, if* $0 < P$*, then if* $0 < m$*, then if* $\mathrm{centerperiodic}(T,P)$*, then if* $T \leq t$*, then* $\mathrm{rowbit}(\mathrm{observerdefect}(m\cdot P,t),t+m\cdot P) = \mathrm{true}$ *and* $\mathrm{rowbit}(\mathrm{observerdefect}(m\cdot P,t),0) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{periodicobserveredges}(T,P,m,t)$. □

Theorem 3.157. $\mathrm{observerdefect}(2,0) = 7$ *and* $\mathrm{vala}(\mathrm{observerdefect}(2,0)) = \mathrm{false} \wedge \mathrm{valb}(\mathrm{observerdefect}(2,0)) = \mathrm{false} \wedge \mathrm{rowbit}(\mathrm{observerdefect}(2,0),0) = \mathrm{true} \wedge \mathrm{observerdefect}(4,0) = 24 \wedge \mathrm{vala}(\mathrm{observerdefect}(4,0)) = \mathrm{true} \wedge \mathrm{valb}(\mathrm{observerdefect}(4,0)) = \mathrm{true} \wedge \mathrm{rowbit}(\mathrm{observerdefect}(4,0),0) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{observerseedzeroval}$. □

4 Cyclotomic observer transport

Theorem 4.1. *For every natural number r and natural number a,* $\mathrm{qdivides}(r,\mathrm{wittlinpow}(r,a))$.

Proof. The goal follows from $\mathrm{qdividesintro}(r,a)$. □

Theorem 4.2. *For every natural number r and natural number d, if* $\mathrm{qdivides}(r,d)$*, then there exists a natural number a such that* $\mathrm{wittlinpow}(r,a) = d$.

Proof. The goal follows from $\mathrm{qdivideselim}(r,d)$. □

Theorem 4.3. *For every natural number r and natural number d and natural number n,* $\mathrm{qdivides}(r, \mathrm{wittshl}(d, n))$ *if and only if* $\mathrm{qdivides}(r, d)$.

Proof. The goal follows from $\mathrm{qdividesshl}(r, d, n)$. □

Theorem 4.4. *For every natural number r and natural number d, if $0 < r$, then if* $\mathrm{qdivides}(r, d)$, *then* $\mathrm{vala}(d) = \mathrm{false}$ *and* $\mathrm{valb}(d) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{qdividesval}(r, d)$. □

Theorem 4.5. *For every natural number r and natural number d, if $d \neq 0$, then if* $\mathrm{qdivides}(r, d)$, *then* $2 \cdot r \leq \mathrm{binarydegree}(d)$.

Proof. The goal follows from $\mathrm{qdividesdegreebound}(r, d)$. □

Theorem 4.6. *For every natural number d, if $d \neq 0$, then there exists a natural number r such that* $\mathrm{qdivides}(r, d)$ *and not* $\mathrm{qdivides}(r + 1, d)$.

Proof. The goal follows from $\mathrm{qorderexists}(d)$. □

Theorem 4.7. *For every natural number r and natural number d, if $d \neq 0$, then* $\mathrm{qdivides}(r, d)$ *if and only if* $r \leq \mathrm{qorder}(d)$.

Proof. The goal follows from $\mathrm{qdividesifforder}(r, d)$. □

Theorem 4.8. *For every natural number Q and natural number t,* $\mathrm{observersourceflux}(Q, t, 0) = 0$.

Proof. The goal follows from $\mathrm{observersourcefluxzero}(Q, t)$. □

Theorem 4.9. *For every natural number Q and natural number t and natural number n,* $\mathrm{observersourceflux}(Q, t, n{+}1) = \mathrm{wittxor}(\mathrm{wittlinear}(\mathrm{observersourceflux}(Q, t, n)), \mathrm{wittshl}(\mathrm{ob}$ $n), n + 1))$.

Proof. The goal follows from $\mathrm{observersourcefluxstep}(Q, t, n)$. □

Theorem 4.10. *For every natural number Q and natural number t and natural number n and natural number s,* $\mathrm{observersourceflux}(Q, t, n + s) = \mathrm{wittxor}(\mathrm{wittlinpow}(s, \mathrm{observersourceflux}(Q, t, n)), \mathrm{wittshl}(\mathrm{observersourceflux}(Q, t{+}$ $n, s), n))$.

Proof. The goal follows from $\mathrm{observersourcefluxappend}(Q, t, n, s)$. □

Theorem 4.11. *For every natural number Q and natural number t,* $\mathrm{observerqrank}(Q, t) = t + Q - 2 \cdot \mathrm{qorder}(\mathrm{observerdefect}(Q, t))$.

Proof. The goal follows from observerqrankdefinition(Q, t). □

Theorem 4.12. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then* wittshl(observerdefect$(m \cdot P, t+1), 1) =$ wittxor(wittlinear(observerdefect$(m \cdot P, t)),$ wittshl(observersource$(m \cdot P, t), 1)$).

Proof. The goal follows from periodicobserverpolynomialstep(T, P, m, t). □

Theorem 4.13. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then* observersource$(m \cdot P, t) =$ wittxor(wittxor(wittcarry(observercode$(t + m \cdot P)),$ wittcarry(observercode$(t))),$ naturalbit(leftdefect$(m \cdot P, t, 1))$).

Proof. The goal follows from periodicobserversource(T, P, m, t). □

Theorem 4.14. *For every natural number T and natural number P and natural number m and natural number t and natural number n, if* centerperiodic(T, P)*, then if $T \leq t$, then* wittshl(observerdefect$(m \cdot P, t+n), n) =$ wittxor(wittlinpow$(n,$ observerdefect$(m \cdot P, t)),$ observersourceflux$(m \cdot P, t, n))$.

Proof. The goal follows from periodicobserverpolynomialwindow(T, P, m, t, n). □

Theorem 4.15. *For every natural number T and natural number P and natural number m and natural number t and natural number n and natural number r and natural number k, if* centerperiodic(T, P)*, then if $T \leq t$, then if* qdivides$(r,$ observerdefect$(m \cdot P, t))$*, then if $k \leq n+r$, then* qdivides$(k,$ observerdefect$(m \cdot P, t + n))$ *if and only if* qdivides$(k,$ observersourceflux$(m \cdot P, t, n))$.

Proof. The goal follows from periodicobserverwindowdivisibility(T, P, m, t, n, r, k). □

Theorem 4.16. *For every natural number T and natural number P and natural number m and natural number t and natural number r, if* centerperiodic(T, P)*, then if $T \leq t$, then if* qdivides$(r,$ observerdefect$(m \cdot P, t))$*, then* qdivides$(r + 1,$ observerdefect$(m \cdot P, t+1))$ *if and only if* qdivides$(r+1,$ observersource$(m \cdot P, t))$.

Proof. The goal follows from periodicobserverqgain(T, P, m, t, r). □

Theorem 4.17. *For every natural number Q and natural number t, if $0 < Q$, then $2 \cdot$* qorder(observerdefect$(Q, t)) \leq t + Q$.

Proof. The goal follows from observerqorderbound(Q, t). □

Theorem 4.18. *For every natural number T and natural number P and natural number m and natural number t and natural number n, if $0 < P$, then if $0 < m$, then if* centerperiodic(T, P)*, then if $T \leq t$, then if $0 < n$, then* observerqrank$(m \cdot P, t + n) <$ observerqrank$(m \cdot P, t)$ *if and only if* qdivides(qorder(observerdefect$(m \cdot P, t)) + n/2 + 1,$ observersourceflux$(m \cdot P, t, n))$.

Proof. The goal follows from periodicobserverqrankdrop(T, P, m, t, n). □

Theorem 4.19. observerdefect$(5, 2) = 216$ *and* observerdefect$(5, 3) = 412 \wedge$ qdivides$(1,$ observerdefect$(5, 2)) \wedge \neg$ qdivides$(1,$ observerdefect$(5, 3)) \wedge$ center$(2) =$ false $\wedge$ center$(7) =$ false $\wedge$ center$(3) =$ true $\wedge$ center$(8) =$ true.

Proof. The goal follows from observerseedqloss. □

Theorem 4.20. *For every natural number r and natural number a and natural number d, if* wittlinpow$(r, a) = d$*, then $a \leq d$.*

Proof. The goal follows from qquotientbound(r, a, d). □

Theorem 4.21. *For every natural number d and natural number r, if* qdivides$(r + 1, d)$*, then* qordersearch$(d, r + 1) = r + 1$.

Proof. The goal follows from qordersearchhit(d, r). □

Theorem 4.22. *For every natural number d and natural number r, if not* qdivides$(r + 1, d)$*, then* qordersearch$(d, r + 1) =$ qordersearch(d, r).

Proof. The goal follows from qordersearchmiss(d, r). □

Theorem 4.23. *For every natural number d,* qorder$(d) =$ qordersearch$(d,$ binarydegree$(d))$.

Proof. The goal follows from qorderdefinition(d). □

Theorem 4.24. *For every natural number r and natural number d, if* qdivides(r, d)*, then there exists a natural number a such that $a \leq d$ and* wittlinpow$(r, a) = d$.

Proof. The goal follows from qdivisesboundedwitness(r, d). □

Theorem 4.25. *For every natural number d,* qordersearch$(d, 0) = 0$.

Proof. The goal follows from qordersearchzero(d). □

Theorem 4.26. *For every natural number a and natural number y,* wittlinpow(twopow(a), y) = wittxor(wittxor(y, wittshl(y, twopow(a))), wittshl(y, twopow(a)+twopow(a))).

Proof. The goal follows from qfrobeniustwopow(a, y). □

Theorem 4.27. wittlinpow(16, 1) = wittxor(wittxor(1, wittshl(1, 16)), wittshl(1, 32)) *and* wittlinpow(16, wittxor(1, wittshl(1, 16))) = wittxor(1, wittshl(1, 48)).

Proof. The goal follows from qsixteenpage. □

Theorem 4.28. *For every natural number a and natural number y,* qdivides(twopow(a), wittxor(y, witts twopow(a)))).

Proof. The goal follows from qdividespageshifttwopow(a, y). □

Theorem 4.29. *For every natural number a and natural number n and natural number H,* qdivides(twopow(a), wittxor(H, wittxor(pagefold(3·twopow(a), H, n), pagetail(3· twopow(a), H, n)))).

Proof. The goal follows from qdividespagefoldtwopow(a, n, H). □

Theorem 4.30. *For every natural number L and natural number H and natural number p, if $0 < L$, then* page(L, wittcarry(H), p) = wittxor(wittcarry(page(L, H, p)), naturalbit(pa

Proof. The goal follows from pagecarry(L, H, p). □

Theorem 4.31. *For every natural number L and natural number H and natural number n, if $0 < L$, then* pagefold(L, wittcarry(H), n) = carryledger(L, H, n).

Proof. The goal follows from carryfoldeqledger(L, H, n). □

Theorem 4.32. *For every natural number a and natural number H and natural number n,* qdivides(twopow(a), wittxor(wittcarry(H), wittxor(carryledger(3· twopow(a), H, n), pagetail(3 · twopow(a), wittcarry(H), n)))).

Proof. The goal follows from carryledgercongruencetwopow(a, H, n). □

Theorem 4.33. *For every natural number T and natural number P and natural number m and natural number t and natural number a, if* centerperiodic(T, P), *then if $T \leq t$, then* qdivides(twopow(a), wittxor(observersource(m·P, t), wittxor(wittxor(carryledger(3· twopow(a), observercode(t+m·P), t+m·P+1), carryledger(3·twopow(a), observercode(t), t+ $m \cdot P + 1$)), naturalbit(leftdefect($m \cdot P$, t, 1))))).

Proof. The goal follows from periodicobserversourceledger(T, P, m, t, a). □

Theorem 4.34. *For every natural number T and natural number P and natural number m and natural number t and natural number r and natural number a, if* centerperiodic(T, P)*, then if* $T \leq t$*, then if* $r+1 \leq$ twopow(a)*, then if* qdivides$(r,$ observerdefect$(m \cdot P, t))$*, then* qdivides$(r+1,$ observerdefect$(m \cdot P, t+1))$ *if and only if* qdivides$(r+1,$ wittxor$($wittxor$($carryledger$(3 \cdot$twopow$(a),$ observercode$(t+ m \cdot P), t + m \cdot P + 1),$ carryledger$(3 \cdot$ twopow$(a),$ observercode$(t), t + m \cdot P + 1)),$ naturalbit$($leftdefect$(m \cdot P, t, 1))))$.

Proof. The goal follows from periodicobserverqgainledger(T, P, m, t, r, a). □

Theorem 4.35. qdivides$(16,$ wittlinpow$(16, 1))$ *and* wittxor$($wittphi$($wittxor$(2,$ wittlinpow$(16, 1))),$ witt wittxor$($wittlinpow$(17, 1), 4) \wedge \neg$ qdivides$(16,$ wittxor$($wittphi$($wittxor$(2,$ wittlinpow$(16, 1))),$ wittphi$(2)))$

Proof. The goal follows from pageresiduenotclosed. □

Theorem 4.36. *For every natural number a and natural number $A0$ and natural number $A1$ and natural number $A2$,* qdivides$($twopow$(a),$ wittxor$($wittxor$($wittxor$(A0,$ wittshl$(A1$ twopow$(a))),$ wittxor$($wittxor$(A0, A2),$ wittshl$($wittxor$(A1, A2),$ twopow$(a)))))$.

Proof. The goal follows from qblockresidue$(a, A0, A1, A2)$. □

Theorem 4.37. *For every natural number a and natural number $A0$ and natural number $A1$ and natural number $A2$, if $A0 <$* twopow$($twopow$(a))$*, then if $A1 <$* twopow$($twopow$(a))$*, then if $A2 <$* twopow$($twopow$(a))$*, then* qdivides$($twopow$(a),$ wittxor$($wittxor$(A0,$ wittshl$(A1,$ twopow$(a))),$ wittshl$(A2,$ twopow$(a)+$ twopow$(a)))$ *if and only if $A0 = A2$ and $A1 = A2$.*

Proof. The goal follows from qblockchargezeroiff$(a, A0, A1, A2)$. □

Theorem 4.38. *For every natural number a and natural number A, if $A <$* twopow$($twopow$(a))$*, then* wittcarry$($wittxor$($wittxor$(A,$ wittshl$(A,$ twopow$(a))),$ wittshl$(A,$ twopow$(a)+$ twopow$(a))))$ = wittxor$($wittxor$($wittxor$($wittcarry$(A),$ wittshl$($wittcarry$(A),$ twopow$(a))),$ wittshl$($witt twopow$(a))),$ wittxor$($wittshl$($naturalbit$($blockendpoint$($twopow$(a), A)),$ twopow$(a)),$ wittshl$($naturalbit twopow$(a)))))$.

Proof. The goal follows from repeatedblockcarry(a, A). □

Theorem 4.39. *For every natural number a and natural number A, if $A <$* twopow$($twopow$(a))$*, then* qdivides$($twopow$(a),$ wittxor$($wittcarry$($wittxor$($wittxor$(A,$ wittshl$(A,$ twopow twopow$(a)))),$ naturalbit$($blockendpoint$($twopow$(a), A))))$.

Proof. The goal follows from repeatedblockcarrycongruence(a, A). □

Theorem 4.40. *For every natural number a and natural number H and natural number p and natural number A, if $A <$ twopow(twopow(a)), then if* page(3·twopow(a), H, p) = wittxor(wittxor(A, wittshl(A, twopow(a))), wittshl(A, twopow(a)+ twopow(a))), *then* qdivides(twopow(a), wittxor(page(3·twopow(a), wittcarry(H), p), wittxor(naturalbit twopow(a), H, p))))).

Proof. The goal follows from pagecarryrepeated(a, H, p, A). □

Theorem 4.41. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P), *then if $T \leq t$, then* observersource($m \cdot P$, t) = wittxor(wittxor(wittcarry(observerdefect($m \cdot P$, t)), crosscarry(observercode(t), observerdefect($m \cdot P$, t))), naturalbit(leftdefect($m \cdot P$, t, 1))).

Proof. The goal follows from periodicobserversourcecrosscarry(T, P, m, t). □

Theorem 4.42. *There exists a natural number s such that there exists a natural number $s2$ such that* qdivides(16, wittxor(s, $s2$)) *and not* qdivides(16, wittxor(wittphi(s), wittphi($s2$)

Proof. The goal follows from residuenotcongruence. □

Theorem 4.43. *For every natural number Q and natural number t and natural number j,* rowbit(observerdefect(Q, t), j) = leftdefect(Q, t, j).

Proof. The goal follows from observerdefectleftreadout(Q, t, j). □

Theorem 4.44. *For every natural number r and natural number e and natural number k, if $e \neq 0$, then if* qdivides(r, wittshl(e, k)), *then $k + 2 \cdot r \leq$* binarydegree(wittshl(e, k)).

Proof. The goal follows from qfrontdegreebudget(r, e, k). □

Theorem 4.45. *For every natural number n and natural number e and natural number o, if $e \neq 0$, then if* twopow(n) $\leq o$, *then if* wittshl(e, o) $<$ twopow(3 · twopow(n)), *then not* qdivides(twopow(n), wittshl(e, o)).

Proof. The goal follows from qpageoffsetcharge(n, e, o). □

Theorem 4.46. *For every natural number Q and natural number t and natural number k, if $0 < Q$, then if* rightmostleft(Q, t, k), *then $k + 2 \cdot$* qorder(observerdefect(Q, t)) $\leq t + Q$.

Proof. The goal follows from observerfrontqbudget(Q, t, k). □

Theorem 4.47. *For every natural number Q and natural number t and natural number k, if $0 < Q$, then if* rightmostleft(Q, t, k)*, then* $k \leq$ observerqrank(Q, t).

Proof. The goal follows from observerfrontqrank(Q, t, k). □

Theorem 4.48. *For every natural number a and natural number Q and natural number t and natural number p and natural number o, if* twopow$(a) \leq o$*, then if $o < 3 \cdot$* twopow(a)*, then if* rightmostleft$(Q, t, 3 \cdot$ twopow$(a) \cdot p + o)$*, then not* qdivides(twopow(a), page$(3 \cdot$ twopow(a), observerdefect$(Q, t), p))$.

Proof. The goal follows from observerfrontpagecharge(a, Q, t, p, o). □

Theorem 4.49. *For every natural number Q and natural number t, if* leftdefect$(Q, t, 0) =$ false*, then* leftdefect$(Q, t + 1, 0) =$ bxor(leftdefect$(Q, t, 1)$, centergate$(Q, t))$.

Proof. The goal follows from centerdefectcoupling(Q, t). □

Theorem 4.50. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then* leftdefect$(m \cdot P, t, 1) =$ centergate$(m \cdot P, t)$.

Proof. The goal follows from periodiccentercoupling(T, P, m, t). □

Theorem 4.51. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then if* center$(t) =$ true*, then* leftdefect$(m \cdot P, t, 1) =$ false.

Proof. The goal follows from periodicblacknoarrival(T, P, m, t). □

Theorem 4.52. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then if* center$(t) =$ false*, then* leftdefect$(m \cdot P, t, 1) =$ rightdefect$(m \cdot P, t, 1)$.

Proof. The goal follows from periodicwhitemirror(T, P, m, t). □

Theorem 4.53. *For every natural number T and natural number P, if* center$(T + P) =$ center(T)*, then if for every natural number t, if $T \leq t$, then* leftdefect$(P, t, 1) =$ centergate(P, t)*, then* centerperiodic(T, P).

Proof. The goal follows from centerperiodicofcoupling(T, P). □

Theorem 4.54. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if $T \leq t$, then* observersource$(m \cdot P, t) =$ wittxor(wittxor(wittcarry(observercode$(t + m \cdot P))$, wittcarry(observercode$(t))$), naturalbit(centergate$(m \cdot P, t))$).

Proof. The goal follows from periodicobserversourcecentergate(T, P, m, t). □

Theorem 4.55. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if* $T \leq t$*, then if* center$(t) =$ true*, then* observersource$(m \cdot P, t) =$ wittxor(wittcarry(observercode$(t + m \cdot P)$), wittcarry(observercode(t))).

Proof. The goal follows from periodicblackobserversource(T, P, m, t). □

Theorem 4.56. *For every natural number T and natural number P and natural number m and natural number t, if* centerperiodic(T, P)*, then if* $T \leq t$*, then if* center$(t) =$ false*, then* observersource$(m \cdot P, t) =$ wittxor(wittxor(wittcarry(observercode$(t + m \cdot P)$), wittcarry(observercode(t))), naturalbit(rightdefect$(m \cdot P, t, 1)$)).

Proof. The goal follows from periodicwhiteobserversource(T, P, m, t). □

Theorem 4.57. *For every natural number T and natural number P and natural number m and natural number b and natural number k and natural number s, if* centerperiodic(T, P)*, then if* $T \leq b$*, then if* rightmostleft$(m \cdot P, b, k)$*, then if* $s + 1 < k$*, then* leftdefect$(m \cdot P, b + s, 1) =$ false *and* observersource$(m \cdot P, b + s) =$ wittxor(wittcarry(observercode$(b + s + m \cdot P)$), wittcarry(observercode$(b + s)$)).

Proof. The goal follows from periodicfrontgapobserversource(T, P, m, b, k, s). □

Theorem 4.58. *For every natural number T and natural number P and natural number m and natural number b and natural number k, if* centerperiodic(T, P)*, then if* $T \leq b$*, then if* rightmostleft$(m \cdot P, b, k)$*, then* leftdefect$(m \cdot P, b + (k - 1), 1) =$ true *and* center$(b+(k-1)) =$ false $\wedge$ rightdefect$(m \cdot P, b+(k-1), 1) =$ true $\wedge$ observersource$(m \cdot P, b+(k-1)) =$ wittxor(wittxor(wittcarry(observercode$(b + (k-1) + m \cdot P)$), wittcarry(observercode$(b + (k-1))$)), 1).

Proof. The goal follows from periodicfrontarrivalobserversource(T, P, m, b, k). □

Theorem 4.59. *For every natural number Q and natural number start, if* center$(start) =$ false*, then if* center$(start + Q) =$ false*, then if* center$(start + 1) =$ false*, then if* center$(start + 1 + Q) =$ false*, then if* rightcell$(start, 1) =$ false*, then if* rightcell$(start + Q, 1) =$ false*, then* leftgatephase$(Q, start + 1, 0) =$ rightphase$(Q, start, 2)$ *and* leftgatephase$(Q, start+1, 0) =$ centersign$(start, 1) \cdot$ rightphase$(Q, start, 2)$.

Proof. The goal follows from coupledarrivalgatephase$(Q, start)$. □

Theorem 4.60. *For every natural number Q and natural number t, if $0 < Q$, then* rowbit(aligneddefect$(Q, t), 0) =$ true.

Proof. The goal follows from aligneddefectbitzero(Q, t). □

Theorem 4.61. *For every natural number T and natural number P and natural number m and natural number b and natural number k, if* centerperiodic(T, P)*, then if $T \leq b$, then if* rightmostleft$(m \cdot P, b, k)$*, then* leftdefect$(m \cdot P, b + (k - 1), 1) =$ true *and* centergate$(m \cdot P, b + (k - 1)) =$ true $\wedge$ leftdefect$(m \cdot P, b + (k - 1) + 1, 0) =$ false.

Proof. The goal follows from periodicarrivalnocrossing(T, P, m, b, k). □

Theorem 4.62. *For every natural number Q and natural number t and natural number r, if* leftadjacentdefect$(Q, t) =$ true*, then if* centercol$(t+r) =$ true*, then if* centercol$(t + r + Q) =$ true*, then* nearcollision$(Q, t, r + 1, 2 \cdot (r + 1))$.

Proof. The goal follows from unitdefectforcesnearcollisionkit(Q, t, r). □

Theorem 4.63. *For every natural number t,* center$(t) =$ centercol(t).

Proof. The goal follows from centereqkit(t). □

Theorem 4.64. *For every natural number k,* $0 <$ twopow(k).

Proof. The goal follows from twopowpos(k). □

Theorem 4.65. *For every natural number b and natural number d,* defectstep$(b, d) =$ wittxor(wittphi(wittxor(b, d)), wittphi(b)).

Proof. The goal follows from defectstepeq(b, d). □

Theorem 4.66. *For every natural number y and natural number Q,* wittphi(wittshl$(y, Q)) =$ wittshl(wittphi$(y), Q)$.

Proof. The goal follows from phishl(y, Q). □

Theorem 4.67. *For every natural number Q and natural number t,* wittxor(alignedbackground(Q, t), ali wittrow$(t + Q)$.

Proof. The goal follows from reconstruct(Q, t). □

Theorem 4.68. *For every natural number T and natural number P, if $T \leq 512$, then if $0 < P$, then if $P \leq 128$, then there exists a natural number t such that $T \leq t$ and $t < T + 32 \wedge \neg \mathrm{center}(t + P) = \mathrm{center}(t)$.*

Proof. The goal follows from horizonthirtytwo(T, P). □

Theorem 4.69. *For every natural number T and natural number P, if $T \leq 512$, then if $0 < P$, then if $P \leq 128$, then not* centerperiodic(T, P).

Proof. The goal follows from nocenterperiod512128(T, P). □

Theorem 4.70. *For every natural number T and natural number P and natural number m and natural number t and natural number k, if* centerperiodic(T, P), *then if $T \leq t$, then if* leftdefect$(m \cdot P, t, k) = \mathrm{true}$, *then there exists a natural number s such that $t \leq s$ and $s < t + k \wedge$* rightdefect$(m \cdot P, s, 1) = \mathrm{true}$.

Proof. The goal follows from leftdefectsourced(T, P, m, t, k). □

Theorem 4.71. *For every natural number T and natural number P and natural number m and natural number t and natural number k, if* centerperiodic(T, P), *then if $T \leq t$, then if* rightagrees$(m \cdot P, t, k)$, *then* leftdefect$(m \cdot P, t, k) = \mathrm{false}$.

Proof. The goal follows from leftconeofrightagrees(T, P, m, t, k). □

Theorem 4.72. *For every natural number Q and natural number t and integer p, if* defect$(Q, t + 1, p) = \mathrm{false}$, *then if* defect$(Q, t, p) = \mathrm{false}$, *then if* defect$(Q, t, p + 1) = \mathrm{false}$, *then* defect$(Q, t, p - 1) = \mathrm{false}$.

Proof. The goal follows from defectleftofagree(Q, t, p). □

Theorem 4.73. shortperiodsfail$(512, 128, 32) = \mathrm{true}$.

Proof. The goal follows from certificate51212832. □

Theorem 4.74. *For every natural number $T0$ and natural number $P0$ and natural number H and natural number T and natural number P, if* shortperiodsfail$(T0, P0, H) =$ true, *then if $T \leq T0$, then if $0 < P$, then if $P \leq P0$, then* centerdisagreeswithin$(T, P, H) =$ true.

Proof. The goal follows from checkerofcertificate$(T0, P0, H, T, P)$. □

Theorem 4.75. *For every natural number T and natural number P and natural number H, if* centerdisagreeswithin$(T, P, H) = \mathrm{true}$, *then there exists a natural number t such that $T \leq t$ and $t < T + H \wedge \neg \mathrm{center}(t + P) = \mathrm{center}(t)$.*

Proof. The goal follows from disagreementofchecker(T, P, H). □

Theorem 4.76. *For every natural number T and natural number P and natural number H, if* centerdisagreeswithin$(T, P, H) =$ true*, then not* centerperiodic(T, P).

Proof. The goal follows from notcenterperiodicofchecker(T, P, H). □

Theorem 4.77. *For every natural number Q and natural number t,* rightdefect$(Q, t, 1) =$ defect$(Q, t, 1)$.

Proof. The goal follows from rightdefecteqdefect(Q, t). □

Theorem 4.78. *For every natural number t,* center$(t) =$ not(ledgerdiagonal(t)).

Proof. The goal follows from centereqnotledgerdiagonal(t). □

Theorem 4.79. *For every natural number T and natural number P,* centerperiodic(T, P) *if and only if* ledgerperiodic(T, P).

Proof. The goal follows from centerperiodiciffledgerperiodic(T, P). □

Theorem 4.80. *For every natural number t,* wittrow$(t) =$ wittxor(wittlinpow$(t, 1)$, wittwin$(0, t)$).

Proof. The goal follows from roweqlinearxorledger(t). □

Theorem 4.81. *For every natural number y,* wittphi$(y) =$ wittxor(wittlinear(y), wittshl(wittcarry(y), 1

Proof. The goal follows from phieqlinearxorcarry(y). □

Theorem 4.82. *For every natural number a and natural number b,* wittlinear(wittxor$(a, b)) =$ wittxor(wittlinear(a), wittlinear(b)).

Proof. The goal follows from philinearxor(a, b). □

Theorem 4.83. *For every natural number m,* trailingparity$(2 \cdot m) =$ false.

Proof. The goal follows from trailingparityeven(m). □

Theorem 4.84. *For every natural number m,* trailingparity$(2 \cdot m + 1) =$ not(trailingparity(m)).

Proof. The goal follows from trailingparityodd(m). □

Theorem 4.85. *For every natural number t,* rowbit(philinpow$(1, t, 1), t) =$ true.

Proof. The goal follows from linearcenter(t). □

Theorem 4.86. *For every natural number t,* $\mathrm{rowbit}(\mathrm{philinpow}(1, t, 1), t + 1) = \mathrm{trailingparity}(t)$.

Proof. The goal follows from $\mathrm{linearabovecenter}(t)$. □

Theorem 4.87. *For every natural number t, if $1 \leq t$, then* $\mathrm{rowbit}(\mathrm{philinpow}(1, t, 1), t - 1) = \mathrm{trailingparity}(t)$.

Proof. The goal follows from $\mathrm{linearbelowcenter}(t)$. □

Theorem 4.88. *For every natural number s and natural number y,* $\mathrm{philin}(s, \mathrm{philin}(s, y)) = \mathrm{philin}(2 \cdot s, y)$.

Proof. The goal follows from $\mathrm{philinphilin}(s, y)$. □

Theorem 4.89. *For every natural number m and natural number y,* $\mathrm{philinpow}(1, 2 \cdot m, y) = \mathrm{philinpow}(2, m, y)$.

Proof. The goal follows from $\mathrm{philinpowtwomul}(m, y)$. □

Theorem 4.90. *For every natural number m and natural number y,* $\mathrm{wittlinpow}(m, y) = \mathrm{philinpow}(1, m, y)$.

Proof. The goal follows from $\mathrm{philinearpoweqphilinpow}(m, y)$. □

Theorem 4.91. *For every natural number t, if $1 \leq t$, then* $\mathrm{center}(t + 1) = \mathrm{rule}(\mathrm{bxor}(\mathrm{trailingparity}(t), \mathrm{ledgerbit}(t, t+1)), \mathrm{not}(\mathrm{ledgerbit}(t, t)), \mathrm{bxor}(\mathrm{trailingparity}(t), \mathrm{ledgerbit}(t, t - 1)))$.

Proof. The goal follows from $\mathrm{centersuccledgerpaper}(t)$. □

Theorem 4.92. *For every natural number j and natural number t,* $\mathrm{rdiag}(j, t) = \mathrm{bxor}(\mathrm{rowbit}(\mathrm{philinpow}(1, t, 1), j), \mathrm{ledgerbit}(t, j))$.

Proof. The goal follows from $\mathrm{rdiageqlinearbxorledger}(j, t)$. □

Theorem 4.93. *For every natural number t,* $\mathrm{cell}(t, -1) = \mathrm{bxor}(\mathrm{trailingparity}(t), \mathrm{ledgerbit}(t, t + 1))$.

Proof. The goal follows from $\mathrm{celllefteqpaper}(t)$. □

Theorem 4.94. *For every natural number t, if $1 \leq t$, then* $\mathrm{cell}(t, 1) = \mathrm{bxor}(\mathrm{trailingparity}(t), \mathrm{ledgerbit}(t, t - 1))$.

Proof. The goal follows from $\mathrm{cellrighteqpaper}(t)$. □

Theorem 4.95. *For every natural number T and natural number P, if $0 < P$, then if* centerperiodic(T, P)*, then* centerkernelfinite*.*

Proof. The goal follows from centerkernelfiniteofcenterperiodic(T, P). □

Theorem 4.96. *For every natural number T and natural number P, if* centerinfinitekernel*, then if $0 < P$, then not* centerperiodic(T, P)*.*

Proof. The goal follows from notcenterperiodicofinfinitekernel(T, P). □

Theorem 4.97. *If* centerinfinitekernel*, then* problemone*.*

Proof. Assume h. The goal follows from problem1ofinfinitekernel(h). □

Theorem 4.98. *For every natural number k and natural number r and natural number t,* kernelseq$(k, r, t{+}1) =$ bxor(bxor(kernelseq(k, r, t), cell(twopow$(k)\cdot t + r$, negpow(k))), response(twopow$(k) \cdot t + r, k)$)*.*

Proof. The goal follows from kernelseqstep(k, r, t). □

Theorem 4.99. *For every natural number t and natural number k, if $t <$* twopow(k)*, then* response$(t, k) =$ bxor(center(t), center$(t +$ twopow$(k))$)*.*

Proof. The goal follows from responsecone(t, k). □

Theorem 4.100. *For every natural number k and natural number r and natural number t,* kernelseq$(k + 1, r, t) =$ kernelseq$(k, r, 2 \cdot t)$*.*

Proof. The goal follows from kernelseqeven(k, r, t). □

Theorem 4.101. *For every natural number k and natural number r and natural number t,* kernelseq$(k{+}1, r{+}$twopow$(k), t) =$ kernelseq$(k, r, 2{\cdot}t{+}1)$*.*

Proof. The goal follows from kernelseqodd(k, r, t). □

Theorem 4.102. *For every natural number k and natural number r and natural number t, if* twopow$(k) \leq$ twopow$(k){\cdot}t{+}r$*, then* kernelseq$(k, r, t{+}1) =$ bxor(kernelseq(k, r, t), bxor(bxor(cell(twopow$(k){\cdot}t{+}r$, negpow(k))), rdiag(twopow$(k)\cdot t + r -$ twopow(k), twopow$(k) \cdot t + r)$), wittresp(twopow$(k) \cdot t + r, k)$))*.*

Proof. The goal follows from kernelseqsteplate(k, r, t). □

Theorem 4.103. *For every natural number d and natural number k, if $d \leq k$, then* center$(d) =$ bxor(wittresp$(d +$ twopow$(k), k)$, response$(d +$ twopow$(k), k)$)*.*

Proof. The goal follows from centerrecoveredatscale(d, k). □

Theorem 4.104. *If* nodyadicselfsimilarity*, then* centerinfinitekernel*.*

Proof. Assume h. The goal follows from centerinfinitekernelofnodyadicselfsimilarity(h). □

Theorem 4.105. *If* nodyadicselfsimilarity*, then* problemone*.*

Proof. Assume h. The goal follows from problem1ofnodyadicselfsimilarity(h). □

Theorem 4.106. *For every natural number* k*,* center(twopow($k + 1$)) = bxor(center(twopow(k)), wittresp(twopow(k), k))*.*

Proof. The goal follows from centertwopowsucc(k). □

Theorem 4.107. *For every natural number* a *and natural number* b*,* twopow($a + b$) = twopow(a) $\cdot$ twopow(b)*.*

Proof. The goal follows from twopowadd(a, b). □

Theorem 4.108. *If* somedilationaperiodic*, then* centerinfinitekernel*.*

Proof. Assume h. The goal follows from centerinfinitekernelofdilationaperiodic(h). □

Theorem 4.109. *If* somedilationaperiodic*, then* problemone*.*

Proof. Assume h. The goal follows from problem1ofdilationaperiodic(h). □

Theorem 4.110. *If* dyadiccolumnaperiodic*, then* problemone*.*

Proof. Assume h. The goal follows from problem1ofdyadiccolumnaperiodic(h). □

Theorem 4.111. *For every natural number* b *and natural number* a *and natural number* c*,* bpow($b, a + c$) = bpow(b, a) $\cdot$ bpow(b, c)*.*

Proof. The goal follows from bpowadd(b, a, c). □

Theorem 4.112. *For every natural number* b*, if* $2 \leq b$*, then if* centerinfinitekernelin(b)*, then* problemone*.*

Proof. The goal follows from problem1ofinfinitekernelin(b). □

Theorem 4.113. *For every natural number b and natural number k, if $2 \leq b$, then $k + 1 \leq \mathrm{bpow}(b, k)$.*

Proof. The goal follows from $\mathrm{bpowgesuccpaper}(b, k)$. □

Theorem 4.114. *For every natural number b and natural number T and natural number P, if $2 \leq b$, then if $0 < P$, then if* $\mathrm{centerperiodic}(T, P)$*, then* $\mathrm{centerkernelfinitein}(b)$.

Proof. The goal follows from $\mathrm{centerkernelfiniteinofcenterperiodicpaper}(b, T, P)$. □

Theorem 4.115. *For every natural number b and natural number t, if $2 \leq b$, then if* $\mathrm{dilationaperiodicin}(b, t)$*, then* $\mathrm{centerinfinitekernelin}(b)$.

Proof. The goal follows from $\mathrm{centerinfinitekernelinofdilationaperiodicpaper}(b, t)$. □

Theorem 4.116. *For every natural number b and natural number t, if $2 \leq b$, then if* $\mathrm{dilationaperiodicin}(b, t)$*, then* problemone.

Proof. The goal follows from $\mathrm{problem1ofdilationaperiodicinpaper}(b, t)$. □

Theorem 4.117. *For every natural number K,* $\mathrm{kappacollapse}(K)$ *if and only if* $\mathrm{centerperiodic}(0, \mathrm{twopow}(K))$.

Proof. The goal follows from $\mathrm{kappacollapseiff}(K)$. □

Theorem 4.118. *For every natural number K, if* $\mathrm{kappacollapse}(K)$*, then* $\mathrm{centerperiodic}(0, \mathrm{twopow}(K))$.

Proof. The goal follows from $\mathrm{centerperiodicofkappacollapse}(K)$. □

Theorem 4.119. *For every natural number K, if* $\mathrm{centerperiodic}(0, \mathrm{twopow}(K))$*, then* $\mathrm{kappacollapse}(K)$.

Proof. The goal follows from $\mathrm{kappacollapseofcenterperiodic}(K)$. □

Theorem 4.120. *For every natural number K, if $K \leq 7$, then not* $\mathrm{kappacollapse}(K)$.

Proof. The goal follows from $\mathrm{nokappacollapsesmall}(K)$. □

Theorem 4.121. *For every natural number K, if* problemone*, then not* $\mathrm{kappacollapse}(K)$.

Proof. The goal follows from $\mathrm{nokappacollapseofproblem1paper}(K)$. □

Theorem 4.122. *For every natural number j,* $\mathrm{center}(\mathrm{twopow}(j+1)) = \mathrm{bxor}(\mathrm{not}(\mathrm{center}(\mathrm{twopow}(j))), \mathrm{response}(\mathrm{twopow}(j), j))$.

Proof. The goal follows from $\mathrm{dyadiccolumnstep}(j)$. □

Theorem 4.123. *For every natural number K and natural number d, if* $\mathrm{dilationperiodic}(1, K, d)$, *then* $\mathrm{dyadicresponseperiodic}(K, d)$.

Proof. The goal follows from $\mathrm{dyadicresponseperiodicofdilationperiodic}(K, d)$. □

Theorem 4.124. *If* dyadicresponseaperiodic, *then* dyadiccolumnaperiodic.

Proof. Assume h. The goal follows from $\mathrm{dyadiccolumnaperiodicofresponse}(h)$. □

Theorem 4.125. *If* dyadicresponseaperiodic, *then* problemone.

Proof. Assume h. The goal follows from $\mathrm{problem1ofdyadicresponseaperiodic}(h)$. □

Theorem 4.126. *For every natural number j,* $\mathrm{response}(\mathrm{twopow}(j+1), j+1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{window}(\mathrm{twopow}(j+1), \mathrm{twopow}(j), 0), \mathrm{window}(\mathrm{twopow}(j+1), \mathrm{twopow}(j), \mathrm{negpow}(j))), \mathrm{window}(\mathrm{twopow}(j+1)+\mathrm{twopow}(j), \mathrm{twopow}(j), 0))$.

Proof. The goal follows from $\mathrm{dyadicresponsehalvespaper}(j)$. □

Theorem 4.127. *For every natural number K and natural number d, if* $\mathrm{dyadicresponseperiodic}(K, d)$, *then* $\mathrm{dilationperiodic}(1, K, 2 \cdot d)$.

Proof. The goal follows from $\mathrm{dilationperiodicofdyadicresponseperiodic}(K, d)$. □

Theorem 4.128. *If* dyadiccolumnaperiodic, *then* dyadicresponseaperiodic.

Proof. Assume h. The goal follows from $\mathrm{dyadicresponseaperiodicofcolumn}(h)$. □

Theorem 4.129. dyadiccolumnaperiodic *if and only if* dyadicresponseaperiodic.

Proof. The goal follows from dyadiccolumnaperiodiciffresponse. □

Theorem 4.130. *For every truth value a and truth value b and truth value c,* $\mathrm{rule}(a, b, c) = \mathrm{wolfram}(30, a, b, c)$.

Proof. The goal follows from $\mathrm{ruleeqwolframthirty}(a, b, c)$. □

Theorem 4.131. *For every truth value a and truth value b and truth value c,* $\mathrm{rule}(a, b, c) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{bxor}(a, b), c), b \,\mathrm{and}\, c)$.

Proof. The goal follows from $\mathrm{ruleeqlinearxorcarry}(a, b, c)$. □

Theorem 4.132. *For every natural number t,* $\mathrm{wittcarry}(\mathrm{wittrow}(t)) = \mathrm{wittxor}(\mathrm{edgesource}(t), \mathrm{interior}(t$

Proof. The goal follows from $\mathrm{carrysplit}(t)$. □

Theorem 4.133. *For every natural number t,* $\mathrm{rowbit}(\mathrm{wittcarry}(\mathrm{wittrow}(t)), 1) = \mathrm{rdiag}(1, t)$.

Proof. The goal follows from $\mathrm{carryrowrightpair}(t)$. □

Theorem 4.134. *For every natural number t,* $\mathrm{wittwin}(0, t) = \mathrm{wittxor}(\mathrm{edgeledger}(t), \mathrm{interiorledger}(t))$.

Proof. The goal follows from $\mathrm{ledgersplit}(t)$. □

Theorem 4.135. *For every natural number t,* $\mathrm{center}(t) = \mathrm{not}(\mathrm{bxor}(\mathrm{edgediagonal}(t), \mathrm{interiordiagonal}(t)$

Proof. The goal follows from $\mathrm{centereqnotbxordiagonals}(t)$. □

Theorem 4.136. *For every natural number n and natural number d,* $\mathrm{rowbit}(\mathrm{philinpow}(1, n, 1), d) = \mathrm{trinomialparity}(n, d)$.

Proof. The goal follows from $\mathrm{testbitphilinpoweqtrinomialparity}(n, d)$. □

Theorem 4.137. *For every natural number t,* $\mathrm{edgediagonal}(t) = \mathrm{edgeformula}(t, t)$.

Proof. The goal follows from $\mathrm{edgediagonaleqformula}(t)$. □

Theorem 4.138. *For every natural number t,* $\mathrm{center}(t) = \mathrm{not}(\mathrm{bxor}(\mathrm{edgeformula}(t, t), \mathrm{interiordiagonal}(t$

Proof. The goal follows from $\mathrm{centereqnotformulaxorinterior}(t)$. □

Theorem 4.139. *For every natural number t,* $\mathrm{interiordiagonal}(t) = \mathrm{interiorformula}(t, t)$.

Proof. The goal follows from $\mathrm{interiordiagonaleqformula}(t)$. □

Theorem 4.140. *For every natural number t,* $\mathrm{center}(t) = \mathrm{not}(\mathrm{bxor}(\mathrm{edgeformula}(t, t), \mathrm{interiorformula}(t,$

Proof. The goal follows from $\mathrm{centereqnotformulas}(t)$. □

Theorem 4.141. *For every natural number T and natural number P,* $\mathrm{centerperiodic}(T, P)$ *if and only if* $\mathrm{interioragainstedge}(T, P)$.

Proof. The goal follows from $\mathrm{centerperiodiciffinterioragainstedge}(T, P)$. □

Theorem 4.142. *For every natural number Q and natural number z and natural number t, if* zerodiagonal(z)*, then if* settledby$(z+2, t)$*, then if* $z+3 \leq t$*, then* diagonalphase$(Q, z+3, t+1)$ + diagonalflux$(Q, z+3, t+1, 1)$ = diagonalphase$(Q, z+1, t)$ + diagonalflux$(Q, z+1, t, 1)$.

Proof. The goal follows from diagonalresetsuminvariant(Q, z, t). □

Theorem 4.143. *For every natural number n and natural number d, if* $2 \cdot n < d$*, then* trinomialparity(n, d) = false.

Proof. The goal follows from trinomialparityeqfalseofgt(n, d). □

Theorem 4.144. *For every natural number t,* edgediagonal(t) = edgeformula$(t, t/2+1)$.

Proof. The goal follows from edgediagonalhalf(t). □

Theorem 4.145. *For every natural number T and natural number P and natural number t, if* centerperiodic(T, P)*, then if* $T \leq t$*, then if* center(t) = true*, then* bxor(ledgerbit$(t+P, t+P+1)$, ledgerbit$(t, t+1)$) = bxor(trailingparity$(t+P)$, trailingparity(t)).

Proof. The goal follows from periodicblackledgerabsorbs(T, P, t). □

Theorem 4.146. *For every natural number n and natural number d, if* $d \leq 2 \cdot n$*, then* trinomialparity(n, d) = trinomialparity$(n, 2 \cdot n - d)$.

Proof. The goal follows from trinomialparitypalindrome(n, d). □

Theorem 4.147. *For every natural number n,* trinomialparity(n, n) = true.

Proof. The goal follows from trinomialparitycentral(n). □

Theorem 4.148. *For every natural number m and natural number e,* trinomialparity$(2 \cdot m, 2 \cdot e)$ = trinomialparity(m, e).

Proof. The goal follows from trinomialparityeveneven(m, e). □

Theorem 4.149. *For every natural number m and natural number e,* trinomialparity$(2 \cdot m, 2 \cdot e + 1)$ = false.

Proof. The goal follows from trinomialparityevenodd(m, e). □

Theorem 4.150. *For every natural number m and natural number e,* trinomialparity$(2 \cdot m + 1, 2 \cdot e + 1)$ = trinomialparity(m, e).

Proof. The goal follows from trinomialparityoddodd(m, e). □

Theorem 4.151. *For every natural number m and natural number e,* trinomialparity($2 \cdot m + 1, 2 \cdot e + 2$) = bxor(trinomialparity($m, e + 1$), trinomialparity(m, e)).

Proof. The goal follows from trinomialparityoddevensucc(m, e). □

Theorem 4.152. columnparity(0) = true.

Proof. The goal follows from columnparityzero. □

Theorem 4.153. *For every natural number e,* columnparity($2 \cdot e + 1$) = columnparity(e).

Proof. The goal follows from columnparityodd(e). □

Theorem 4.154. *For every natural number e,* columnparity($2 \cdot e + 2$) = not(columnparity(e)).

Proof. The goal follows from columnparityevensucc(e). □

Theorem 4.155. *For every natural number N,* sameparitycolumn(N) = columnparity($N/2$).

Proof. The goal follows from sameparitycolumneq(N). □

Theorem 4.156. *For every natural number t,* edgediagonal(t) = timeparity(t).

Proof. The goal follows from edgediagonaleqtimeparity(t). □

Theorem 4.157. *For every natural number t,* center(t) = not(bxor(timeparity(t), interiordiagonal(t))).

Proof. The goal follows from centereqnottimeparityxorinterior(t). □

Theorem 4.158. *For every natural number t,* interiordiagonal(t) = bxor(not(center(t)), timeparity(t)).

Proof. The goal follows from interiordiagonaleqcenter(t). □

Theorem 4.159. *For every natural number t and natural number P,* timeparity($t + 2 \cdot P$) = timeparity(t).

Proof. The goal follows from timeparityaddtwomul(t, P). □

Theorem 4.160. *For every natural number T and natural number P, if* centerperiodic(T, P)*, then* interiorperiodic($T, 2 \cdot P$).

Proof. The goal follows from interiorperiodicofcenterperiodic(T, P). □

Theorem 4.161. *For every natural number T and natural number P, if* interiorperiodic(T, P)*, then* centerperiodic$(T, 2 \cdot P)$.

Proof. The goal follows from centerperiodicofinteriorperiodic(T, P). □

Theorem 4.162. problemone *if and only if* interioraperiodic.

Proof. The goal follows from problem1iffinterioraperiodic. □

Theorem 4.163. *If for every natural number T and natural number P, if* $0 < P$*, then not* interiorperiodic(T, P)*, then* problemone.

Proof. Assume h. The goal follows from problem1ofinterioraperiodic(h). □

Theorem 4.164. centerkernelfinite *if and only if* interiorkernelfinite.

Proof. The goal follows from centerkernelfiniteiffinterior. □

Theorem 4.165. centerinfinitekernel *if and only if not* interiorkernelfinite.

Proof. The goal follows from centerinfinitekerneliffinterior. □

Theorem 4.166. *If not* interiorkernelfinite*, then* problemone.

Proof. Assume h. The goal follows from problem1ofinteriorinfinitekernel(h). □

Theorem 4.167. *For every natural number n,* trinomialparity$(n, 0) =$ true.

Proof. The goal follows from trinomialparityzeroright(n). □

Theorem 4.168. *For every natural number m,* wittcarry(philinpow$(1, 2 \cdot m, 1)) = 0$.

Proof. The goal follows from linearselfintersectioneven(m). □

Theorem 4.169. *For every natural number m and natural number i,* rowbit(wittcarry(philinpow$(1, 2 \cdot m{+}1, 1)), 2 \cdot (i{+}1){+}1) =$ trinomialparity$(m, i{+}1)$ and not(trinomialparity(m, i)).

Proof. The goal follows from linearselfintersectionoddrise(m, i). □

Theorem 4.170. *For every natural number m and natural number i,* rowbit(wittcarry(philinpow$(1, 2 \cdot m{+}1, 1)), 2 \cdot i{+}2) =$ trinomialparity(m, i) and not(trinomialparity$(m, i{+}1)$).

Proof. The goal follows from linearselfintersectionoddfall(m, i). □

Theorem 4.171. *For every natural number m,* rowbit(wittcarry(philinpow$(1, 2 \cdot m + 1, 1)), 1) =$ true.

Proof. The goal follows from linearselfintersectionoddone(m). □

Theorem 4.172. *For every natural number m,* interior$(m) =$ wittxor(lowsource(m), interiorthree(m)).

Proof. The goal follows from interiorsplitthree(m). □

Theorem 4.173. *For every natural number t,* interiorledger$(t) =$ wittxor(lowledger(t), interiorthreeledg

Proof. The goal follows from interiorledgersplitthree(t). □

Theorem 4.174. *For every natural number t,* interiordiagonal$(t) =$ bxor(lowdiagonal(t), interiorthreedi

Proof. The goal follows from interiordiagonalsplitthree(t). □

Theorem 4.175. *For every natural number e,* bxor(columnparity$(e+1)$, columnparity$(e)) =$ not(trailingparity$(e + 1)$).

Proof. The goal follows from columnparitysuccbxor(e). □

Theorem 4.176. *For every natural number s,* lowdiagonal$(s + 2 + 1) =$ not(bxor(columnparity(s), columnparity$(s/2))$).

Proof. The goal follows from lowdiagonalthree(s). □

Theorem 4.177. *For every natural number t, if $2 \leq t$, then* bxor(timeparity(t), lowdiagonal$(t)) =$ not(trailingparity$(t - 2)$).

Proof. The goal follows from lowdiagonaleq(t). □

Theorem 4.178. *For every natural number t, if $2 \leq t$, then* center$(t) =$ bxor(trailingparity$(t - 2)$, interiorthreediagonal(t)).

Proof. The goal follows from centereqtrailingparitybxorthree(t). □

Theorem 4.179. *For every natural number T and natural number P, if $2 \leq T$, then* centerperiodic(T, P) *if and only if* threeagainsttrailing(T, P).

Proof. The goal follows from centerperiodiciffthreeagainsttrailing(T, P). □

Theorem 4.180. problemone *if and only if* threeaperiodic.

Proof. The goal follows from problem1iffthreeaperiodic. □

Theorem 4.181. *If* threeaperiodic*, then* problemone.

Proof. Assume h. The goal follows from problem1ofthreeaperiodic(h). □

Theorem 4.182. *For every natural number T and natural number P, if $0 < P$, then not* trailingparityperiodic(T, P).

Proof. The goal follows from trailingparitynotperiodic(T, P). □

Theorem 4.183. *For every natural number T and natural number P, if $0 < P$, then if* centerperiodic(T, P)*, then not* threeperiodic(T, P).

Proof. The goal follows from interiorthreeaperiodicofcenterperiodic(T, P). □

Theorem 4.184. *For every natural number k and natural number s, if $s + 1 <$* twopow(k)*, then* trailingparity(twopow(k) + s) = trailingparity(s).

Proof. The goal follows from trailingparitytwopowadd(k, s). □

Theorem 4.185. *For every natural number j and natural number m and natural number d, if $m <$* twopow(j)*, then if $d <$* twopow(j)*, then* trinomialparity(twopow(j+ 1) − 1 − m, d) = trinomialparity(twopow(j) − 1 − m, d).

Proof. The goal follows from trinomialparitydyadicstable(j, m, d). □

Theorem 4.186. *For every natural number j and natural number m and natural number d, if $m <$* twopow(j)*, then if $d <$* twopow(j)*, then* trinomialparity(twopow(j)− 1 − m, d) = stablekernel(m, d).

Proof. The goal follows from trinomialparityeqstablekernel(j, m, d). □

Theorem 4.187. *For every natural number m and natural number d,* stablekernel(m, d+ 2) = bxor(bxor(stablekernel(m+1, d+2), stablekernel(m+1, d+1)), stablekernel(m+ 1, d)).

Proof. The goal follows from stablekernelsucc(m, d). □

Theorem 4.188. *For every natural number m,* stablekernel(m, 0) = stablekernel(m+ 1, 0).

Proof. The goal follows from stablekernelsucczero(m). □

Theorem 4.189. *For every natural number m,* stablekernel(m, 1) = bxor(stablekernel(m+ 1, 1), stablekernel($m + 1$, 0)).

Proof. The goal follows from stablekernelsuccone(m). □

Theorem 4.190. *For every natural number n and natural number d,* trinomialparity(n+1, d+2) = bxor(bxor(trinomialparity(n, d+2), trinomialparity(n, d+1)), trinomialparity(n, d)).

Proof. The goal follows from trinomialparityrowsucc(n, d). □

Theorem 4.191. *For every natural number j, if* $1 \leq j$, *then* timeparity(twopow(j)) = false.

Proof. The goal follows from timeparitytwopow(j). □

Theorem 4.192. *For every natural number j and natural number n and natural number y and natural number t, if* $n <$ twopow(j), *then if* $t <$ twopow(j), *then* conv(twopow(j) − 1 − n, y, t) = stableconv(n, y, t).

Proof. The goal follows from conveqstableconv(j, n, y, t). □

Theorem 4.193. *For every natural number j, if* $1 \leq j$, *then* center(twopow(j)) = not(stableformula(j, twopow(j))).

Proof. The goal follows from centertwopoweqstableformula(j). □

Theorem 4.194. dyadiccolumnaperiodic *if and only if* stableformulaaperiodic.

Proof. The goal follows from dyadiccolumnaperiodiciffstable. □

Theorem 4.195. *If* stableformulaaperiodic, *then* problemone.

Proof. Assume h. The goal follows from problem1ofstableformulaaperiodic(h). □

Theorem 4.196. *For every natural number S and natural number P and natural number Q, if* $0 < P$, *then if* $0 < Q$, *then not* trailingdiffperiodic(S, P, Q).

Proof. The goal follows from trailingdiffnotperiodic(S, P, Q). □

Theorem 4.197. *For every natural number T and natural number P and natural number S and natural number Q, if* $0 < P$, *then if* $0 < Q$, *then if* centerperiodic(T, P), *then not* threeperiodic(S, Q).

Proof. The goal follows from notthreeperiodicofcenterperiodic(T, P, S, Q). □

Theorem 4.198. *For every natural number T and natural number P and natural number k, if* centerperiodic(T, P), *then* centerperiodic(T, twopow(k)·P).

Proof. The goal follows from centerperiodictwopowmul(T, P, k). □

Theorem 4.199. *For every natural number T and natural number P and natural number k, if* centerperiodic(T, P)*, then* threeoffclassperiodic$(k, T + 2, P)$.

Proof. The goal follows from threeoffclassofcenterperiodic(T, P, k). □

Theorem 4.200. *For every natural number T and natural number P and natural number k, if* centerperiodic(T, P)*, then* threeclassagainsttrailing$(k, T+1, P)$.

Proof. The goal follows from threeclassofcenterperiodic(T, P, k). □

Theorem 4.201. *For every natural number a and natural number b,* wittcarry(wittxor(a, b)) = wittxor(wittxor(wittcarry(a), wittcarry(b)), cross(a, b)).

Proof. The goal follows from carryxor(a, b). □

Theorem 4.202. *For every natural number m,* wittcarry(wittrow(m)) = wittxor(wittxor(wittcarry(wittlinpow$(m, 1)$), wittcarry(wittwin$(0, m)$))), cross(wittlinpow$(m, 1)$, wittwir

Proof. The goal follows from carryrowsplit(m). □

Theorem 4.203. *For every natural number j and natural number m, if $m <$* twopow(j)*, then* rowterm(twopow$(j + 1), m)$ = rowterm(twopow$(j), m)$.

Proof. The goal follows from rowtermtwopowsucc(j, m). □

Theorem 4.204. *For every natural number j and natural number m, if $m <$* twopow(j)*, then* rowterm(twopow$(j), m)$ = dyadicrow(m).

Proof. The goal follows from rowtermtwopoweqdyadicrow(j, m). □

Theorem 4.205. *For every natural number j, if $1 \leq j$, then* center(twopow$(j+1))$ = bxor(center(twopow(j)), blockparity(j)).

Proof. The goal follows from centertwopowsucceq(j). □

Theorem 4.206. *For every natural number j, if $1 \leq j$, then* wittresp(twopow$(j), j)$ = blockparity(j).

Proof. The goal follows from wittrespbooltwopoweqblockparity(j). □

Theorem 4.207. *For every natural number j, if $1 \leq j$, then* response(twopow$(j), j)$ = not(blockparity(j)).

Proof. The goal follows from responsetwopoweqnotblockparity(j). □

Theorem 4.208. *For every natural number K and natural number d, if* dilationperiodic$(1, K, d)$*, then* blockparityperiodic$(K + 1, d)$*.*

Proof. The goal follows from blockparityperiodicofdilationperiodic(K, d). □

Theorem 4.209. *If* blockparityaperiodic*, then* dyadiccolumnaperiodic*.*

Proof. Assume h. The goal follows from dyadiccolumnaperiodicofblockparityaperiodic(h). □

Theorem 4.210. *If* blockparityaperiodic*, then* problemone*.*

Proof. Assume h. The goal follows from problem1ofblockparityaperiodic(h). □

Theorem 4.211. *For every natural number t, if $2 \leq t$, then* ldiag$(2, t) =$ false*.*

Proof. The goal follows from ldiagtwoeq(t). □

Theorem 4.212. *For every natural number t, if $2 \leq t$, then* ldiag$(3, t) =$ timeparity(t)*.*

Proof. The goal follows from ldiagthreeeq(t). □

Theorem 4.213. *For every natural number t, if $2 \leq t$, then* ldiag$(4, t) =$ true*.*

Proof. The goal follows from ldiagfoureq(t). □

Theorem 4.214. *For every natural number t, if $3 \leq t$, then* ldiag$(5, t) =$ timeparity(t)*.*

Proof. The goal follows from ldiagfiveeq(t). □

Theorem 4.215. *For every natural number t, if $3 \leq t$, then* ldiag$(6, t) =$ timeparity(t)*.*

Proof. The goal follows from ldiagsixeq(t). □

Theorem 4.216. *For every natural number t,* ldiag$(7, t) =$ false*.*

Proof. The goal follows from ldiagseveneq(t). □

Theorem 4.217. *For every natural number n,* trinomialparity$(n + 2, n) =$ trailingparity$((n + 2)/2)$.

Proof. The goal follows from trinomialparitybelowtwo(n). □

Theorem 4.218. *For every natural number t,* interiorthreeledger$(t) =$ wittxor(leftbandledger(t), interio

Proof. The goal follows from interiorthreeledgerspliteight(t). □

Theorem 4.219. *For every natural number t, if $6 \leq t$, then* leftbanddiagonal$(t) =$ bxor(not(timeparity(t)), bxor(trailingparity$(t-4)$, trailingparity$((t-4)/2)$)).

Proof. The goal follows from leftbanddiagonaleq(t). □

Theorem 4.220. *For every natural number t, if $6 \leq t$, then* center$(t) =$ bxor(trailingparity$(t-2)$, bxor(bxor(not(timeparity(t)), bxor(trailingparity$(t-4)$, trailingparity$((t-4)/2)$)), interioreightdiagonal(t))).

Proof. The goal follows from centereqleftband(t). □

Theorem 4.221. *For every natural number t, if $6 \leq t$, then* center$(t) =$ bxor(rulerbackground(t), interioreightdiagonal(t)).

Proof. The goal follows from centereqrulerbackgroundbxoreight(t). □

Theorem 4.222. *For every natural number T and natural number P, if $6 \leq T$, then* centerperiodic(T, P) *if and only if* eightagainstrulers(T, P).

Proof. The goal follows from centerperiodiciffeightagainstrulers(T, P). □

Theorem 4.223. problemone *if and only if* eightaperiodic.

Proof. The goal follows from problem1iffeightaperiodic. □

Theorem 4.224. *If* eightaperiodic*, then* problemone.

Proof. Assume h. The goal follows from problem1ofeightaperiodic(h). □

Theorem 4.225. *For every natural number k and natural number r and natural number u and natural number m, if $m <$* twopow(k)*, then if $r <$* twopow(k)*, then if $1 \leq u$, then* rowterm(twopow$(k) \cdot u + r, m) =$ bxor(frozen(k, r, m), trailingparity$(u-1)$ and second(k, r, m)).

Proof. The goal follows from rowtermkernellaw(k, r, u, m). □

Theorem 4.226. *For every natural number k and natural number r and natural number u, if $r < \mathrm{twopow}(k)$, then if $1 \leq u$, then* $\mathrm{interiorformula}(\mathrm{twopow}(k) \cdot u + r, \mathrm{twopow}(k) \cdot u + r) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{frozenprefix}(k, r), \mathrm{trailingparity}(u - 1) \mathrm{\ and\ } \mathrm{secondprefix}(k, r)), \mathrm{latetail}(k, r, u))$.

Proof. The goal follows from $\mathrm{interiorformulakernel}(k, r, u)$. □

Theorem 4.227. *For every natural number k and natural number r and natural number u, if $r < \mathrm{twopow}(k)$, then if $1 \leq u$, then if $1 \leq k$, then* $\mathrm{kernelseq}(k, r, u) = \mathrm{not}(\mathrm{bxor}(\mathrm{timeparity}(r), \mathrm{bxor}(\mathrm{bxor}(\mathrm{frozenprefix}(k, r), \mathrm{trailingparity}(u - 1) \mathrm{\ and\ } \mathrm{secondprefix}(k, r)), \mathrm{latetail}(k, r, u))))$.

Proof. The goal follows from $\mathrm{kernelseqexplicit}(k, r, u)$. □

Theorem 4.228. *For every natural number k and natural number u and natural number m, if $m < \mathrm{twopow}(k)$, then if $1 \leq u$, then* $\mathrm{rowterm}(\mathrm{twopow}(k) \cdot u, m) = \mathrm{dyadicrow}(m)$.

Proof. The goal follows from $\mathrm{rowtermtwopowmul}(k, u, m)$. □

Theorem 4.229. *For every natural number k and natural number u, if $1 \leq k$, then if $1 \leq u$, then* $\mathrm{center}(\mathrm{twopow}(k) \cdot u) = \mathrm{bxor}(\mathrm{center}(\mathrm{twopow}(k)), \mathrm{latetail}(k, 0, u))$.

Proof. The goal follows from $\mathrm{centertwopowmul}(k, u)$. □

Theorem 4.230. *For every natural number k and natural number u, if $1 \leq k$, then if $1 \leq u$, then* $\mathrm{center}(\mathrm{twopow}(k) \cdot u + 1) = \mathrm{bxor}(\mathrm{center}(\mathrm{twopow}(k) + 1), \mathrm{bxor}(\mathrm{latetail}(k, 1, 1), \mathrm{latetail}(k, 1, u)))$.

Proof. The goal follows from $\mathrm{centertwopowmuladdone}(k, u)$. □

Theorem 4.231. *For every natural number s,* $\mathrm{trinomialparity}(s, s + 1) = \mathrm{trailingparity}(s)$.

Proof. The goal follows from $\mathrm{trinomialparitysuccself}(s)$. □

Theorem 4.232. *For every natural number s, if $1 \leq s$, then* $\mathrm{trinomialparity}(s, s - 1) = \mathrm{trailingparity}(s)$.

Proof. The goal follows from $\mathrm{trinomialparitypredself}(s)$. □

Theorem 4.233. *For every natural number k and natural number u and natural number r, if $1 \leq k$, then* $\mathrm{timeparity}(\mathrm{twopow}(k) \cdot u + r) = \mathrm{timeparity}(r)$.

Proof. The goal follows from $\mathrm{timeparitytwopowmuladd}(k, u, r)$. □

Theorem 4.234. *For every natural number k and natural number m, if $1 \leq k$, then if $m <$ twopow(k), then* second$(k, 1, m) =$ false.

Proof. The goal follows from secondone(k, m). □

Theorem 4.235. *For every natural number k, if $1 \leq k$, then* secondprefix$(k, 1) =$ false.

Proof. The goal follows from secondprefixone(k). □

Theorem 4.236. rulercoefficient$(1) =$ true.

Proof. The goal follows from rulercoefficientone. □

Theorem 4.237. rulercoefficient$(2) =$ true.

Proof. The goal follows from rulercoefficienttwo. □

Theorem 4.238. rulercoefficient$(3) =$ false.

Proof. The goal follows from rulercoefficientthree. □

Theorem 4.239. *For every natural number r, if $4 \leq r$, then* rulercoefficient$(r) =$ false.

Proof. The goal follows from rulercoefficientofgefour(r). □

Theorem 4.240. *For every natural number k and natural number r and natural number u, if $3 \leq k$, then if $r <$ twopow(k), then if $1 \leq u$, then* rulerbackground(twopow$(k) \cdot u + r) =$ bxor(frozenruler(k, r), trailingparity$(u - 1)$ and rulercoefficient(r)).

Proof. The goal follows from rulerbackgroundkernel(k, r, u). □

Theorem 4.241. *For every natural number T and natural number P and natural number k and natural number r, if* centerperiodic(T, P), *then if $3 \leq k$, then if $r <$* twopow(k), *then if $r = 1$ or $r = 2$, then* eightclassagainstruler$(k, r, T + 1, P)$.

Proof. The goal follows from eightclassofcenterperiodic(T, P, k, r). □

Theorem 4.242. *For every natural number T and natural number P and natural number k and natural number r, if* centerperiodic(T, P), *then if $3 \leq k$, then if $r <$* twopow(k), *then if $r \neq 1$, then if $r \neq 2$, then* eightclassperiodic$(k, r, T + 1, P)$.

Proof. The goal follows from eightoffclassofcenterperiodic(T, P, k, r). □

Theorem 4.243. *For every natural number T and natural number P and natural number k and natural number r and natural number u, if* centerperiodic(T, P), *then if* $1 \leq k$, *then if* $r <$ twopow(k), *then if* $1 \leq u$, *then if* $T \leq$ twopow$(k) \cdot u+r$, *then* bxor(latetail$(k, r, u+P)$, latetail(k, r, u)) = bxor(trailingparity$(u+P-1)$, trailingparity$(u-1)$) and secondprefix(k, r).

Proof. The goal follows from latetailobligation(T, P, k, r, u). □

Theorem 4.244. *For every natural number k and natural number v, if* $1 \leq k$, *then if* $1 \leq v$, *then* latetail$(k, 0, 2 \cdot v)$ = bxor(latetail$(k+1, 0, v)$, blockparity(k)).

Proof. The goal follows from latetailscalezero(k, v). □

Theorem 4.245. *For every natural number k and natural number r and natural number v, if* $1 \leq k$, *then if* $r <$ twopow(k), *then if* $1 \leq v$, *then* bxor(latetail$(k+1, r, v)$, latetail$(k, r, 2 \cdot v)$) = bxor(bxor(frozenprefix$(k+1, r)$, frozenprefix(k, r)), bxor(trailingparity$(v-1)$ and secondprefix$(k+1, r)$, not(trailingparity$(v-1)$) and secondprefix(k, r))).

Proof. The goal follows from latetailscale(k, r, v). □

Theorem 4.246. *For every natural number k and natural number m, if* $m <$ twopow(k), *then* second$(k, 2, m)$ = false.

Proof. The goal follows from secondtwo(k, m). □

Theorem 4.247. *For every natural number k,* secondprefix$(k, 2)$ = false.

Proof. The goal follows from secondprefixtwo(k). □

Theorem 4.248. *For every natural number k and natural number m, if* $m <$ twopow(k), *then* second$(k, 3, m)$ = pairrow(m).

Proof. The goal follows from secondthree(k, m). □

Theorem 4.249. *For every natural number k, if* $2 \leq k$, *then* secondprefix$(k, 3)$ = true.

Proof. The goal follows from secondprefixthree(k). □

Theorem 4.250. *For every natural number k,* latetail$(k, 1, 1)$ = rowterm(twopow$(k)+1$, twopow(k)).

Proof. The goal follows from latetailoneatone(k). □

Theorem 4.251. *For every natural number k, if $1 \leq k$, then* rowterm(twopow(k)+ 1, twopow(k)) = center(twopow(k)) and cell(twopow(k), 1).

Proof. The goal follows from rowtermtwopowaddone(k). □

Theorem 4.252. *For every natural number k, if $1 \leq k$, then* latetail($k, 1, 1$) = center(twopow(k)) and cell(twopow(k), 1).

Proof. The goal follows from latetailoneatonecell(k). □

Theorem 4.253. *For every natural number k and natural number v, if $2 \leq k$, then if $1 \leq v$, then* latetail($k + 1, 3, v$) = bxor(latetail($k, 3, 2 \cdot v$), bxor(bxor(frozenprefix($k + 1, 3$), frozenprefix($k, 3$)), true)).

Proof. The goal follows from latetailthreescale(k, v). □

Theorem 4.254. *For every natural number T and natural number P and natural number k, if* centerperiodic(T, P)*, then if $2 \leq k$, then if $T \leq$* twopow(k)+ 3*, then* latetail($k, 3, 1 + P$) = bxor(latetail($k, 3, 1$), trailingparity(P)).

Proof. The goal follows from latetailthreeobligationone(T, P, k). □

Theorem 4.255. *For every natural number K, if $4 \leq K$, then* rowterm($K+$ $3, K + 2$) = cell($K + 2, 0$) and cell($K + 2, 1$).

Proof. The goal follows from rowtermthreerowtwo(K). □

Theorem 4.256. *For every natural number K, if $4 \leq K$, then* rowterm($K+$ $3, K+1$) = bxor(bxor(cell($K+1, -1$) and cell($K+1, 0$), cell($K+1, 0$) and cell($K+$ $1, 1$)), cell($K + 1, 1$) and cell($K + 1, 2$)).

Proof. The goal follows from rowtermthreerowone(K). □

Theorem 4.257. *For every natural number K, if $4 \leq K$, then* rowterm($K+$ $3, K$) = bxor(bxor(cell($K, -2$) and cell($K, -1$), cell($K, 0$) and cell($K, 1$)), cell($K, 2$) and cell($K, 3$)).

Proof. The goal follows from rowtermthreerowzero(K). □

Theorem 4.258. *For every natural number k, if $2 \leq k$, then* latetail($k, 3, 1$) = bxor(bxor(bxor(bxor(cell(twopow(k), -2) and cell(twopow(k), -1), cell(twopow(k), 0) and cell(twopow(k)+ $1, -1$) and cell(twopow(k)+$1, 0$), cell(twopow(k)+$1, 0$) and cell(twopow(k)+ $1, 1$)), cell(twopow(k) + $1, 1$) and cell(twopow(k) + $1, 2$))), cell(twopow(k) + $2, 0$) and cell(twopow(k) + $2, 1$)).

Proof. The goal follows from latetailthreeatonecells(k). □

Theorem 4.259. *For every natural number T and natural number P and natural number k, if* centerperiodic(T, P)*, then if* $2 \leq k$*, then if* $T \leq$ twopow$(k)+3$*, then* bxor(latetail$(k, 3, 1+P)$, bxor(bxor(bxor(bxor(cell(twopow$(k), -2)$ and cell(twopow$(k), -1)$, cell($1, -1)$ and cell(twopow$(k)+1, 0)$, cell(twopow$(k)+1, 0)$ and cell(twopow$(k)+1, 1)$), cell(twopow$(k) + 1, 1)$ and cell(twopow$(k) + 1, 2)$))), cell(twopow$(k) + 2, 0)$ and cell(twopow$(k) + 2, 1)$))) = trailingparity(P).

Proof. The goal follows from threerowsagainstruler(T, P, k). □

Theorem 4.260. *For every natural number t,* center$(t) =$ rdiag(t, t).

Proof. The goal follows from centereqrdiagself(t). □

Theorem 4.261. *For every natural number T and natural number P,* centerperiodic(T, P) *if and only if* readoutperiodic(T, P).

Proof. The goal follows from centerperiodiciffcenterper(T, P). □

Theorem 4.262. problemone *if and only if* antidiagonalaperiodic.

Proof. The goal follows from problem1iffantidiagonal. □

Theorem 4.263. *If* antidiagonalaperiodic*, then* problemone.

Proof. Assume h. The goal follows from problem1ofantidiagonal(h). □

Theorem 4.264. *For every natural number j and natural number t,* rdiag$(j, t) =$ towerword(j, t).

Proof. The goal follows from rdiageqtowerword(j, t). □

Theorem 4.265. *For every natural number t,* center$(t) =$ towerword(t, t).

Proof. The goal follows from centereqtowerword(t). □

Theorem 4.266. *For every natural number k and natural number s,* towerword$(k+2, s+1) =$ bxor(towerword$(k+2, s)$, towerword$(k+1, s)$ or towerword(k, s)).

Proof. The goal follows from towerwordstep(k, s). □

Theorem 4.267. *For every natural number s,* towerword$(0, s) =$ true.

Proof. The goal follows from towerwordzero(s). □

Theorem 4.268. *For every natural number s,* towerword$(1, s) =$ timeparity(s).

Proof. The goal follows from towerwordone(s). □

Theorem 4.269. *For every natural number j,* $\mathrm{towerword}(j+1,0) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{towerwordsucczero}(j)$. □

Theorem 4.270. *For every natural number j and natural number q and natural number t,* $\mathrm{rdiag}(j, t + \mathrm{twopow}(j) \cdot q) = \mathrm{rdiag}(j,t)$.

Proof. The goal follows from $\mathrm{rdiagaddtwopowmul}(j,q,t)$. □

Theorem 4.271. *For every natural number k and natural number s,* $\mathrm{orword}(k, s + \mathrm{twopow}(k+1)) = \mathrm{orword}(k,s)$.

Proof. The goal follows from $\mathrm{orwordperiodic}(k,s)$. □

Theorem 4.272. *For every natural number k and natural number s,* $\mathrm{towerword}(k+2, s + \mathrm{twopow}(k+1)) = \mathrm{bxor}(\mathrm{towerword}(k+2,s), \mathrm{wordweight}(k))$.

Proof. The goal follows from $\mathrm{towerwordhalfperiod}(k,s)$. □

Theorem 4.273. $\mathrm{wordweight}(0) = \mathrm{false}$.

Proof. The goal follows from wordweightzero. □

Theorem 4.274. $\mathrm{wordweight}(1) = \mathrm{false}$.

Proof. The goal follows from wordweightone. □

Theorem 4.275. $\mathrm{wordweight}(2) = \mathrm{false}$.

Proof. The goal follows from wordweighttwo. □

Theorem 4.276. $\mathrm{wordweight}(3) = \mathrm{false}$.

Proof. The goal follows from wordweightthree. □

Theorem 4.277. $\mathrm{wordweight}(4) = \mathrm{false}$.

Proof. The goal follows from wordweightfour. □

Theorem 4.278. $\mathrm{wordweight}(5) = \mathrm{false}$.

Proof. The goal follows from wordweightfive. □

Theorem 4.279. *For every natural number j and natural number s,* $\mathrm{towerword}(j, s + \mathrm{twopow}(j)) = \mathrm{towerword}(j,s)$.

Proof. The goal follows from $\mathrm{towerwordaddtwopow}(j,s)$. □

Theorem 4.280. halfperiodic(2).

Proof. The goal follows from halfperiodictwo. □

Theorem 4.281. halfperiodic(3).

Proof. The goal follows from halfperiodicthree. □

Theorem 4.282. *For every natural number k, if* halfperiodic$(k+1)$*, then if* halfperiodic$(k+2)$*, then* wordweight$(k+1) =$ false.

Proof. The goal follows from wordweightsuccfalse(k). □

Theorem 4.283. *For every natural number k, if* halfperiodic$(k+1)$*, then if* halfperiodic$(k+2)$*, then* halfperiodic$(k+3)$.

Proof. The goal follows from halfperiodicsucc(k). □

Theorem 4.284. *For every natural number j, if $2 \leq j$, then* halfperiodic(j).

Proof. The goal follows from halfperiodicoftwole(j). □

Theorem 4.285. *For every natural number j and natural number t, if $2 \leq j$, then* rdiag$(j, t + \text{twopow}(j-1)) =$ rdiag(j, t).

Proof. The goal follows from rdiagperiodhalf(j, t). □

Theorem 4.286. *For every natural number k,* wordweight$(k) =$ false.

Proof. The goal follows from wordweightfalse(k). □

Theorem 4.287. *For every natural number k and natural number s,* towerword$(k+2, s + \text{twopow}(k+1)) =$ towerword$(k+2, s)$.

Proof. The goal follows from towerwordaddtwopowhalf(k, s). □

Theorem 4.288. *For every natural number k and natural number i,* orword$(k, i) =$ bxor(towerword$(k+1, i)$, exposed(k, i)).

Proof. The goal follows from orwordeqexposed(k, i). □

Theorem 4.289. *For every natural number k and natural number s, if $2 \leq k$, then* exposed$(k, s + \text{twopow}(k)) =$ exposed(k, s).

Proof. The goal follows from exposedperiodic(k, s). □

Theorem 4.290. *For every natural number n and natural number s,* $\mathrm{lineartower}(n, s) = \mathrm{constprefix}(n + 1, s)$.

Proof. The goal follows from $\mathrm{lineartowereqconstprefix}(n, s)$. □

Theorem 4.291. *For every natural number n and natural number s, if $s < n$, then* $\mathrm{constprefix}(n, s) = \mathrm{false}$.

Proof. The goal follows from $\mathrm{constprefixbelow}(n, s)$. □

Theorem 4.292. *For every natural number n,* $\mathrm{constprefix}(n, n) = \mathrm{true}$.

Proof. The goal follows from $\mathrm{constprefixdiag}(n)$. □

Theorem 4.293. *For every natural number t, if $1 \leq t$, then* $\mathrm{lineartower}(t - 1, t) = \mathrm{true}$.

Proof. The goal follows from $\mathrm{lineartowerdiag}(t)$. □

Theorem 4.294. *For every natural number t, if $2 \leq t$, then* $\mathrm{center}(t) = \mathrm{not}(\mathrm{masktransport}(t - 2, t))$.

Proof. The goal follows from $\mathrm{centereqnotmasktransport}(t)$. □

Theorem 4.295. *For every natural number T and natural number P and natural number t, if* $\mathrm{centerperiodic}(T, P)$*, then if $T \leq t$, then if $2 \leq t$, then* $\mathrm{masktransport}(t + P - 2, t + P) = \mathrm{masktransport}(t - 2, t)$.

Proof. The goal follows from $\mathrm{masktransportcenterperiodic}(T, P, t)$. □

Theorem 4.296. *For every natural number k and natural number s,* $\mathrm{towerword}(k + 2, s) = \mathrm{bxor}(\mathrm{prefixword}(k + 1, s), \mathrm{exposedprefix}(k, s))$.

Proof. The goal follows from $\mathrm{towerwordsplit}(k, s)$. □

Theorem 4.297. *For every natural number t, if $2 \leq t$, then* $\mathrm{center}(t) = \mathrm{bxor}(\mathrm{prefixword}(t - 1, t), \mathrm{exposedprefix}(t - 2, t))$.

Proof. The goal follows from $\mathrm{centersplit}(t)$. □

Theorem 4.298. *For every natural number k and natural number s,* $\mathrm{towerword}(k + 2, s) = \mathrm{bxor}(\mathrm{lineartower}(k + 1, s), \mathrm{masktransport}(k, s))$.

Proof. The goal follows from $\mathrm{towerwordunfolded}(k, s)$. □

Theorem 4.299. *For every natural number t, if $2 \leq t$, then* $\mathrm{center}(t) = \mathrm{bxor}(\mathrm{lineartower}(t - 1, t), \mathrm{masktransport}(t - 2, t))$.

Proof. The goal follows from centertransport(t). □

Theorem 4.300. *For every natural number n,* pascalparity$(n, 0) =$ true.

Proof. The goal follows from pascalparityzeroright(n). □

Theorem 4.301. *For every natural number a and natural number b,* pascalparity$(2 \cdot a, 2 \cdot b) =$ pascalparity(a, b).

Proof. The goal follows from pascalparityeveneven(a, b). □

Theorem 4.302. *For every natural number a and natural number b,* pascalparity$(2 \cdot a, 2 \cdot b + 1) =$ false.

Proof. The goal follows from pascalparityevenodd(a, b). □

Theorem 4.303. *For every natural number a and natural number b,* pascalparity$(2 \cdot a + 1, 2 \cdot b) =$ pascalparity(a, b).

Proof. The goal follows from pascalparityoddeven(a, b). □

Theorem 4.304. *For every natural number a and natural number b,* pascalparity$(2 \cdot a + 1, 2 \cdot b + 1) =$ pascalparity(a, b).

Proof. The goal follows from pascalparityoddodd(a, b). □

Theorem 4.305. *For every natural number n and natural number k,* pascalparity$(n + 1, k + 1) =$ bxor(pascalparity(n, k), pascalparity$(n, k + 1)$).

Proof. The goal follows from pascalparitysucc(n, k). □

Theorem 4.306. *For every natural number n and natural number k, if $n < k$, then* pascalparity$(n, k) =$ false.

Proof. The goal follows from pascalparitygt(n, k). □

Theorem 4.307. *For every natural number n,* pascalparity$(n, n) =$ true.

Proof. The goal follows from pascalparityself(n). □

Theorem 4.308. *For every natural number n and natural number s,* constprefix$(n, s) =$ pascalparity(s, n).

Proof. The goal follows from constprefixeqlucas(n, s). □

Theorem 4.309. *For every natural number n and natural number s,* lineartower$(n, s) =$ pascalparity$(s, n + 1)$.

Proof. The goal follows from lineartowereqlucas(n, s). □

Theorem 4.310. *For every natural number k and natural number s,* masktransport$(k, s) =$ lucassum(k, s).

Proof. The goal follows from masktransportlucas(k, s). □

Theorem 4.311. *For every natural number t, if $2 \leq t$, then* center$(t) =$ not(lucassum$(t - 2, t)$).

Proof. The goal follows from centereqlucas(t). □

Theorem 4.312. *For every natural number T and natural number k and natural number m, if* centerperiodic$(T,$ twopow$(m))$*, then if $2 \leq k$, then if $T \leq$* twopow$(k)\cdot$twopow$(m)+3$*, then* latetail$(k, 3,$ twopow$(m)+$twopow$(m)) =$ not(latetail$(k, 3,$ twopow$(m)))$.

Proof. The goal follows from latetailthreedyadicflip(T, k, m). □

Theorem 4.313. *For every natural number K and natural number k, if* kappacollapse(K)*, then if $2 \leq k$, then* latetail$(k, 3,$ twopow$(K)+$twopow$(K)) =$ not(latetail$(k, 3,$ twopow$(K)))$.

Proof. The goal follows from kappacollapselatetailflip(K, k). □

Theorem 4.314. *For every natural number k, if $2 \leq k$, then* lateruler$(k, 0) =$ false *and* lateruler$(k, 1) =$ true $\wedge$ lateruler$(k, 2) =$ true $\wedge$ lateruler$(k, 3) =$ true.

Proof. The goal follows from laterulervalues(k). □

Theorem 4.315. *For every natural number i,* exposed$(0, i) =$ not(timeparity(i)).

Proof. The goal follows from exposedzero(i). □

Theorem 4.316. *For every natural number i,* exposed$(1, i) =$ false.

Proof. The goal follows from exposedone(i). □

Theorem 4.317. *For every natural number m and natural number n and natural number d and natural number q, if $d <$* twopow(m)*, then* pascalparity$(n+$ twopow$(m) \cdot q, d) =$ pascalparity(n, d).

Proof. The goal follows from pascalparityaddtwopowmul(m, n, d, q). □

Theorem 4.318. *For every natural number k and natural number J, if $1 \leq k$, then if* rightthreshold(k, J), *then $k + 1 \leq J$.*

Proof. The goal follows from rightthresholdge(k, J). □

Theorem 4.319. *For every natural number n and natural number k, if $k \leq n$, then* pascalparity$(n, k) =$ pascalparity$(n, n - k)$.

Proof. The goal follows from pascalparitysymm(n, k). □

Theorem 4.320. *For every natural number t, if $2 \leq t$, then* center$(t) =$ not(leftcone(t)).

Proof. The goal follows from centereqleftcone(t). □

Theorem 4.321. *For every natural number T and natural number P and natural number t, if* centerperiodic(T, P), *then if $T \leq t$, then if $2 \leq t$, then* leftcone$(t + P) =$ leftcone(t).

Proof. The goal follows from leftconecenterperiodic(T, P, t). □

Theorem 4.322. *For every natural number m and natural number n and natural number j and natural number q and natural number c, if $q <$* twopow(m), *then* coneterm$(n +$ twopow$(m) \cdot c, j, q) =$ coneterm(n, j, q).

Proof. The goal follows from conetermaddtwopowmul(m, n, j, q, c). □

Theorem 4.323. *For every natural number t, if $2 \leq t$, then* lucassum$(t - 2, t) =$ leftcone(t).

Proof. The goal follows from lucassumeqleftcone(t). □

Theorem 4.324. leftperiodic$(0, 0, 1)$.

Proof. The goal follows from leftperiodiczero. □

Theorem 4.325. leftperiodic$(1, 1, 1)$.

Proof. The goal follows from leftperiodicone. □

Theorem 4.326. *For every natural number k and natural number T and natural number p, if* leftperiodic(k, T, p), *then if* leftperiodic$(k+1, T, p)$, *then* leftperiodic$(k + 2, T + p, 2 \cdot p)$.

Proof. The goal follows from leftperiodicstep(k, T, p). □

Theorem 4.327. *For every natural number j,* $\mathrm{leftperiod}(j) = \mathrm{twopow}(j)$.

Proof. The goal follows from $\mathrm{leftperiodeqtwopow}(j)$. □

Theorem 4.328. *For every natural number j,* $\mathrm{leftperiodic}(j, \mathrm{lefttransient}(j), \mathrm{leftperiod}(j))$.

Proof. The goal follows from $\mathrm{ldiageventuallyperiodic}(j)$. □

Theorem 4.329. *For every natural number t,* $\mathrm{center}(t) = \mathrm{ldiag}(t, t)$.

Proof. The goal follows from $\mathrm{centereqldiagself}(t)$. □

Theorem 4.330. *For every natural number T and natural number P and natural number t, if* $\mathrm{centerperiodic}(T, P)$*, then if* $T \leq t$*, then* $\mathrm{ldiag}(t + P, t + P) = \mathrm{ldiag}(t, t)$.

Proof. The goal follows from $\mathrm{ldiagselfperiodicofcenterperiodic}(T, P, t)$. □

Theorem 4.331. *For every natural number k and natural number q, if* $\mathrm{orperiodic}(k, q)$*, then if* $\mathrm{towerword}(k + 2, q) = \mathrm{false}$*, then* $\mathrm{wordperiodic}(k + 2, q)$.

Proof. The goal follows from $\mathrm{wordperiodicofor}(k, q)$. □

Theorem 4.332. *For every natural number k and natural number q, if* $\mathrm{orperiodic}(k, q)$*, then* $\mathrm{wordperiodic}(k + 2, 2 \cdot q)$.

Proof. The goal follows from $\mathrm{wordperiodicdouble}(k, q)$. □

Theorem 4.333. *For every natural number k and natural number q, if* $\mathrm{wordperiodic}(k + 1, q)$*, then if* $\mathrm{wordperiodic}(k, q)$*, then* $\mathrm{orperiodic}(k, q)$.

Proof. The goal follows from $\mathrm{orperiodicofwords}(k, q)$. □

Theorem 4.334. *For every natural number j,* $\mathrm{wordperiodic}(j, \mathrm{twopow}(\mathrm{jenexp}(j)))$.

Proof. The goal follows from $\mathrm{towerwordjenperiodic}(j)$. □

Theorem 4.335. $\mathrm{jenexp}(2) = 1$ *and* $\mathrm{jenexp}(3) = 2 \wedge \mathrm{jenexp}(4) = 3 \wedge \mathrm{jenexp}(5) = 3 \wedge \mathrm{jenexp}(6) = 4 \wedge \mathrm{jenexp}(7) = 5 \wedge \mathrm{jenexp}(8) = 5 \wedge \mathrm{jenexp}(9) = 6$.

Proof. The goal follows from jenexpvalues. □

Theorem 4.336. *For every natural number j,* $\mathrm{jenexp}(j) \leq \mathrm{jenexp}(j + 1)$.

Proof. The goal follows from $\mathrm{jenexplesucc}(j)$. □

Theorem 4.337. *For every natural number j,* $\mathrm{jenexp}(j+1) \leq \mathrm{jenexp}(j)+1$.

Proof. The goal follows from $\mathrm{jenexpsuccle}(j)$. □

Theorem 4.338. *For every natural number k,* $\mathrm{jenexp}(k+2) = \mathrm{jenexp}(k+1) + \mathrm{boolnat}(\mathrm{towerword}(k+2, \mathrm{twopow}(\mathrm{jenexp}(k+1))))$.

Proof. The goal follows from $\mathrm{jenexpsuccsucceq}(k)$. □

Theorem 4.339. *For every natural number j,* $\mathrm{jenexp}(j) \leq j$.

Proof. The goal follows from $\mathrm{jenexple}(j)$. □

Theorem 4.340. *For every natural number t,* $\mathrm{leftcone}(t+1) = \mathrm{bxor}(\mathrm{leftcone}(t), \mathrm{bxor}(\mathrm{maskedcolumn}(t),$

Proof. The goal follows from $\mathrm{leftconesucc}(t)$. □

Theorem 4.341. *For every natural number t,* $\mathrm{bxor}(\mathrm{center}(t+1), \mathrm{center}(t)) = \mathrm{bxor}(\mathrm{maskedcolumn}(t), \mathrm{conezero}(t))$.

Proof. The goal follows from $\mathrm{centerincrementall}(t)$. □

Theorem 4.342. *For every natural number T and natural number P and natural number S and natural number t, if* $\mathrm{centerperiodic}(T, P)$*, then if* $\mathrm{conezeroperiodic}(S, P)$*, then if $T \leq t$, then if $S \leq t$, then* $\mathrm{maskedcolumn}(t+P) = \mathrm{maskedcolumn}(t)$.

Proof. The goal follows from $\mathrm{maskedcolumnofperiodic}(T, P, S, t)$. □

Theorem 4.343. *For every natural number T and natural number P and natural number S, if $0 < P$, then if* $\mathrm{centerperiodic}(T, P)$*, then not* $\mathrm{conezeroperiodic}(S, P)$.

Proof. The goal follows from $\mathrm{notconezeroperiodicfrom}(T, P, S)$. □

Theorem 4.344. *For every natural number t,* $\mathrm{conezero}(t) = \mathrm{cell}(t, -1)$.

Proof. The goal follows from $\mathrm{conezeroeqcell}(t)$. □

Theorem 4.345. *For every natural number t,* $\mathrm{bxor}(\mathrm{center}(t+1), \mathrm{center}(t)) = \mathrm{bxor}(\mathrm{maskedcolumn}(t), \mathrm{cell}(t, -1))$.

Proof. The goal follows from $\mathrm{centerincrementeqrule}(t)$. □

Theorem 4.346. *If* noperiodicblackconsistent*, then* problemone.

Proof. Assume h. The goal follows from $\mathrm{problem1ofnoperiodicblackconsistent}(h)$. □

Theorem 4.347. *For every natural number d and natural number t, if* $\mathrm{twopow}(d) \leq t$, *then* $\mathrm{diagonalcone}(d, t+3 \cdot \mathrm{twopow}(d)) = \mathrm{diagonalcone}(d, t)$.

Proof. The goal follows from $\mathrm{diagonalconeperiodic}(d, t)$. □

Theorem 4.348. *For every natural number w and natural number K and natural number z, if* $\mathrm{doublesat}(w \cdot \mathrm{twopow}(K), z)$, *then* $K \leq z$.

Proof. The goal follows from $\mathrm{leofdoublesatmultwopow}(w, K, z)$. □

Theorem 4.349. *For every natural number Q and natural number z and natural number y, if* $0 < Q$, *then if* $\mathrm{doublesat}(Q, z)$, *then if* $\mathrm{doublesat}(2 \cdot Q, y)$, *then* $z < y$.

Proof. The goal follows from $\mathrm{doublesattwomullt}(Q, z, y)$. □

Theorem 4.350. *For every natural number k,* problemone *if and only if* $\mathrm{noblockconeperiod}(k)$.

Proof. The goal follows from $\mathrm{problem1iffnoblockconeperiod}(k)$. □

Theorem 4.351. *For every natural number t and integer p,* $\mathrm{cell}(t, p) = \mathrm{bxor}(\mathrm{seedat}(t, p), \mathrm{coneat}(p, t))$.

Proof. The goal follows from $\mathrm{celleqseedatbxorconeat}(t, p)$. □

Theorem 4.352. *For every natural number t,* $\mathrm{center}(t) = \mathrm{not}(\mathrm{coneat}(0, t))$.

Proof. The goal follows from $\mathrm{centereqnotconeatzero}(t)$. □

Theorem 4.353. *For every natural number j and natural number m,* $\mathrm{wordperiodic}(j, \mathrm{twopow}(m))$ *if and only if* $\mathrm{wordmahlervanish}(j, m)$.

Proof. The goal follows from $\mathrm{wordperiodiciffmahler}(j, m)$. □

Theorem 4.354. *For every natural number j and natural number m, if not* $\mathrm{wordperiodic}(j, \mathrm{twopow}(m))$, *then* $\mathrm{wordmahlerabove}(j, m)$.

Proof. The goal follows from $\mathrm{towerwordmahlerdegree}(j, m)$. □

Theorem 4.355. noperiodicblackconsistent *if and only if* singleseedmask.

Proof. The goal follows from noperiodicblackconsistentiffsingleseedmask. □

Theorem 4.356. *If* noperiodicblackconsistent, *then* nozcenterperiod.

Proof. Assume h. The goal follows from nozcenterperiodofnoperiodicblackconsistent(h). □

Theorem 4.357. noperiodicisolatedblackconsistent *if and only if not* walledap.

Proof. The goal follows from noperiodicisolatedblackconsistentiff. □

Theorem 4.358. *If* noperiodicblackconsistent, *then* noperiodicisolatedblackconsistent.

Proof. Assume h. The goal follows from noperiodicisolatedofnoperiodicblackconsistent(h). □

Theorem 4.359. whitedefectinwindow.

Proof. The goal follows from whitedefectinwindowholds. □

Theorem 4.360. *For every natural number* t, $\mathrm{rdiag}(2, t) = \mathrm{timeparity}(t)$.

Proof. The goal follows from rdiag2eqtimeparity(t). □

Theorem 4.361. *For every natural number* t, $\mathrm{ldiag}(3, t) = \mathrm{bxor}(\mathrm{timeparity}(t), \mathrm{ldiag}(2, t))$.

Proof. The goal follows from ldiag3eqtimeparityxorldiag2(t). □

Theorem 4.362. *For every natural number* t, $\mathrm{ldiag}(4, t) = \mathrm{bxor}(\mathrm{ldiag}(1, t), \mathrm{ldiag}(2, t))$.

Proof. The goal follows from ldiag4eqldiag1xorldiag2(t). □

Theorem 4.363. *For every natural number* t, $\mathrm{ldiag}(5, t) = \mathrm{bxor}(\mathrm{timeparity}(t), \mathrm{ldiag}(2, t))$.

Proof. The goal follows from ldiag5eqtimeparityxorldiag2(t). □

Theorem 4.364. *For every natural number* t, $\mathrm{ldiag}(6, t) = \mathrm{bxor}(\mathrm{timeparity}(t), \mathrm{ldiag}(2, t))$.

Proof. The goal follows from ldiag6eqtimeparityxorldiag2(t). □

Theorem 4.365. *For every natural number* t, $\mathrm{ldiag}(8, t) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(3, t), \mathrm{ldiag}(1, t)), \mathrm{ldiag}(2, t))$

Proof. The goal follows from ldiag8eqrdiag3xorldiag1xorldiag2(t). □

Theorem 4.366. *For every natural number* t, $\mathrm{rdiag}(3, t{+}1) = \mathrm{bxor}(\mathrm{timeparity}(t), \mathrm{rdiag}(3, t))$.

Proof. The goal follows from rdiag3succeqtimeparityxorrdiag3(t). □

Theorem 4.367. *For every natural number* t, $\mathrm{rdiag}(4, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(3, t), \mathrm{rdiag}(4, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag4succeqrdiag3xorrdiag4xorexposed2(t). □

Theorem 4.368. *For every natural number t,* $\mathrm{rdiag}(5, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(4, t), \mathrm{rdiag}(5, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag5succeqrdiag4xorrdiag5xorexposed3(t). □

Theorem 4.369. *For every natural number t,* $\mathrm{rdiag}(6, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(5, t), \mathrm{rdiag}(6, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag6succeqrdiag5xorrdiag6xorexposed4(t). □

Theorem 4.370. *For every natural number t,* $\mathrm{rdiag}(7, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(6, t), \mathrm{rdiag}(7, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag7succeqrdiag6xorrdiag7xorexposed5(t). □

Theorem 4.371. *For every natural number t,* $\mathrm{rdiag}(8, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(7, t), \mathrm{rdiag}(8, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag8succeqrdiag7xorrdiag8xorexposed6(t). □

Theorem 4.372. *For every natural number t,* $\mathrm{rdiag}(9, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(8, t), \mathrm{rdiag}(9, t)), \mathrm{exposed}$

Proof. The goal follows from rdiag9succeqrdiag8xorrdiag9xorexposed7(t). □

Theorem 4.373. *For every natural number t,* $\mathrm{rdiag}(10, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(9, t), \mathrm{rdiag}(10, t)), \mathrm{expo}$

Proof. The goal follows from rdiag10succeqrdiag9xorrdiag10xorexposed8(t). □

Theorem 4.374. *For every natural number t,* $\mathrm{ldiag}(2, t{+}1) = \mathrm{not}(\mathrm{ldiag}(1, t))$.

Proof. The goal follows from ldiag2succeqnotldiag1(t). □

Theorem 4.375. *For every natural number t,* $\mathrm{ldiag}(10, t{+}1) = \mathrm{bxor}(\mathrm{bxor}(\mathrm{bxor}(\mathrm{rdiag}(3, t), \mathrm{ldiag}(1, t)), \mathrm{l}$

Proof. The goal follows from ldiag10succeqrdiag3xorldiag1xorldiag2xorldiag9(t). □

Theorem 4.376. *For every natural number t,* $\mathrm{ldiag}(12, t{+}1) = \mathrm{bxor}(\mathrm{ldiag}(10, t), \mathrm{ldiag}(12, t))$.

Proof. The goal follows from ldiag12succeqldiag10xorldiag12(t). □

Theorem 4.377. *For every natural number t,* $\mathrm{exposed}(2, t{+}1) = \mathrm{not}(\mathrm{bxor}(\mathrm{rdiag}(3, t), \mathrm{exposed}(2, t)))$.

Proof. The goal follows from exposed2succeqnotrdiag3xorexposed2(t). □

Theorem 4.378. *For every natural number t,* exposed$(3, t+1)$ = bxor(bxor(bxor(rdiag$(3, t)$, rdiag$(4, t)$))

Proof. The goal follows from exposed3succeqrdiag3xorrdiag4xorexposed3xorexposed4(t). □

Theorem 4.379. *For every natural number t,* exposed$(4, t+1)$ = bxor(bxor(timeparity(t), rdiag$(4, t)$), e

Proof. The goal follows from exposed4succeqtimeparityxorrdiag4xorexposed4(t). □

Theorem 4.380. *For every natural number t,* rdiag$(3, 2 \cdot t)$ = timeparity(t).

Proof. The goal follows from rdiag3eveneqtimeparity(t). □

Theorem 4.381. *For every natural number t,* rdiag$(4, 2 \cdot t)$ = bxor(timeparity(t), rdiag$(3, t)$).

Proof. The goal follows from rdiag4eveneqtimeparityxorrdiag3(t). □

Theorem 4.382. *For every natural number t,* rdiag$(5, 2 \cdot t)$ = false.

Proof. The goal follows from rdiag5eveneqfalse(t). □

Theorem 4.383. *For every natural number t,* rdiag$(6, 2 \cdot t)$ = bxor(bxor(bxor(timeparity(t), rdiag$(3, t)$)),

Proof. The goal follows from rdiag6eveneqtimeparityxorrdiag3xorrdiag4xorexposed2(t). □

Theorem 4.384. *For every natural number t,* rdiag$(7, 2 \cdot t)$ = bxor(bxor(bxor(bxor(bxor(timeparity(t), r

Proof. The goal follows from relationf42bb48334ad(t). □

Theorem 4.385. *For every natural number t,* ldiag$(1, 2 \cdot t)$ = ldiag$(1, t)$.

Proof. The goal follows from ldiag1eveneqldiag1(t). □

Theorem 4.386. *For every natural number t,* ldiag$(2, 2 \cdot t)$ = false.

Proof. The goal follows from ldiag2eveneqfalse(t). □

Theorem 4.387. *For every natural number t,* ldiag$(11, 2 \cdot t)$ = bxor(bxor(bxor(bxor(ldiag$(2, t)$, ldiag$(9, t$

Proof. The goal follows from relation9b630bd3f3ff(t). □

Theorem 4.388. *For every natural number t,* ldiag$(12, 2 \cdot t)$ = bxor(timeparity(t), ldiag$(2, t)$).

Proof. The goal follows from ldiag12eveneqtimeparityxorldiag2(t). □

Theorem 4.389. *For every natural number t,* exposed$(2, 2 \cdot t)$ = false.

Proof. The goal follows from exposed2eveneqfalse(t). □

Theorem 4.390. *For every natural number t,* exposed$(3, 2{\cdot}t)$ = bxor(timeparity(t), exposed$(2, t)$).

Proof. The goal follows from exposed3eveneqtimeparityxorexposed2(t). □

Theorem 4.391. *For every natural number t,* exposed$(4, 2{\cdot}t)$ = bxor(timeparity(t), rdiag$(3, t)$).

Proof. The goal follows from exposed4eveneqtimeparityxorrdiag3(t). □

Theorem 4.392. *For every natural number t,* exposed$(5, 2 \cdot t)$ = false.

Proof. The goal follows from exposed5eveneqfalse(t). □

Theorem 4.393. *For every natural number t,* exposed$(7, 2{\cdot}t)$ = bxor(bxor(bxor(bxor(bxor(rdiag$(6, t)$, e
1)), rdiag$(8, 2 \cdot t)$), exposed$(6, 2 \cdot t)$).

Proof. The goal follows from relation1de4355b13b0(t). □

Theorem 4.394. *For every natural number t,* ldiag$(11, 2{\cdot}t{+}1)$ = bxor(bxor(bxor(bxor(timeparity(t), ldi

Proof. The goal follows from relation084ca6f82327(t). □

Theorem 4.395. *For every natural number t,* exposed$(5, 2{\cdot}t{+}1)$ = bxor(bxor(bxor(timeparity(t), rdiag(

Proof. The goal follows from exposed5oddeqtimeparityxorrdiag5xorexposed2xorexposed3(t).
□

Theorem 4.396. *For every natural number s,* ldiag$(1, \mathrm{succ}(s))$ = true.

Proof. The goal follows from ldiagone(s). □