

# Basics of Applied Analysis A Report

Alifian Mahardhika Maulana

June 12, 2018

**Problem 1.** Let  $N \in \mathbb{N}, N \geq 2$ , and  $\alpha > 0$  be given.

We define  $A = (a_{ij})$  as finite difference discretization of :

$$\begin{cases} \alpha u - u'' = f \text{ in } (0, 1) \\ u(0) = u(1) = 0 \end{cases} \quad (1)$$

with entries

$$a_{ij} = \begin{cases} \alpha + \frac{2}{h^2}, & i = j \\ -\frac{1}{h^2}, & |i - j| = 1 \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

where  $h = \frac{1}{N}$

(a) To find eigenvalues  $\lambda$  s.t.  $Av = \lambda v$ , We define  $v_i := \sin\left(\frac{m\pi i}{N}\right)$  for some  $m \in \mathbb{N}$ , hence the discrete form of eigen problem becomes:

$$\begin{aligned} \sum_{i=1}^N a_{ij} v_j &= \lambda v_i \\ \left(\alpha + \frac{2}{h^2}\right) v_i - \frac{1}{h^2} v_{i+1} - \frac{1}{h^2} v_{i-1} &= \lambda v_i \\ \left(\alpha + \frac{2}{h^2} - \lambda\right) v_i - \frac{1}{h^2} (v_{i+1} + v_{i-1}) &= 0 \end{aligned}$$

substitute  $v_i$  to the discrete eigen problem, it becomes

$$\begin{aligned} \left(\alpha + \frac{2}{h^2} - \lambda\right) \sin\left(\frac{m\pi i}{N}\right) - \frac{1}{h^2} \left(\sin\left(\frac{m\pi(i+1)}{N}\right) + \sin\left(\frac{m\pi(i-1)}{N}\right)\right) &= 0 \\ \left(\alpha + \frac{2}{h^2} - \lambda\right) \sin\left(\frac{m\pi i}{N}\right) - \frac{1}{h^2} \left(2 \sin\left(\frac{m\pi i}{N}\right) \cos\left(\frac{m\pi}{N}\right)\right) &= 0 \\ \sin\left(\frac{m\pi i}{N}\right) \left(\alpha + \frac{2}{h^2} - \lambda - \frac{1}{h^2} 2 \cos\left(\frac{m\pi}{N}\right)\right) &= 0 \end{aligned}$$

to find  $\lambda$ , we set  $\sin\left(\frac{m\pi i}{N}\right)$  equal to some constant  $C$ , so that:

$$\begin{aligned} \left(\alpha + \frac{2}{h^2} - \lambda - \frac{2}{h^2} \cos\left(\frac{m\pi}{N}\right)\right) &= 0 \\ \therefore \lambda &= \alpha + \frac{2}{h^2} \left(1 - \cos\left(\frac{m\pi}{N}\right)\right) \end{aligned} \quad (3)$$

With Eigenvectors:

$$\therefore v = \sin\left(\frac{m\pi i}{N}\right)$$

(b) We define spectral radius  $\sigma(A) := \max\{|\lambda|\}$ , using  $\lambda$  computed in (3), we get:

$$\begin{aligned}\sigma(A) &= \max\{|\lambda|\} \\ &= \alpha + \frac{2}{h^2} \left( 1 - \cos \left( \frac{m\pi}{N} \right) \right)\end{aligned}$$

then, thanks to the symmetricity of  $\lambda$ , we can use taylor expansion to approximate the value of

$$\cos \left( \frac{m\pi}{N} \right) \approx 1 - \frac{1}{2} \left( \frac{\pi}{N} \right)^2$$

, hence the spectral radius  $\sigma(A)$  can be calculated by:

$$\begin{aligned}\sigma(A) &\approx \alpha + \frac{2}{h^2} \left( 1 - \left( 1 - \frac{m^2\pi^2}{2N^2} \right) \right) \\ &\approx \alpha + \frac{2}{h^2} \frac{m^2\pi^2}{2N^2} \\ \therefore \sigma(A) &\approx \alpha + (m\pi)^2\end{aligned}$$

(c) Let:

$$R := -D^{-1}(L + U)$$

be the Jacobi iteration matrix, we want to find the eigenvalues and eigenvectors s.t.  $Rv = \lambda v$ .

$$\begin{aligned}Rv &= \lambda v \\ -D^{-1}(L + U)v &= \lambda v \\ -(L + U)v &= \lambda Dv\end{aligned}$$

the discrete form of the eigen problem is:

$$\frac{1}{h^2}(v_{i-1} + v_{i+1}) = \lambda \left( \alpha + \frac{2}{h^2} \right) v_i$$

and then we set,  $v_i := \sin \left( \frac{m\pi i}{N} \right)$  and substitute to the discrete form of the eigen problem, thus:

$$\begin{aligned}\frac{1}{h^2} \left( \sin \left( \frac{m\pi(i+1)}{N} \right) + \sin \left( \frac{m\pi(i-1)}{N} \right) \right) &= \lambda \left( \alpha + \frac{2}{h^2} \right) \sin \left( \frac{m\pi i}{N} \right) \\ \frac{1}{h^2} \left( 2 \sin \left( \frac{m\pi i}{N} \right) \cos \left( \frac{m\pi}{N} \right) \right) &= \lambda \left( \alpha + \frac{2}{h^2} \right) \sin \left( \frac{m\pi i}{N} \right) \\ \sin \frac{m\pi i}{N} \left( \frac{2}{h^2} \cos \left( \frac{m\pi}{N} \right) - \lambda \left( \alpha + \frac{2}{h^2} \right) \right) &= 0\end{aligned}$$

to find  $\lambda$ , we set  $\sin \frac{m\pi i}{N}$  equal to some constant  $D$ , so that:

$$\begin{aligned}\left( \frac{2}{h^2} \cos \left( \frac{m\pi}{N} \right) - \lambda \left( \alpha + \frac{2}{h^2} \right) \right) &= 0 \\ \lambda \left( \alpha + \frac{2}{h^2} \right) &= \frac{2}{h^2} \cos \left( \frac{m\pi}{N} \right)\end{aligned}$$

$$\text{with } h = \frac{1}{N}$$

$$\lambda = \frac{\frac{2}{h^2} \cos \left( \frac{m\pi}{N} \right)}{\alpha + \frac{2}{h^2}}$$

(4)

with  $h = \frac{1}{N}$ ,

$$\therefore \lambda = \frac{2N^2 \cos\left(\frac{m\pi}{N}\right)}{\alpha + 2N^2}$$

(d) We define spectral radius  $\sigma(R) := \max\{|\lambda|\}$ , using  $\lambda$  computed in (4), we get:

$$\begin{aligned}\sigma(R) &= \max\{|\lambda|\} \\ &= \frac{2N^2 \cos\left(\frac{m\pi}{N}\right)}{\alpha + 2N^2}\end{aligned}$$

again, thanks to the symmetricity of  $\lambda$ , we can use taylor expansion to approximate the value of

$$\cos\left(\frac{m\pi}{N}\right) \approx 1 - \frac{1}{2}\left(\frac{m\pi}{N}\right)^2$$

, hence the spectral radius  $\sigma(R)$  can be calculated by:

$$\therefore \sigma(R) \approx \frac{2N^2\left(1 - \frac{m^2\pi^2}{2N^2}\right)}{\alpha + 2N^2}$$

**Problem 2.** Let  $N, h = \frac{1}{N}, \alpha$  and  $A$  be as in Problem 1. We define:

$$\begin{cases} \alpha u - u'' = \sin(\pi x) & \text{in } (0, 1) \\ u(0) = u(1) = 0 \end{cases} \quad (5)$$

(a) To find the exact solution of  $u$ , we use general solution for Ordinary Differential Equation (ODE):

$$u(x) = A \sin(\pi x) + B \cos(\pi x) \quad (6)$$

and take the second derivative of  $u$

$$u''(x) = -A\pi^2 \sin(\pi x) + B\pi^2 \cos(\pi x) \quad (7)$$

then substitute (6) and (7) to (5), we get:

$$\begin{aligned} \alpha(A \sin(\pi x) + B \cos(\pi x)) + A\pi^2 \sin(\pi x) + B\pi^2 \cos(\pi x) &= \sin(\pi x) \\ A(\alpha + \pi^2) \sin(\pi x) + B(\alpha + \pi^2) \cos(\pi x) &= \sin(\pi x) \end{aligned} \quad (8)$$

from (8) we know that

$$A(\alpha + \pi^2) \sin(\pi x) = \sin(\pi x)$$

and

$$B(\alpha + \pi^2) \cos(\pi x) = 0$$

$$\therefore A = \frac{1}{(\alpha + \pi^2)} \quad \therefore B = 0 \quad (9)$$

then we substitute (9) to (6), we get the exact solution as:

$$u(x) = \frac{\sin(\pi x)}{(\alpha + \pi^2)} \quad (10)$$

(b) To find exact solution s.t.  $Av = b$ , with  $b_i = \sin(\pi hi)$ , we set

$$v_i = C \sin(\pi hi)$$

thus,

$$\begin{aligned} Av &= b \\ \sum_{i=1}^{N-1} a_{ij} v_i &= b_i \\ \left( \left( \alpha + \frac{2}{h^2} \right) v_i - \frac{1}{h^2} (v_{i+1} + v_{i-1}) \right) &= \sin(\pi hi) \\ \left( \left( \alpha + \frac{2}{h^2} \right) C \sin(\pi hi) - \frac{1}{h^2} (C \sin(\pi h(i+1)) + C \sin(\pi h(i-1))) \right) &= \sin(\pi hi) \\ \left( \left( \alpha + \frac{2}{h^2} \right) C \sin(\pi hi) - C \frac{1}{h^2} (2 \sin(\pi hi) \cos(\pi h)) \right) &= \sin(\pi hi) \end{aligned}$$

divide by  $\sin(\pi hi)$  on the both side we get:

$$C \left( \left( \alpha + \frac{2}{h^2} \right) - \frac{2}{h^2} \cos(\pi h) \right) = 1$$

$$\therefore C = \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)}$$

so the exact solution in terms of  $v_i$  is:

$$v_i = \frac{\sin(\pi h i)}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)}$$

(c) We define

$$\epsilon(h) := \max_{1 \leq i \leq N-1} |u(hi) - v_i|$$

then we use  $u(x)$  and  $v_i$  that we derived before to find the explicit formula of  $\epsilon(h)$ :

$$\begin{aligned} \epsilon(h) &:= \max_{1 \leq i \leq N-1} \left| \frac{\sin(\pi h i)}{(\alpha + \pi^2)} - \frac{\sin(\pi h i)}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)} \right| \\ \epsilon(h) &:= \max_{1 \leq i \leq N-1} \left| \sin(\pi h i) \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)} \right) \right| \end{aligned}$$

for simplicity, we rewrite:

$$\begin{aligned} D &:= \sin(\pi h i) \\ E &:= \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)} \right) \\ \max_{1 \leq i \leq N-1} |DE| &= \max_{1 \leq i \leq N-1} |D| \max_{1 \leq i \leq N-1} |E| \\ &= 1 \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)} \right) \end{aligned}$$

then, the explicit formula for  $\epsilon(h)$  is:

$$\epsilon(h) = \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \cos(\pi h)\right)} \right) \quad (11)$$

around  $h = 0$ , leading order of taylor expansion of  $\epsilon(h)$  can be obtained by expand  $\cos(\pi h)$ , so that

$$\cos(\pi h) \approx \cos(a) + \frac{\cos'(a)(\pi h - a)}{1!} + \frac{\cos''(a)(\pi h - a)^2}{2!} + \dots$$

then we take  $a = 0$ , we get

$$\cos(\pi h) \approx 1 - \frac{1}{2}(\pi h)^2$$

substitute it to (11) we get:

$$\begin{aligned} \epsilon(h) &= \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{\left(\left(\alpha + \frac{2}{h^2}\right) - \frac{2}{h^2} \left(1 - \frac{1}{2}(\pi h)^2\right)\right)} \right) \\ \epsilon(h) &= \left( \frac{1}{(\alpha + \pi^2)} - \frac{1}{(\alpha + \pi^2)} \right) \end{aligned}$$

*which is just a constant, so the leading order is depend on what order of taylor expansion we choose to approximate the value of  $\cos(\pi h)$  around  $h = 0$ . In this case if we choose taylor expansion orde 2, we get the leading term of  $\epsilon$  is  $\mathbb{O}(F)$  with  $F$  is a constant.*

**Problem 3.** We consider system of linear equations:

$$\begin{cases} -v_{i-1,j} - v_{i+1,j} - v_{i,j-1} - v_{i,j+1} + 4v_{i,j} = b_{i,j} & i, j = 1, \dots, N-1 \\ v_{0,j} = v_{0,N} = v_{i,0} = v_{i,N} = 0 & i, j = 1, \dots, N-1 \end{cases} \quad (12)$$

for unknowns  $v_{i,j}, i, j = 1, \dots, N-1$ .

- (a) We want to find the eigenvalues and eigenvectors of the matrix  $A$  for the system (12). Using the idea of discrete separation variables, we set  $w_{i,j} = v_i \tilde{v}_j$  s.t.

$$Aw_{i,j} = \lambda w_{i,j}$$

$$\begin{aligned} 4w_{i,j} - w_{i-1,j} - w_{i+1,j} - w_{i,j-1} - w_{i,j+1} &= \lambda w_{i,j} \\ 4(v_i \tilde{v}_j) - (v_{i-1} \tilde{v}_j) - (v_{i+1} \tilde{v}_j) - (v_i \tilde{v}_{j-1}) - (v_i \tilde{v}_{j+1}) &= \lambda(v_i \tilde{v}_j) \\ 4(v_i \tilde{v}_j) - \tilde{v}_j(v_{i-1} + v_{i+1}) - v_i(\tilde{v}_{j-1} + \tilde{v}_{j+1}) &= \lambda(v_i \tilde{v}_j) \\ (4 - \lambda)(v_i \tilde{v}_j) - \tilde{v}_j(v_{i-1} + v_{i+1}) - v_i(\tilde{v}_{j-1} + \tilde{v}_{j+1}) &= 0 \end{aligned}$$

then, divide by  $v_i \tilde{v}_j$  on both side, we get:

$$\begin{aligned} (4 - \lambda) - \frac{(v_{i-1} + v_{i+1})}{v_i} - \frac{(\tilde{v}_{j-1} + \tilde{v}_{j+1})}{\tilde{v}_j} &= 0 \\ (4 - \lambda) - \frac{(v_{i-1} + v_{i+1})}{v_i} &= \frac{(\tilde{v}_{j-1} + \tilde{v}_{j+1})}{\tilde{v}_j} \end{aligned}$$

now, to satisfies equality, both side should be equal to some constant, let say  $G$ , hence:

$$(4 - \lambda) - \frac{(v_{i-1} + v_{i+1})}{v_i} = G = \frac{(\tilde{v}_{j-1} + \tilde{v}_{j+1})}{\tilde{v}_j}$$

First, we solve for:

$$\begin{aligned} \frac{(\tilde{v}_{j-1} + \tilde{v}_{j+1})}{\tilde{v}_j} &= G \\ (\tilde{v}_{j-1} + \tilde{v}_{j+1}) &= \tilde{v}_j G \end{aligned}$$

set:  $\tilde{v}_j = \varphi^j$

$$\begin{aligned} \varphi^{j-1} + \varphi^{j+1} &= \varphi^j G \\ \varphi^{j-1} + \varphi^{j+1} - \varphi^j G &= 0 \end{aligned}$$

divide both side by  $\varphi^{j-1}$

$$\begin{aligned} 1 + \varphi^2 - \varphi G &= 0 \\ \therefore \varphi_{\pm} &= \frac{G \pm \sqrt{(G^2 - 4)}}{2} \end{aligned} \quad (13)$$

here we recall that eigenvector of  $\tilde{v}_j$  should be a linear combination of  $\varphi$  s.t.

$$\tilde{v}_j = c_1 \varphi_+^j + c_2 \varphi_-^j$$

and we know the boundary condition on (12), when  $j = N$  then  $\tilde{v}_N = 0$ , to satisfies this condition, then  $c_1 = -c_2$ , then

$$\tilde{v}_j = c_1(\varphi_+^j - \varphi_-^j)$$

$\therefore \varphi_+^j$  and  $\varphi_-^j$  should be distinguished.

Hence, we choose  $G < 2$  for (13) we then rewrite (13) as:

$$\begin{aligned} \varphi_{\pm} &= \frac{G \pm \sqrt{-(4 - G^2)}}{2} \\ \varphi_{\pm} &= \frac{G \pm i\sqrt{(4 - G^2)}}{2} \end{aligned}$$

then, the possible solution is

$$\varphi_{\pm}^j = \cos(\theta j) \pm i \sin(\theta j)$$

$$\begin{aligned} \therefore \tilde{v}_j &= c_1(\cos(\theta j) + i \sin(\theta j) - (\cos(\theta j) - i \sin(\theta j))) \\ \tilde{v}_j &= c_1(2i \sin(\theta j)) \end{aligned}$$

we choose  $c_1 = -\frac{i}{2}$  so that

$$\tilde{v}_j = \sin(\theta j)$$

inserting boundary condition, when  $j = N$  then  $\tilde{v}_N = 0$ , we get:

$$\begin{aligned} 0 &= \sin(\theta N) \\ \theta &= m\pi, \quad \forall m : 2, 4, 6 \cdots N, \text{ even} \\ \theta &= \frac{m\pi}{N} \end{aligned}$$

then, we know that from imager triangle

$$\cos \theta = \frac{G}{2}$$

$$\therefore G = 2 \cos(\theta) = 2 \cos\left(\frac{m\pi}{N}\right)$$

Second, we solve for:

$$(4 - \lambda) - \frac{(v_{i-1} + v_{i+1})}{v_i} = G$$

multiplied both side by  $v_i$ , we get:

$$\begin{aligned} v_i(4 - \lambda) - (v_{i-1} + v_{i+1}) &= v_i G \\ v_i(4 - \lambda) - (v_{i-1} + v_{i+1}) - v_i G &= 0 \\ v_i(4 - \lambda - G) - (v_{i-1} + v_{i+1}) &= 0 \end{aligned}$$

set  $v_i = \xi^i$

$$\xi^i(4 - \lambda - G) - (\xi^{i-1} + \xi^{i+1}) = 0$$

divide both side by  $\xi^{i-1}$

$$\begin{aligned} \xi(4 - \lambda - G) - (1 + \xi^2) &= 0 \\ \xi^2 - \xi(4 - \lambda - G) + 1 &= 0 \\ \therefore \xi_{\pm} &= \frac{(4 - \lambda - G) \pm \sqrt{(4 - \lambda - G)^2 - 4}}{2} \end{aligned}$$

take  $2H = (4 - \lambda - G)$

$$\begin{aligned} \xi_{\pm} &= \frac{2H \pm \sqrt{(2H)^2 - 4}}{2} \\ \xi_{\pm} &= H \pm \sqrt{H^2 - 1} \end{aligned} \tag{14}$$

here we recall that eigenvector of  $v_i$  should be a linear combination of  $\xi$  s.t.

$$v_i = d_1 \xi_+^i + d_2 \xi_-^i$$

and we know the boundary condition on (12), when  $i = N$  then  $v_N = 0$ , to satisfies this condition, then  $d_1 = -d_2$ , then

$$v_i = d_1(\xi_+^i - \xi_-^i)$$



$\therefore \xi_+^j$  and  $\xi_-^j$  should be distinguished.

Hence, we choose  $H < 1$  for (14) we then rewrite (14) as:

$$\begin{aligned}\xi_{\pm} &= H \pm \sqrt{-(1-H^2)} \\ \xi_{\pm} &= H \pm i\sqrt{(1-H^2)}\end{aligned}$$

then, the possible solution is

$$\xi_{\pm}^i = \cos(\theta i) \pm I \sin(\theta i)$$

here i use "I" as imaginer number, so it is not confusing between "i" index and "I" imaginer number.

$$\begin{aligned}\therefore v_i &= d_1(\cos(\theta i) + I \sin(\theta i) - (\cos(\theta i) - I \sin(\theta i))) \\ v_i &= d_1(2I \sin(\theta i))\end{aligned}$$

we choose  $d_1 = -\frac{I}{2}$  so that

$$v_i = \sin(\theta i)$$

inserting boundary condition, when  $i = N$  then  $v_N = 0$ , we get:

$$\begin{aligned}0 &= \sin(\theta N) \\ \theta &= m\pi, \quad \forall m : 2, 4, 6 \cdots N, \text{ even} \\ \theta &= \frac{m\pi}{N}\end{aligned}$$

then, we know that from imaginer triangle

$$\cos \theta = H$$

$$\therefore H = \cos(\theta) = \cos\left(\frac{m\pi}{N}\right)$$

substitute  $H$  and  $G$  to  $2H = (4 - \lambda - G)$ , we get

$$\begin{aligned}2 \cos\left(\frac{m\pi}{N}\right) &= 4 - \lambda - 2 \cos\left(\frac{m\pi}{N}\right) \\ \lambda &= 4 - 2 \cos\left(\frac{m\pi}{N}\right) - 2 \cos\left(\frac{m\pi}{N}\right) \\ \therefore \lambda &= 4 - 2\left(\cos\left(\frac{m\pi}{N}\right) + \cos\left(\frac{m\pi}{N}\right)\right)\end{aligned}$$

for the eigenvectors, we already know that

$$\begin{aligned}w_{i,j} &= v_i \tilde{v}_j \\ v_i &= \sin(\theta i) \\ \tilde{v}_j &= \sin(\theta j) \\ \therefore w_{i,j} &= \sin\left(\frac{m\pi i}{N}\right) \sin\left(\frac{m\pi j}{N}\right)\end{aligned}$$

(b) I attached the Python Code that solves (12) in 1

(c) Minimal Number of Iterations  $K$  such the:

$$\max_{1 \leq i,j \leq N-1} \left| v_{i,j}^{(K+1)} - v_{i,j}^{(K)} \right| \leq 10^{-4}$$

(a)  $N = 10$

- i. Jacobi,  $K = 53$
- ii. Gauss-Seidel,  $K = 33$
- iii. SOR ( $\omega=1.5$ ),  $K = 12$

(b)  $N = 20$

- i. Jacobi,  $K = 137$
- ii. Gauss-Seidel,  $K = 94$
- iii. SOR ( $\omega=1.5$ ),  $K = 45$

(c)  $N = 50$

- i. Jacobi,  $K = 1$
- ii. Gauss-Seidel,  $K = 173$
- iii. SOR ( $\omega=1.5$ ),  $K = 148$

(d) Optimal  $\omega_b$  for SOR:

- (a)  $N = 10$ ,  $\omega_b = 1.16527511683289$
- (b)  $N = 20$ ,  $\omega_b = 1.2933223938735654$
- (c)  $N = 50$ ,  $\omega_b = 1.3746381523099196$

# 1 Attachment

```
import numpy as np

#Function Definition
#Function to create Right Hand Side and Left Hand Side Matrix
def const_mat(N):
    h = 1/N
    A = np.zeros((N,N))
    b = np.zeros((N,N))
    for i in range(N):
        for j in range(N):
            A[i][i] = 4
            b[i][j] = h*h*(np.sin(np.pi*i/N)*np.sin(np.pi*j/N))
            #b[i][j] = 1
            if (i==j-1 and i%4==3) or (i==j+1 and i%4==0):
                A[i][j] = 0
            elif i==j-1:
                A[i][j] = -1
            elif i==j+1:
                A[i][j] = -1
            elif i==j-4:
                A[i][j] = -1
            elif i==j+4:
                A[i][j] = -1
    return A,b

#Jacobi
def jacobi(b):
    v = np.zeros_like(b)
    N = len(b)
    for k in range(1000):
        v_old = np.copy(v)
        for i in range(N-2):
            for j in range(N-2):
                v[i+1][j+1] = 0.25 * (v_old[i][j+1] + v_old[i+2][j+1] + v_old[i+1][j] + v_old[i+1][j+2] + b[i+1][j+1])
        #print(np.max(v))
        if np.max(np.abs(v - v_old)) < 1e-4:
            break
    return (k)

#Gauss-Seidel
def gauss_seidel(b):
    v = np.zeros_like(b)
    N = len(b)
    for k in range(1000):
        v_old = np.copy(v)
        for i in range(N-2):
            for j in range(N-2):
                v[i+1][j+1] = 0.25 * (v[i][j+1] + v[i+2][j+1] + v[i+1][j] + v[i+1][j+2] + b[i+1][j+1])
        #print(np.max(v))
        if np.max(np.abs(v - v_old)) < 1e-4:
            break
    return (k)

#SOR
def sor(omega,b):
    v = np.zeros_like(b)
    N = len(b)
    for k in range(1000):
        v_old = np.copy(v)
        for i in range(N-2):
            for j in range(N-2):
                v[i+1][j+1] = (1-omega)*v_old[i+1][j+1] + omega*(0.25 * (v[i][j+1] + v[i+2][j+1] + v[i+1][j] + v[i+1][j+2] + b[i+1][j+1]))
        #print(np.max(v))
        if np.max(np.abs(v - v_old)) < 1e-4:
            break
    return (k)

#Calculate Optimum Omega for SOR
def calc_op_omega(A):
    D = np.diag(np.diag(A))
    L = np.tril(A) - D
    U = np.triu(A) - D
    T = 1/np.diag(A)*(L+U)
    rho = np.max(np.linalg.eigvals(T))
    opt_omega = 2/(1+np.sqrt(1-rho*rho))

    return opt_omega

#main
Ns = [10,20,30,40,50]
om = 1.5
for n in Ns:
    matA, matB = const_mat(n)
    iterj = jacobi(matB)
    iterg = gauss_seidel(matB)
    iters = sor(om,matB)
    op_om = calc_op_omega(matA)
    print("N=",n)
    print("Minimum Iteration Jacobi=",iterj)
    print("Minimum Iteration Gauss-Seidel=",iterg)
    print("Minimum Iteration SOR using omega(1.5)=",iters)
    print("Optimal omega for SOR=",op_om)
```