


Utilizing the 25m Dwingeloo Radio Telescope (DRT) to Study Fast Radio Bursts (FRBs)

August 21st, 2024


Maxwell A. Fine
Advisors: Dr. Tammo Jan
Dijkema & Prof Jason
Hessels



Here is me working at the DRT. Image credit Prof Jason Hessels.

Fast Radio Bursts (FRBs)

- **Fast Radio Bursts (FRBs):** brief, intense radio emission from unknown extragalactic sources that last milliseconds
- **FRB Characteristics:**
 - Detected across a wide frequency range 110 MHz to 8 GHz
 - Dispersion Measure (DM) causes a frequency-dependent arrival time delay
 - $E_{\text{iso}} \sim 10^{36-41}$ erg
 - ~2% are known to repeat
- **Possible Origins:** typical theories involve neutron stars & magnetars



Dynamic spectrum of the first detected Fast Radio Burst, known as the Lorimer burst. (Lorimer et al. 2007)

Dispersion (Measure)

- Free electrons in space cause a frequency dependent arrival time delay – known as a **dispersive** sweep
- DM has units of pc/cm³ (**column density**)
- Probe of the free electron density between us, and the source

$$\text{DM} = \int_0^L n_e \, dl$$

$$\Delta t = \frac{(e^-)^2}{2\pi m_e c} \left(\frac{1}{\nu_2^2} - \frac{1}{\nu_1^2} \right) \text{DM}$$

Question:

The Crab Pulsar is located $\sim 1,800$ pc away and has a DM of ~ 60 pc/cm³.

What is the average n_e between us and the Crab Pulsar?

First correct answer gets a box of Kraft!*

$$\text{DM} = \int_0^L n_e \, dl$$

Must be in person to claim



A link between repeating and non-repeating fast radio bursts through their energy distributions

Received: 17 January 2023

Accepted: 30 October 2023

Published online: 4 January 2024




F. Kirsten^{1,2}✉, O. S. Ould-Boukattine^{2,3}, W. Herrmann⁴, M. P. Gawroński⁵, J. W. T. Hessels^{2,3}, W. Lu⁶, M. P. Snelders^{2,3}, P. Chawla³, J. Yang¹, R. Blaauw², K. Nimmo⁷, W. Puchalska⁵, P. Wolak⁵ & R. van Ruiten³

Fast radio bursts (FRBs) are extremely energetic, millisecond-duration radio flashes that reach Earth from extragalactic distances. Broadly speaking, FRBs can be classified as repeating or (apparently) non-repeating. It is still unclear, however, whether the two types share a common physical origin and differ only in their activity rate. Here we report on an observing campaign that targeted one hyperactive repeating source, FRB 20201124A, for more than 2,000 h using four 25–32 m class radio telescopes. We detected 46 high-energy bursts, many more than one would expect given previous observations of lower-energy bursts using larger radio telescopes. We find a high-energy burst distribution that resembles that of the non-repeating FRB population, suggesting that apparently non-repeating FRB sources may simply be the rarest bursts from repeating sources. Also, we discuss how FRB 20201124A contributes strongly to the all-sky FRB rate and how similar sources would be observable even at very high redshift.

Motivation

- Recent interest in using 'small' 25m radio telescopes for high cadence FRB follow-up observations of repeating FRBs
- Limited number of observation time, and telescopes in the world
- The Dwingeloo Radio Telescope (DRT) is 25m
- Why not use the DRT?

Motivation




Broken power law fit for hyperactive repeating FRB 20201124A (Kirsten et al. 2024)

- A Westerbork 25m Dish observed for
 - 512 hours in L-band and detected 24 bursts
 - 650 hours in P-band and detected 0 bursts
- Kirsten et al used four 25–32m radio telescopes located throughout Europe
- Results indicate that (apparently) non-repeating FRBs could be rare bursts from repeating FRBs

The 25m Dwingeloo Radio Telescope (DRT)

- Hydrogen Maser clock
 - It's Westerbork's clock!
- Can observe simultaneously in L and P band
 - L-Band 1200-1400 MHz
 - P-Band 400-420 MHz
- Records data
 - ~ 1 Gb/s for baseband data
 - ~1 Gb/min for filterbank data
- Sensitivity is comparable to a single 25m Westerbork dish
 - 80% at P-Band
 - 20% at L-Band



Here is me with the DRT

Project Goals:

- **Have Fun**
- **Build a real time FRB detection pipeline for the DRT**
- **If lucky, detect an FRB**
- **Observe Crab Pulsar**


Project Goals:

- Have Fun ✓
- Build a real time FRB detection pipeline for the DRT
- If lucky, detect an FRB
- Observe Crab Pulsar




Real-Time FRB detection pipeline for the DRT

- Search's at $\pm 10\%$ DM, Boxcar sizes between 1-30, 30s integration for Radio Frequency Interference (RFI) find
- Records data in **filterbank** format files, and **baseband** (amplitude & phase) data
- Writes data in 10-minute chunks to search
- Pipeline is open source
<https://gitlab.camras.nl/dijkema/rbscripts>



Real-Time FRB detection pipeline for the DRT


- Pipeline is based on presto
 - rfifind
 - ddp1an.py
 - prepsubband
 - single_pulse_search.py
- Candidates are written as .h5 files



Diagnostic plot displaying the contents of a bright (SNR > 200) candidate .h5 file for the Crab Pulsar



Real-Time FRB detection pipeline for the DRT

- Most candidates are still RFI, or spurious, need to mitigate this
- **'Machine Learning' Techniques**
 - DBscan Clustering candidates
 - Fetch's `predict.py` sorts candidates into 'good' and 'bad'
- Save's baseband data if a "good" candidate is made
- Baseband data could be used in VLBI to localize FRBS



Diagnostic plot displaying the contents of a bright (SNR > 200) candidate .h5 file for the Crab Pulsar

DBscan Clustering



after clustering if there are 800+
candidates the filterbank file is not
processed further

Project Goals:

- Have Fun ✓
- Build a real time FRB detection pipeline for the DRT ✓
- If lucky, detect an FRB
- Observe Crab Pulsar






If lucky, detect an FRB

- Unlucky, no super bright hyperactive active repeaters over summer
- FRB 20240619D ~ 20 hours
- Other FRBs ~ 50 hours
- **Even no detections of FRBs is important!**
- Westerbork made several FRB detections that the DRT would have been able to observe

If lucky, detect an FRB

- Even no detections of FRBs is important!
- Westerbork made several FRB detections that the DRT would have been able to observe
- While the DRT did not make any detections, we did contribute to studying burst rates of FRB 20240619D




FRB 20240619D detected with 25m Westerbork dish on 2024-07-21 UTC. Ould-Boukattine et al. 2024, The Astronomer's Telegram, No. 16732.

Project Goals:

- Have Fun ✓
- Build a real time FRB detection pipeline for the DRT ✓
- If lucky, detect an FRB ⚡
- Observe Crab Pulsar



Crab Analysis



- Observed:
 - P-Band 29.28 Hours
 - L-Band 25.91 Hours
- Fitted Power Law Index (α) with maximum likelihood method of James et al. 2018
- Power Law results consistent with Bera et al. 2019

Summary

- The DRT now has a **real-time FRB detection pipeline** that can save **baseband data**
- DRT is capable of joining high cadence FRB follow-up observations
- When, not if the DRT makes an FRB detection its data can be used in the VLBI network to help localize FRBs
- Pipeline is open source:
https://gitlab.camras.nl/dijkema/frbs_cripts

Next Steps for the DRT

- Train Camras volunteers to run the observation program, write directions for them
- Recruit PhD researchers at Astron to use the DRT as their office, if they are willing to observe
- Increase bandwidth in L-band, P-Band
- Post the "good" candidates to the Camras mastodon bot
- Prototype of new receiver?