


Quantum optimization through path-integral molecular dynamics



PSI


Center for Scientific Computing,
Theory and Data

Alfredo Fiorentino¹, Nicola Marzari^{1,2}

¹ PSI Center for Scientific Computing, Theory, and Data, Villigen, Switzerland

² Theory and Simulation of Materials (THEOS), Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

MARVEL



1. Introduction

► Motivation

Global optimization of atomic degrees of freedom is a key problem in physics whenever the number of candidate structures (local minima) is combinatorially large: cluster physics, polymer folding, interfaces...

► The idea

Novel implementation of **Quantum Annealing** (QA) using **Path-Integral Molecular Dynamics** (PIMD), labeled QA-PIMD. While **Classical Annealing** (CA) involves melting the system and then slowly cooling it through a temperature schedule, QA relies on quantum delocalization.




Figure: Summary of the workflow

2. Methods

QA-PIMD is obtained by combining two powerful methods:

► Quantum annealing: exploration through quantum delocalization

$$\hat{H}(t) = -\alpha(t) \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla^2 + U(\hat{\mathbf{x}}_1, \dots, \hat{\mathbf{x}}_N)$$

$$\alpha(t=0) \gg 1, \alpha(t_{\max}) \ll 1$$

$$\lim_{t_{\max} \rightarrow \infty} \text{Tr}[\rho(t_{\max})U] = \min U$$


► Path-Integral Molecular Dynamics

Equilibrium nuclear quantum effects (NQE) from P classical simulations ($P \sim 10 - 100$).

$$H_P = \sum_{i=1}^N \sum_{j=1}^P \left[\frac{1}{2m_i} |\mathbf{p}_i^{(j)}|^2 + \frac{1}{2} m_i \omega_P^2 |\mathbf{x}_i^{(j)} - \mathbf{x}_i^{(j+1)}|^2 \right] + \sum_{j=1}^P U(\mathbf{x}_1^{(j)}, \dots, \mathbf{x}_N^{(j)}),$$
$$\omega_P = \frac{Pk_B T}{\hbar}$$

3. Benchmark: asymmetric double well

The local density from 200-ps PIMD simulations follows the exact instantaneous quantum density adiabatically. $T = 100$ K.



4.1 Lennard-Jones clusters

LJ clusters: millions of estimated local minima with only a few tens of atoms!




Figure: Global minimum of an LJ cluster with $N = 13$.

4.2 LJ clusters: classical vs quantum




Figure: Comparison of QA and CA residual energies as a function of annealing time, for two LJ cluster sizes.

5.1 Missing hydrogen sites

Many experimental structures lack hydrogen sites. We reconstruct 10 H-compounds using QA-PIMD with a foundational MLIP (MACE).





Figure: Example of identifying hydrogen sites in B_4H_{12} through QA-PIMD. Check the QR code for the video!

5.2 Results: missing hydrogen problem




Figure: Top: residual energy with respect to the MC3D database as a function of annealing time. Bottom: probability of finding an equal or lower energy.

6. Conclusion

- QA-PIMD is a novel, unbiased, and general global optimization algorithm using quantum fluctuations.
- It can outperform classical annealing; e.g., shorter annealing times are sufficient for LJ clusters.
- Material discovery: identifies new candidate stable crystals from the MC3D database.

Open questions and challenges:

- Small but **finite** temperature — not a ground-state method.
- PIMD time evolution is not exactly quantum; dynamically, $QA - PIMD \neq QA$.

7. References

1. A. Fiorentino and N. Marzari. *Quantum annealing for materials science*. Manuscript in preparation.
2. R. P. Feynman et al., *Quantum Mechanics and Path Integrals*, Physics Today 19, 89 (1966).
3. T. Gregor et al., *Chem. Phys. Lett.* 412, 125 (2005).
4. L. Stella et al., *Phys. Rev. B* 72, 014303 (2005).

8. Acknowledgements

We thank S. Baroni, L. Vojáček, A. Carta, V. Sanella, and S. Schären for insightful discussions. Financial support was provided by the NCCR MARVEL, a National Centre of Competence in Research funded by the Swiss National Science Foundation (grant number 205602).