



SAPIENZA
UNIVERSITÀ DI ROMA

“SAPIENZA” UNIVERSITÀ DI ROMA
INGEGNERIA DELL'INFORMAZIONE,
INFORMATICA E STATISTICA
DIPARTIMENTO DI INFORMATICA

Discrete Mathematics

TODO non so se scriverò qualcosa qui idk

Author
Alessio Bandiera

1 marzo 2024

Indice

Informazioni e Contatti	1
1 Number Theory	2
1.1 TODO	2
1.1.1 TODO	2

Informazioni e Contatti

Segnalazione errori ed eventuali migliorie:

Per segnalare eventuali errori e/o migliorie possibili, si prega di utilizzare il **sistema di Issues fornito da GitHub** all'interno della pagina della repository stessa contenente questi ed altri appunti (link fornito al di sotto), utilizzando uno dei template già forniti compilando direttamente i campi richiesti.

Gli appunti sono in continuo aggiornamento, pertanto, previa segnalazione, si prega di controllare se l'errore sia ancora presente nella versione più recente.

Licenza di distribuzione:

These documents are distributed under the [GNU Free Documentation License](#), a form of copyleft intended to be used on manuals, textbooks or other types of document in order to assure everyone the effective freedom to copy and redistribute it, with or without modifications, either commercially or non-commercially.

Contatti dell'autore e ulteriori link:

- Github: <https://github.com/aflaag>
- Email: alessio.bandiera02@gmail.com
- LinkedIn: [Alessio Bandiera](#)

1

Number Theory

1.1 TODO

1.1.1 TODO

Definizione 1.1.1.1: Peano axioms

The **Peano axioms** are 5 axioms which define the set \mathbb{N} of the **natural numbers**, and they are the following:

- i) $0 \in \mathbb{N}$
- ii) $\exists \text{succ} : \mathbb{N} \rightarrow \mathbb{N}$, or equivalently, $\forall x \in \mathbb{N} \quad \text{succ}(x) \in \mathbb{N}$
- iii) $\forall x, y \in \mathbb{N} \quad x \neq y \implies \text{succ}(x) \neq \text{succ}(y)$
- iv) $\nexists x \in \mathbb{N} \mid \text{succ}(x) = 0$
- v) $\forall S \subseteq \mathbb{N} \quad (0 \in S \wedge (\forall x \in S \quad \text{succ}(x) \in S)) \implies S = \mathbb{N}$

Principio 1.1.1.1: Induction principle

Let P be a property which is true for $n = 0$, thus $P(0)$ is true; also, for every $n \in \mathbb{N}$ we have that $P(n) \implies P(n + 1)$; then $P(n)$ is true for every $n \in \mathbb{N}$.

With symbols, using the formal logic notation, we have that

$$\frac{P(0) \quad P(n) \implies P(n + 1)}{\forall n \quad P(n)}$$

Osservazione 1.1.1.1: The fifth Peano axiom

Note that the fifth Peano axiom is equivalent to the induction principle, since, it states that for every subset S of \mathbb{N} containing 0 and closed under succ must be equal to \mathbb{N} itself.

Problema 1.1.1.1: Cardinality of the power set

Show that for every given set S such that $n := |S|$ it holds true that $|\mathcal{P}(S)| = 2^n$.

Dimostrazione. The statement will be shown by induction over n , the number of elements contained into S .

Caso base. $n = 0 \implies S = \emptyset \implies \mathcal{P}(S) = \mathcal{P}(\emptyset) = \{\emptyset\} \implies |\mathcal{P}(S)| = 1 = 2^0 = 2^n$.

Ipotesi induttiva. Assume that the statement is true for some fixed integer n .

Passo induttivo. It must be shown that the statement is true for $n + 1$ as well.

□

Definizione 1.1.1.2: Integers

TODO

Definizione 1.1.1.3: Divisor

TODO

Esempio 1.1.1.1 (Divisors). TODO

Proposizione 1.1.1.1: \mathbb{P} is infinite

There are infinitely many primes. With symbols

$$|\mathbb{P}| = +\infty$$

Dimostrazione. By way of contradiction, assume that \mathbb{P} is finite, thus

$$\exists n \in \mathbb{N} \mid \mathbb{P} = \{p_1, \dots, p_n\}$$

and let $x = p_1 \cdot \dots \cdot p_n$. Since $x \neq p_1, \dots, p_n$, then $x \notin \mathbb{P}$, so x is not a prime number; but x can't be divided by any of the p_1, \dots, p_n either, because the remainder will always be 1. This means that x is neither prime nor non-prime, which is a contradiction \nmid . □

Problema 1.1.1.2: $n^2 + n$ is even

Show that $\forall n \in \mathbb{N} \quad n^2 + n$ is an even number.

Dimostrazione. Note that $n^2 + n = n \cdot (n + 1)$, hence:

- if n is even, then

$$\exists k \in \mathbb{N} \mid n = 2k \implies n(n + 1) = 2k(2k + 1) = 4k^2 + 2k = 2(k^2 + k)$$

which is an even number;

- if n is odd, then

$$\exists k \in \mathbb{N} \mid n = 2k + 1 \implies n(n + 1) = (2k + 1)(2k + 2) = 4k^2 + 6k + 2 = 2(2k^2 + 3k + 1)$$

which is an even number.

□

Problema 1.1.1.3: $4n - 1$ is not prime

Show that there are infinitely many numbers of the form $4n - 1$ that are not prime.

Dimostrazione. Note that $\forall x^2 \in \mathbb{N} - \{0\} \quad 4x^2 - 1 = (2x + 1)(2x - 1)$ which is a proper factorization of $4x^2 - 1$, hence every perfect square yields a number of the form $4n - 1$ which is not a prime number. Note that the number of perfect squares is infinite since the set of perfect square has the same cardinality of \mathbb{N} since it's possible to construct a bijective function as follows:

$$f : \mathbb{N} \rightarrow \mathbb{N} : x \mapsto x^2$$

Also, note that this proof does not show *every non-prime number of the form $4n - 1$* , since that is outside the scope of the problem. □