





Post MI Prevalence: 90 to 94 female Baseline Lifestyle 0.065 Outreach Polypill 0.060 0.055 0.050 0.045 0.040 2040.0 2032.5 2025.0 2027.5 2022.5 2037.











Post MI Prevalence: 40 to 44 male Baseline 0.016 Lifestyle Outreach 0.015 Polypill 0.014 0.013 0.012 0.011 0.010 2040.0 -2022.5 2032.5 2025.0 2027.5 2037.


Post MI Prevalence: 45 to 49 male Baseline 0.023 Lifestyle Outreach 0.022 Polypill 0.021 0.020 0.019 0.018 0.017 2040.0 -2032.5 2025.0 2022.5 2027.5 2037.


Post MI Prevalence: 50 to 54 male Baseline 0.031 Lifestyle Outreach 0.030 Polypill 0.029 0.028 0.027 0.026 0.025 0.024 2040.0 -2032.5 2035.0 2025.0 2027.5 2030.0 2022.5 2037.


Post MI Prevalence: 55 to 59 male 0.042 Baseline Lifestyle 0.041 Outreach Polypill 0.040 0.039 0.038 0.037 0.036 2040.0 -2022.5 2032.5 2035.0 2025.0 2027.5 2037.














Post MI Prevalence: 90 to 94 male Baseline Lifestyle 0.11 Outreach Polypill 0.10 0.09 0.08 0.07 2040.0 -2025.0 2022.5 2027.5 2030.0 2035.0 2032.5

Post MI Prevalence: 95 plus male 0.11 Baseline Lifestyle 0.10 Outreach Polypill 0.09 0.08 0.07 0.06 0.05 2040.0 -2022.5 2025.0 2030.0 2027.5 2035.0 2032.