

1 Clonal Transmission of Emerging Novel *Plasmodium falciparum* Kelch13

2 Mutations and Increasing Complexity of Infection in Libreville, Gabon, 2021-2023

3 Jacques Mari Ndong Ngomo^{1,2,†}, Denise Patricia Mawili-Mboumba^{1,2}, Tobbi Ndong Mouity^{1,2},
4 Alec Leonetti³, Dimitri Moussavou Mabika^{1,2}, J'ose Coell Mihindou^{1,2}, Noé Patrick
5 Mbondoukwé^{1,2}, Jonathan J. Juliano^{4,5,6}, Jeffrey A. Bailey^{3,7,*}, Marielle Karine Bouyou-Akotet^{1,2,*},
6 Karamoko Niaré^{3,7,*,†}

7

8 ¹Département des Sciences fondamentales, service de Parasitologie-Mycologie et de
9 Médecine Tropicale, Université des Sciences de la Santé.

10 ²Centre de Recherche biomédicale en pathogènes Infectieux et Pathologies Associées
11 (CREIPA).

12 ³Department of Pathology and Laboratory Medicine, Brown University, Providence, RI,
13 USA.

14 ⁴Institute for Global Health and Infectious Diseases, University of North Carolina,
15 Chapel Hill, NC, USA.

16 ⁵Division of Infectious Diseases, School of Medicine, University of North Carolina,
17 Chapel Hill, NC, 27599, USA.

18 ⁶Department of Epidemiology, Gillings School of Global Public Health, University of
19 North Carolina, Chapel Hill, NC, 27599, USA.

20 ⁷Center for Computational Molecular Biology, Brown University, Providence, RI, USA.

21

22 *equal senior author contribution

23

24 [†]corresponding authors:

25 Karamoko Niare, PhD

26 Department of Pathology and Laboratory Medicine, Brown University, Providence, RI,
27 USA, karamoko_niare@brown.edu

28 Jacques Mari Ndong Ngomo

29 Département des Sciences fondamentales, service de Parasitologie-Mycologie et de
30 Médecine Tropicale, Université des Sciences de la Santé

31

32 ABSTRACT

33 Artemisinin partial resistance (ART-R) in *Plasmodium falciparum*, due to mutations in
34 the Kelch13 (K13) propeller domain, is spreading across Africa. However, data from
35 Central Africa remain sparse. This study performed molecular surveillance in a
36 peri-urban sentinel site in Libreville, Gabon, from 2021 to 2023 to assess emerging
37 resistance markers and parasite population dynamics. Febrile patients with confirmed *P.*
38 *falciparum* infection were enrolled at the Melen sentinel site. Dried blood spots were
39 collected and isolated DNA sequenced using molecular inversion probes (MIPs)
40 targeting drug resistance genes and genome-wide SNPs. We assessed the prevalence
41 of mutations in K13, DHFR, DHPS, CRT, and MDR1. Complexity of infection (COI) and
42 identity-by-descent (IBD) were used to evaluate transmission intensity and parasite
43 relatedness, respectively. Among 468 genotyped samples, no validated or candidate
44 K13 mutations were detected. However, 21 carried K13 mutations of unknown
45 significance, including E433D (n=14), Q613H (n=5), V520I, and V637I. Interestingly,
46 E433D prevalence rose from 0.7% in 2022 to 5.5% in 2023. Parasites with E433D or
47 Q613H showed significantly higher IBD than wild-type ($P<0.001$) and chains of clonal
48 transmission. Recent DHFR and DHPS mutations associated with higher-level
49 sulfadoxine-pyrimethamine resistance were at low prevalence. MDR1 Y184F increased
50 from 52.6% to 68.8%, while CRT K76T remained rare. IBD estimates support clonal
51 transmission of parasites carrying emerging K13 mutations, particularly E433D and
52 Q613H. In parallel, COI estimates increased over time, suggesting intensifying malaria
53 transmission, potentially with a seasonal component. These findings highlight the need
54 for expanded genomic surveillance and functional validation of these novel mutations to
55 inform malaria control strategies in Gabon and Central Africa.

56 Keywords:

57 Malaria, *Plasmodium falciparum*, Kelch13, artemisinin resistance, Gabon, molecular
58 surveillance, drug resistance, identity-by-descent, complexity of infection

59

60 INTRODUCTION

61

62 Malaria continues to pose a significant global health challenge, particularly in
63 sub-Saharan Africa, where it causes high rates of illness and death (1). Despite
64 decades of intervention efforts, progress has been inconsistent, with only minimal
65 declines in malaria burden across the region in recent years. Several factors threaten
66 the effectiveness of malaria control, including resistance to drugs, diagnostics, and
67 insecticides, as well as the spread of the invasive mosquito species *Anopheles*
68 *stephensi* (2–6).(7–9). Another major gap is the lack of vaccines that provide
69 long-lasting immunity (10,11).

70 Artemisinin partial resistance (ART-R) has been linked to partial loss-of-function
71 mutations in the *Plasmodium falciparum* Kelch13 (K13) propeller domain (4). Alarminglly,
72 multiple ART-R mutations have emerged and are increasingly reported across the Great
73 Rift Valley and into Southern Africa (6,12–18). World Health Organization
74 (WHO)-validated and candidate K13 mutations such as P441L, C469Y, C469F, R561H,
75 R622I, and A675V have been identified in these regions and appear to be increasing in
76 frequency and spreading across borders (6,12–17). These appear to be de novo
77 mutations that originated in Africa (6,12,15). Given the full mutational landscape that
78 can impart ART-R has not been fully characterized, it is important to continue to monitor
79 for the emergence of novel mutations that may impart ART-R in addition to previously
80 validated mutations.

81 Artesunate-amodiaquine (ASAQ) and artemether-lumefantrine (AL) were the first
82 artemisinin-based combination therapies (ACT) used as first-line treatment for
83 uncomplicated malaria in Gabon, but recent policy changes in 2024 now recommend
84 ASAQ, AL, artesunate-mefloquine (ASMQ) and dihydroartemisinin-piperaquine (DP).
85 Sulfadoxine-pyrimethamine (SP) is used for intermittent preventive treatment in
86 pregnancy, while severe malaria is treated with injectable artemisinin derivatives or
87 quinine. The national malaria control program, in partnership with the Université des
88 Sciences de la Santé, provides free malaria diagnosis at six sentinel sites and collects
89 blood samples for molecular surveillance. However, to date, molecular drug resistance
90 surveillance in Gabon has relied on less sensitive restriction fragment length

polymorphism (RFLP) methods and was limited to a few loci analyzed by study according to a recent review (19). To date, no validated or candidate ART-R mutations have been reported in Gabon (20,21), and ACT efficacy remains high, with cure rates above 95% for AL and ASAQ (22,23).

To support malaria control, next-generation sequencing (NGS) offers a high-throughput, and detailed approach to understand parasite mutations, their origins, and their spread (24,25). Molecular surveillance also provides insights into malaria transmission dynamics, parasite relatedness, and selective pressures on the parasite population (26–28). This type of genomic surveillance has not been recently conducted in Gabon. Therefore, we conducted a pilot genomic surveillance project using samples collected between 2021 and 2023 from a peri-urban sentinel site in Libreville. We employed molecular inversion probe (MIP) capture and Illumina deep sequencing to assess antimalarial drug resistance, infection complexity, and parasite relatedness. Our analysis revealed non-synonymous mutations in K13, including one mutation that appears to be increasing in frequency over time.

RESULTS


We sequenced 537 samples with parasitemia ≥ 500 parasites/ μ l (by microscopy) collected between 2021-2023 in Melen, a peri-urban sentinel site in Libreville. We successfully genotyped 465 samples to sufficient depth, defined as unique molecular identifier (UMI) counts per MIP of ≥ 5 , for variant calling using DR23KE and IBC2CORE panels (29–31). We analyzed 77 samples from 2021, 169 from 2022 and 219 from 2023 for drug resistance mutations, relatedness, and complexity of infection (COI).

Uncharacterized non-synonymous mutations in K13 BTB/POZ and propeller domains are emerging in Gabon

While no WHO validated or candidate K13 mutations were detected, we found one mutation in the BTB/POZ domain (E433D) and three uncharacterized mutations within the propeller domains (V520I, Q613H and V637I) (**Fig.1A**)(32). The most common mutation was E433D – first being seen in 2022 in one (0.6%) of 154 samples. The mutation prevalence increased to 6.3% (13/207) in 2023, the last year surveyed. The

Q613H mutation was detected in 5 isolates (5/204, 2.4%) in 2023. The V520I (1/67) and V637I (1/79) only occurred in only one participant each in 2021. Many (10/21) of these propeller domain and BTB/POZ mutations were either the only or the dominant allele found within an infection and there was no evidence of multiple mutations within a haplotype (**Fig1.B**). The prevalence of the A578S, which is a wide-spread mutation not associated with ART-R, was low with 1.2% (1/81, CI) in 2021 and 1.4% (3/221) in 2023. Outside the BTB/POZ and propeller domain, ten polymorphisms were detected (**Table S1**).

To understand the relatedness of parasites carrying the same K13 mutation, we performed pairwise identity-by-descent (IBD) analysis between them using IBC2CORE data which targeted SNPs outside the drug resistance genes (29). The mean IBD fractions between sample pairs with E433D (mean IBD=0.42) and Q613H (mean IBD=0.2) were significantly higher ($P<0.0001$, **Fig.1C**) compared to wildtype (WT) sample pairs (mean IBD=0.04) and A578S sample pairs (mean IBD=0.03). IBD network analysis showed clustering of samples carrying E433D and Q613H at IBD>0.9, consistent with clonal chains of transmission (**Fig1.D**).



136

Figure 1: Prevalence of non-synonymous mutations detected in the *k13* gene and parasite relatedness. **A)** Prevalence of mutations by year. Sample sizes are shown on top of the barplot. **B)** Fraction of unique molecular identifiers (UMIs) supporting each mutation within-sample defined as within-sample allele frequency. The minimum total UMI count by mutation site was 17. **C)** Relatedness between isolates carrying the same mutations versus wildtype (WT) alleles based on IBD. **D)** IBD networks showing clusters of samples carrying E433D and Q613H despite at IBD \geq 0.90. IBD analysis was performed with all samples (n=465). Clusters of samples carrying the same K13 mutations are indicated by dotted lines.

146

147 Prevalence of most antimalarial drug resistance mutations remains stable over 148 time


149 We analyzed SP resistance markers and found that dihydrofolate reductase (DHFR)
150 mutations N51I, C59R, and S108N were fixed, with prevalence consistently above 97%
151 over time (**Fig. 2A & B**). The I164L mutation in DHFR was not detected. Similarly, the
152 dihydropteroate synthase (DHPS) A437G mutation was fixed (>97% prevalence across
153 all three years). Other common DHPS mutations included S436A and K540E (**Fig. 2B**).
154 While K540E remained stable at ~5%, S436A increased from 7.6% (5/66) in 2021 to
155 18.2% (28/154) in 2022 and 23.1% (49/212) in 2023. A581G and A613S were detected
156 only in 2022 and 2023, with prevalence below 1% (2/217) and 3% (4/156), respectively.
157 The triple DHFR mutation combination (N51I+C59R+S108N, or IRN) was present in
158 over 90% of samples each year (**Table I**). The quintuple mutation IRN+A437G+K540E
159 (IRN+GE) declined from 9.7% (6/62) in 2021 to 5.3% (11/209) in 2023. No sextuple
160 mutations were detected.

161 To assess resistance to partner drugs amodiaquine and lumefantrine, we examined the
162 chloroquine resistance transporter (CRT) and multidrug resistance 1 (MDR1) mutations,
163 where these drugs select for mutations and wildtype, respectively. CRT mutations were
164 rare, with the highest prevalence of the K76T mutation at 6.5% (9/139) in 2022 (**Fig.**
165 **2C**). Among MDR1 mutations, N86Y was found only in 2021 and 2023 at <3%, while
166 Y184F increased from 52.6% (40/76) in 2021 to 68.8% (148/215) in 2023 (**Fig. 3C**).
167 D650N and N652D remained stable between 34.6% and 47.7%. Rare MDR1 variants
168 are listed in **Table S1**. No gene duplications were observed (**Fig. S1**).

169 We also investigated additional resistance-associated loci relevant in Africa, including
170 including putative amino acid transporter (AAT1), plasmepsin 2 and 3 (PM2 and PM3),
171 cysteine proteinase falcipain 2a (FP2a), GTP cyclohydrolase 1 (GCH1), and coronin
172 (CORONIN) (**Table S2**). In AAT1, the S258L mutation was nearly fixed (96.7–100%),
173 while F313S was absent. No duplications of *pm2* or *pm3*—linked to piperaquine
174 resistance—were found. Although PM3 SNPs were rare, PM2 Q442H was frequent
175 (86.7–90.6%). In FP2A, only Q414H was detected (59.3–66.4%). GCH1 mutations were

176 rare, with no evidence of copy number variation. In coronin, mutations G50E, R100K,
177 and E107V were absent, while S183G was the most common (73.3–83.3%).

178



179

180 **Figure2 : Prevalence of common antimalarial drug resistance mutations from**
181 **2021 to 2023. Prevalence of DHFR (A), DHPS (B), CRT (C) and MDR1 (D) mutations,**
182 **respectively (n=465). For MDR1, only major mutations, N86Y, Y184F, D650N and**

183 N652D, are shown. The rest of mutations detected are shown in supplementary Table
184 S2.

185

186 **Table 1: Prevalence of DHFR triple mutations and DHFR and DHPS quintuple**
187 **mutations by year.**

Haplotype	Frequency % (n/N)			Number of Mutations
	2021	2022	2023	
IRN	90.3 (56/62)	92.9 (145/156)	94.7 (198/209)	Triple
IRN+GE	9.7 (6/62)	7.1 (11/156)	5.3 (11/209)	Quintuple

188

189 **Complexity of infection varies by year**

190


191 To assess recent changes in malaria transmission intensity in western Gabon, we
192 analyzed the distribution of Complexity of Infection (COI) over time at our sentinel site
193 (**Fig. 3**). Overall, COI showed a significant upward trend across the study period, with a
194 strong positive correlation with time ($P < 0.0009$, linear regression). Although the
195 increase from 2022 to 2023 approached significance, it still suggests a potential
196 intensification of transmission during this period (**Fig. 3A**).

197

198 Given that malaria transmission in Gabon occurs year-round (33–35), we further
199 investigated whether the observed rise in COI was seasonally driven. We calculated the
200 proportion of polygenomic infections (samples with $COI > 1$) in quarterly intervals and
201 compared these proportions across the three survey years. From January to
202 September, polygenomic infections generally increased between 2021 and 2023 (**Fig.**

203 **3B**), with a statistically significant rise during January–March ($P = 0.043$, linear
204 regression). However, from October to December, the proportions remained relatively
205 stable across years.

206



207


208 **Figure 3: Temporal trend of complexity of infection (COI).** **A)** continuous COI
209 distribution by year. Significant pairwise comparisons of COI between years are
210 indicated by the P values displayed (Wilcoxon Test). The temporal increase of COI
211 between 2021–2023 was significant ($P < 0.0009$) using a linear regression model. Dots

represent samples and sample size is indicated below each box. **B)** Comparison of proportions of polygenomic infections (fraction of samples with COI>1) between 2021–2023 after every three-month period. Significance of the linear regression analysis between proportion of polygenomic infections and year is indicated on top of the bars as well as polygenomic sample count and sample size.

Genetic relatedness between samples suggests some clonal transmission of parasites from the same year and those with E433D and Q613H mutations

To assess the genetic relatedness of malaria parasites within and across years, we analyzed identity-by-descent (IBD) fractions across all 71,610 sample pairs, which provides insights into parasite interrelatedness (36). This analysis was based on 669 genome-wide SNPs generated using the IBC2CORE panel. Within-year pairwise IBD fractions—representing the proportion of the genome with shared IBD between sample pairs—increased steadily from 2021 to 2023 (**Fig. 4A**). To further explore population structure, we conducted an IBD network analysis at varying thresholds of relatedness. At $IBD \geq 0.50$, the majority of isolates formed a single interconnected network, regardless of the year of collection (**Fig. 4B**). However, when the threshold was raised to $IBD \geq 0.90$, clustering became more year-specific, revealing multiple groups of highly related (near-clonal) parasites, mostly from the same year (**Fig. 4C**). A few clusters spanned multiple years, including those containing E433D mutant samples.

We also performed t-distributed Stochastic Neighbor Embedding (t-SNE) analysis using the same SNP dataset to investigate potential clustering by year and/or K13 mutation status. While most samples did not form distinct clusters, several year-specific clusters were observed (**Fig. 4D**). Additionally, some clusters included samples from different years, such as a group of E433D mutants. Q613H isolates also formed a distinct cluster, whereas A578S mutants were dispersed among the unclustered samples (**Fig. 4D**).



239

Figure 4: Relatedness between samples within and across years. **A)** Comparison of pairwise IBD fractions across years. IBD fractions between isolate pairs significantly over time (* indicates significance). **B)** IBD network analysis at IBD ≥ 0.50 showing the vast majority of samples clustering into a single network of relatives. **C)** IBD network analysis at IBD ≥ 0.9 showing samples clustering by year. Each dot represents an isolate and color codes for sample collection year. All the samples were included in IBD analysis (n=465). **D)** Population structure partially defined by sample collection year and K13 mutation (n=465). Clusters of samples carrying the same K13 mutations are circled by colored dotted lines.

249

250 DISCUSSION

251 This three-year pilot genomic surveillance project in Gabon identified emerging
252 mutations in the K13 gene, particularly E433D in the BTB/POZ domain and Q613H in
253 the propeller domain. While these are not validated or candidate ART-R mutations, their
254 appearance is concerning. Particularly, E433D showed a notable increase in frequency
255 between 2022 and 2023, while Q613H appeared in the most recent year. Both showed
256 evidence of clonal chains of transmission. These findings, along with a rising COI
257 suggestive of increasing transmission and spreading markers MDR1 N86 and 184F
258 associated with decreased sensitivity to lumefantrine, highlight the importance of
259 continued monitoring and further investigation into the potential implications for malaria
260 control in the region.

261 The E433D substitution appears to be a newly emerging mutation in Gabon, first
262 detected in 2022 and rising to a prevalence of 6% by 2023. This ten-fold increase over
263 a single year suggests strong selective pressure, potentially driven by widespread use
264 of ACTs. The mutation was not previously reported in Gabon or neighboring countries,
265 indicating a local emergence (19–21,37). Genetic analyses revealed that E433D was
266 primarily found in mixed infections and showed high IBD among carriers, suggesting
267 clonal expansion. In addition to E433D, three propeller domain mutations—V520I,
268 Q613H, and V637I—were identified. Among these, Q613H was the most prevalent
269 (2.4%) and may warrant close monitoring. While V520I and V637I were rare and only
270 detected in 2021, Q613H showed signs of clonal transmission similar to E433D.
271 Although different amino acid substitutions at positions 613 and 637 have been reported
272 in other African countries, these specific variants are novel in Gabon (37–41).

273 The study site, a peri-urban area of Libreville, is characterized by mesoendemic and
274 unstable malaria transmission, which presents a unique epidemiological setting for the
275 emergence and spread of drug-resistant parasites (33,34). Historical data show
276 fluctuating malaria prevalence in this region, with a notable decline from 34% in 2005 to
277 25% in 2011, followed by a resurgence to 31% in 2021, indicating a recent upward trend

(33,34). This pattern of instability may reflect changes in vector control efforts, environmental conditions, and healthcare access over time. Such fluctuations can lead to inconsistent levels of population immunity, particularly in children and other vulnerable groups, which in turn may increase susceptibility to infection and facilitate the selection of resistant strains (42,43). The combination of unstable transmission and high drug pressure—driven by widespread ACT use and easy access to antimalarials—creates favorable conditions for resistance to emerge (12,42,43). Informal drug markets and self-medication practices may contribute to suboptimal treatment, intensifying selective pressure (44,45).

Surveillance also revealed concerning trends in resistance to partner drugs used in ACTs, as well as SP. The MDR1 N86 wild-type allele was found to be fixed in the population, a shift that likely reflects long-term selection pressure from the widespread use of AL, the most commonly used ACT in Gabon (46). In parallel, the Y184N mutation in MDR1 showed a rising prevalence, which may further indicate adaptation to lumefantrine exposure (47–49). Although the functional implications of Y184N remain under investigation, its increasing frequency suggests it may play a role in modulating drug response or compensating for fitness costs associated with other resistance mutations. In contrast, the CRT K76T mutation—associated with resistance to amodiaquine (48)—remained rare, suggesting that ASAQ is not widely used or has not exerted strong selective pressure in this setting. This supports the continued efficacy of ASAQ as an effective treatment option in Gabon. SP is primarily used in pregnant women for intermittent preventive treatment although trimethoprim and sulfamethoxazole is preferred in HIV-infected women to prevent opportunistic infections. Markers of SP resistance were widespread, with fixation of DHFR mutations (N51I, C59R, S108N) and high prevalence of DHPS A437G. Notably, the DHPS S436A mutation associated with resistance to sulfadoxine showed a rising trend from 7.6% in 2021 to 23.1% in 2023, indicating increasing SP pressure. However, key late-stage mutations associated with high-level SP resistance—such as DHFR I164L and DHPS K540E, A581G, and A613S (50–61)—remained rare or absent. This suggests that, despite the widespread presence of early SP resistance markers, SP may still retain some utility in Gabon, particularly for intermittent preventive treatment in pregnancy.

309 These results are also in keeping with well controlled use of SP, which is rarely utilized
310 for self-medication in Gabon.

311 This study was limited to a single geographic site in northwestern Gabon, which may
312 not fully represent the genetic diversity or resistance dynamics occurring in other
313 regions of the country or Central Africa more broadly. Sample collection was mainly
314 disrupted by the COVID-19 pandemic, leading to relatively small sample sizes,
315 particularly in 2021; which may limit the statistical power to detect low-frequency
316 mutations or subtle trends over time. While identity-by-descent and population structure
317 analyses suggested clonal transmission of key mutations such as E433D and Q613H,
318 these findings would benefit from confirmation through whole genome sequencing to
319 better understand the broader genomic context and evolutionary history of these
320 variants.

321 In conclusion, this pilot study provides important early insights into the evolving genetic
322 landscape of malaria parasites in Gabon. Although none of the K13 mutations
323 identified—E433D, Q613H, V520I, and V637I—have been validated as markers of
324 artemisinin resistance, the prevalence of E433D and Q613H with evidence of clonal
325 transmission warrant further examination. Functional studies, such as
326 CRISPR-Cas9–based gene editing and *in vitro* drug susceptibility assays, are needed to
327 determine whether these mutations confer reduced artemisinin sensitivity. Together,
328 these findings emphasize the need for expanded genomic surveillance, phenotypic
329 validation, and integrated monitoring of both artemisinin and partner drug resistance to
330 inform malaria control strategies in Central Africa.

331

332 MATERIALS AND METHODS

333 Study design

334 Surveillance was conducted at the Centre Hospitalier Régional de l'Estuaire in Melen,
335 located in a peri-urban area 11 km from Libreville, the capital city of Gabon. This is one
336 of six sentinel sites supported by the national malaria control program to perform

337 malaria epidemiological surveillance and therapeutic efficacy studies. Between
338 2021-2023, symptomatic ($\geq 37.5^{\circ}\text{C}$) children and adolescents from paediatric wards
339 were screened by microscopy and persons with microscopically confirmed *P. falciparum*
340 infection were enrolled after obtaining informed consent from guardians. Clinical and
341 sociodemographic data were recorded. Dried blood spots (DBS) on filter paper were
342 collected from finger prick ($\sim 40\ \mu\text{L}$) for molecular studies. Each sample was stored in a
343 ziplock bag containing two desiccants and kept at 4°C in a fridge before punching and
344 shipping to Brown University for sequencing. *P. falciparum*-infected samples with at
345 least 1,000 parasites/ μL were included in the study ($n=537$). Ethical approval was
346 obtained from the Scientific Committee of Comité National de Pilotage de la Riposte
347 contre le Coronavirus-Gabon (reference: 0072/P/COPIL-CS-COVID-19). The work at
348 Brown on deidentified samples was considered nonhuman subjects.

349 **Sample Processing and MIP genotyping.**

350 DNA was extracted from DBS using a Chelex-Tween protocol (30,31) before MIP
351 capture. MIP capture and sequencing was conducted as previously described (30,31).
352 Briefly, we combined two panels, DR23KE, targeting drug resistance genes, and
353 IBC2CORE, capturing common SNPs across the genome for relatedness and COI
354 analyses. This MIP pool was used to capture genomic regions of interest into
355 circularized DNA, followed by digestion of single stranded/linearized DNA using
356 exonucleases, to enrich captured DNA Sample barcoding, unique molecular identifier,
357 and library preparation was conducted by PCR amplification to create final Illumina
358 libraries that are pooled and sequenced on an Illumina Nextseq 550 at Brown
359 University. Laboratory strains 3D7 and Dd2 were used as controls.

360 We used the MIPTools (version4.0) (<https://github.com/bailey-lab/MIPTools>) for the data
361 analysis, including read processing and mapping and variant calling using Freebayes
362 (version1.3.5) (Garrison & Marth, 2012). Samples with median unique molecular
363 identifier (UMI) count < 10 across probes were repooled and resequenced to improve
364 read depth. We used the MIPlicorn R package (<https://github.com/bailey-lab/miplicorn>)
365 to process the output of the MIPTools analysis before calculating the prevalence of
366 mutations found in drug resistance genes. Copy number of *mdr1*, *pfpm2* and *pfpm3*

was measured by calculating read depth fold-change per MIP, based on UMI counts, relative to other non-copy number regions captured by the DR23KE panel as previously described (62).

Complexity of infection analysis.

The SNP genotypes generated with IBC2CORE MIPs was used for the discrete COI calculation using THE REAL McCOIL package (version 2.0) (63). We set the total number of Markov chain Monte Carlo and burn-in iterations to 2000 with 500, respectively. Only targeted SNPs for which the MIPs were designed with site and sample missingness of <10% and <20%, respectively, were used. The proportion of polygenomic infections, samples with discrete COI > 1 over total sample size, was calculated by year and for three month intervals. The continuous COI was estimated using the *coiaf* R package based on the same filtered SNPs (64). This COI calculation was corrected for the variation of site-level read depth.

Identity-by-descent analysis.

We used the *hmmIBD* package (v3.0) to compute IBD fractions between sample pairs based on SNPs obtained with IBC2CORE MIPs. The maximum number of fit iterations was set to 5 to detect recent big IBD blocks. A theoretical genotype error rate of 0.1% was adopted and clustering and visualization were performed using the *igraph* R package.

Authors contribution

NNJM, BAKMK, JJJ, JAB and KN designed and conducted the study. KN analyzed data and interpreted results. JJJ and JAB supervised the analysis and interpretation of the data. JJJ and JAB obtained primary funding for the molecular work. MMD, MJC, MNP, KL, MMDP, NNJM and BAKMK performed sample collection and shared metadata. NNJM led the field work in Gabon. KN and AL performed DNA isolation and sequencing. KN wrote the primary draft of the manuscript. All authors contributed to the writing of the manuscript.

Conflicts of Interest:

The authors declare no competing interests.

396 Acknowledgements -

397 The authors are grateful to Malaria Research and Reference Reagent Resource Center
398 (MR4) for sharing laboratory strain DNA controls. The authors thank study participants
399 and caregivers at the Melen sentinel site.

400 Data Availability

401 Parasite sequences that support the findings of this study have been deposited in the
402 Sequence Read Archive repository under the accession numbers SAMN50482924 to
403 SAMN50483388 at <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1302560>

404 **Funding:** This project was supported by the National Institutes for Allergy and
405 Infectious Diseases (R01AI156267 to JAB and JJJ; K24AI134990 to JJJ).

406 References

- 407 1. World Health Organization. World malaria report 2024 [Internet]. World Health
408 Organization; 2024 [cited 2025 May 1]. Available from:
409 <https://www.who.int/publications/i/item/9789240104440>
- 410 2. World Health Organization. World malaria report 2023. World Health Organization;
411 2023.
- 412 3. Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, et al. Artemisinin
413 resistance in *Plasmodium falciparum* malaria. *N Engl J Med*. 2009;361(5):455–67.
- 414 4. Arieu F, Witkowski B, Amaratunga C, Beghain J, Langlois AC, Khim N, et al. A
415 molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature*.
416 2014 Jan 2;505(7481):50–5.
- 417 5. Ashley E a., Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, et al. Spread
418 of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med*.
419 2014;371(5):411–23.
- 420 6. Uwimana A, Legrand E, Stokes BH, Ndikumana JLM, Warsame M, Umulisa N, et
421 al. Emergence and clonal expansion of in vitro artemisinin-resistant *Plasmodium*
422 *falciparum* kelch13 R561H mutant parasites in Rwanda. *Nat Med*. 2020
423 Oct;26(10):1602–8.
- 424 7. Balkew M, Mumba P, Dengela D, Yohannes G, Getachew D, Yared S, et al.
425 Geographical distribution of *Anopheles stephensi* in eastern Ethiopia. *Parasit*
426 *Vectors*. 2020 Jan 20;13(1):35.
- 427 8. Seyfarth M, Khairah BA, Abdi AA, Bouh SM, Faulde MK. Five years following first
428 detection of *Anopheles stephensi* (Diptera: Culicidae) in Djibouti, Horn of Africa:
429 populations established-malaria emerging. *Parasitol Res*. 2019 Mar;118(3):725–32.

- 430 9. Ranson H, Lissenden N. Insecticide Resistance in African Anopheles Mosquitoes:
431 A Worsening Situation that Needs Urgent Action to Maintain Malaria Control.
432 Trends Parasitol. 2016 Mar;32(3):187–96.
- 433 10. RTS,S Clinical Trials Partnership. Efficacy and safety of RTS,S/AS01 malaria
434 vaccine with or without a booster dose in infants and children in Africa: final results
435 of a phase 3, individually randomised, controlled trial. Lancet. 2015 Jul
436 4;386(9988):31–45.
- 437 11. RTSS Clinical Trials Partnership. Efficacy and Safety of the RTS,S/AS01 Malaria
438 Vaccine during 18 Months after Vaccination: A Phase 3 Randomized, Controlled
439 Trial in Children and Young Infants at 11 African Sites. Krishna S, editor. PLoS Med.
440 2014 Jul 29;11(7):e1001685.
- 441 12. Conrad MD, Asua V, Garg S, Giesbrecht D, Niaré K, Smith S, et al. Evolution of
442 Partial Resistance to Artemisinins in Malaria Parasites in Uganda. N Engl J Med.
443 2023 Aug 24;389(8):722–32.
- 444 13. Young NW, Gashema P, Giesbrecht D, Munyaneza T, Maisha F, Mwebembezi F, et
445 al. High frequency of artemisinin partial resistance mutations in the great lake
446 region revealed through rapid pooled deep sequencing [Internet]. bioRxiv. 2024.
447 Available from:
448 <https://www.medrxiv.org/content/10.1101/2024.04.29.24306442v1.abstract>
- 449 14. Ishengoma DS, Mandara CI, Bakari C, Fola AA, Madebe RA, Seth MD, et al.
450 Evidence of artemisinin partial resistance in North-western Tanzania: clinical and
451 drug resistance markers study. medRxiv [Internet]. 2024 Feb 1 [cited 2024 Oct 3];
452 Available from: <https://pubmed.ncbi.nlm.nih.gov/38352311/>
- 453 15. Juliano JJ, Giesbrecht DJ, Simkin A, Fola AA, Lyimo BM, Pereus D, et al. Country
454 wide surveillance reveals prevalent artemisinin partial resistance mutations with
455 evidence for multiple origins and expansion of high level
456 sulfadoxine-pyrimethamine resistance mutations in northwest Tanzania. medRxiv
457 [Internet]. 2023 Nov 30; Available from:
458 <http://dx.doi.org/10.1101/2023.11.07.23298207>
- 459 16. Mihreteab S, Platon L, Berhane A, Stokes BH, Warsame M, Campagne P, et al.
460 Increasing prevalence of artemisinin-resistant HRP2-negative malaria in Eritrea. N
461 Engl J Med. 2023 Sep 28;389(13):1191–202.
- 462 17. Fola AA, Feleke SM, Mohammed H, Brhane BG, Hennelly CM, Assefa A, et al.
463 Plasmodium falciparum resistant to artemisinin and diagnostics have emerged in
464 Ethiopia. Nat Microbiol. 2023 Oct;8(10):1911–9.
- 465 18. Martin AC, Sadler JM, Simkin A, Musonda M, Katowa B, Matoba J, et al.
466 Emergence and rising prevalence of artemisinin partial resistance marker Kelch13
467 P441L in a low malaria transmission setting in southern Zambia [Internet]. medRxiv.
468 2025 [cited 2025 May 2]. p. 2025.01.02.24319706. Available from:

- 469 <https://www.medrxiv.org/content/10.1101/2025.01.02.24319706v1.abstract>
- 470 19. Sima-Biyang YV, Ontoua SS, Longo-Pendy NM, Mbou-Boutambe C,
471 Makouloutou-Nzassi P, Moussadji CK, et al. Epidemiology of malaria in Gabon: A
472 systematic review and meta-analysis from 1980 to 2023. *J Infect Public Health*.
473 2024 Jul;17(7):102459.
- 474 20. Kamau E, Campino S, Amenga-Etego L, Drury E, Ishengoma D, Johnson K, et al.
475 K13-propeller polymorphisms in plasmodium falciparum parasites from
476 sub-saharan Africa. *J Infect Dis*. 2015 Apr 15;211(8):1352–5.
- 477 21. Dinzouna-Boutamba SD, Iroungou BA, Akombi FL, Yacka-Mouele L, Moon Z, Aung
478 JM, et al. Assessment of genetic polymorphisms associated with malaria antifolate
479 resistance among the population of Libreville, Gabon. *Malar J*. 2023 Jun
480 14;22(1):183.
- 481 22. Adegbite BR, Edoa JR, Honkpehedji YJ, Zinsou FJ, Dejon-Agobe JC,
482 Mbong-Ngwese M, et al. Monitoring of efficacy, tolerability and safety of
483 artemether–lumefantrine and artesunate–amodiaquine for the treatment of
484 uncomplicated Plasmodium falciparum malaria in Lambaréné, Gabon: an
485 open-label clinical trial. *Malar J*. 2019 Dec 16;18(1):424.
- 486 23. Ndong Ngomo JM, Ondzagha Megnie GJ, Moutombi Ditombi B, Koumba Lengongo
487 JV, M'Bondoukwé NP, Offouga CL, et al. Persistence of high in vivo efficacy and
488 safety of artesunate-amodiaquine and artemether-lumefantrine as the first- and
489 second-line treatments for uncomplicated Plasmodium falciparum malaria 10 years
490 after their implementation in Gabon. *Acta Parasitol*. 2019 Dec;64(4):898–902.
- 491 24. Volkman SK, Neafsey DE, Schaffner SF, Park DJ, Wirth DF. Harnessing genomics
492 and genome biology to understand malaria biology. *Nat Rev Genet*. 2012 Apr
493 12;13(5):315–28.
- 494 25. Golumbeanu M, Edi CAV, Hetzel MW, Koepfli C, Nsanzabana C. Bridging the gap
495 from molecular surveillance to programmatic decisions for malaria control and
496 elimination. *Am J Trop Med Hyg*. 2025 Jan 7;112(1_Suppl):35–47.
- 497 26. Schaffner SF, Badiane A, Khorgade A, Ndiop M, Gomis J, Wong W, et al. Malaria
498 surveillance reveals parasite relatedness, signatures of selection, and correlates of
499 transmission across Senegal. *Nat Commun*. 2023 Nov 10;14(1):7268.
- 500 27. Wesolowski A, Taylor AR, Chang HH, Verity R, Tessema S, Bailey JA, et al.
501 Mapping malaria by combining parasite genomic and epidemiologic data. *BMC*
502 *Med*. 2018 Oct 18;16(1):190.
- 503 28. Tessema S, Wesolowski A, Chen A, Murphy M, Wilhelm J, Mupiri AR, et al. Using
504 parasite genetic and human mobility data to infer local and cross-border malaria
505 connectivity in Southern Africa. *Elife [Internet]*. 2019 Apr 2 [cited 2025 Jul 3];8.
506 Available from: <http://dx.doi.org/10.7554/eLife.43510>

- 507 29. Niaré K, Crudale R, Fola AA, Wernsman Young N, Asua V, Conrad M, et al. Highly
508 multiplex molecular inversion probe panel in *Plasmodium falciparum* targeting
509 common SNPs approximates whole genome sequencing assessments for selection
510 and relatedness [Internet]. medRxiv. 2025. p. 2025.03.07.25323597. Available
511 from: <https://www.medrxiv.org/content/10.1101/2025.03.07.25323597v1>
- 512 30. Verity R, Aydemir O, Brazeau NF, Watson OJ, Hathaway NJ, Mwandagaliwa MK,
513 et al. The impact of antimalarial resistance on the genetic structure of *Plasmodium*
514 *falciparum* in the DRC. Nat Commun. 2020 Apr 30;11(1):2107.
- 515 31. Aydemir O, Janko M, Hathaway NJ, Verity R, Mwandagaliwa MK, Tshetu AK, et al.
516 Drug-resistance and population structure of *Plasmodium falciparum* across the
517 Democratic Republic of Congo using high-throughput molecular inversion probes. J
518 Infect Dis. 2018 Aug 14;218(6):946–55.
- 519 32. Malaria: Artemisinin partial resistance [Internet]. [cited 2025 Jun 18]. Available from:
520 <https://www.who.int/news-room/questions-and-answers/item/artemisinin-resistance>
- 521 33. Mawili-Mboumba DP, Bouyou Akotet MK, Kendjo E, Nzamba J, Medang MO, Mbina
522 JRM, et al. Increase in malaria prevalence and age of at risk population in different
523 areas of Gabon. Malar J. 2013 Jan 2;12(1):3.
- 524 34. Moutombi Ditombi BC, Pongui Ngondza B, Manomba Boulingui C, Mbang Nguema
525 OA, Ndong Ngomo JM, M'Bondoukwé NP, et al. Malaria and COVID-19 prevalence
526 in a population of febrile children and adolescents living in Libreville. S Afr J Infect
527 Dis. 2022 Oct 26;37(1):459.
- 528 35. Lendongo-Wombo JB, Oyegue-Liabagui SL, Biteghe-Bi-Essone JC, Ngoungou EB,
529 Lekana-Douki JB. Epidemiology of malaria from 2019 to 2021 in the southeastern
530 city of Franceville, Gabon. BMC Public Health. 2022 Dec 10;22(1):2313.
- 531 36. Taylor AR, Schaffner SF, Cerqueira GC, Nkhoma SC, Anderson TJC, Sriprawat K,
532 et al. Quantifying connectivity between local *Plasmodium falciparum* malaria
533 parasite populations using identity by descent. Didelot X, editor. PLoS Genet. 2017
534 Oct 27;13(10):e1007065.
- 535 37. Milong Melong CS, Peloewetse E, Russo G, Tamgue O, Tchoumboungang F,
536 Paganotti GM. An overview of artemisinin-resistant malaria and associated Pfk13
537 gene mutations in Central Africa. Parasitol Res. 2024 Jul 18;123(7):277.
- 538 38. Huang F, Yan H, Xue JB, Cui YW, Zhou SS, Xia ZG, et al. Molecular surveillance of
539 pfcr, pfmdr1 and pfk13-propeller mutations in *Plasmodium falciparum* isolates
540 imported from Africa to China. Malar J. 2021 Feb 6;20(1):73.
- 541 39. Ménard D, Khim N, Beghain J, Adegnik AA, Shafiul-Alam M, Amodu O, et al. A
542 worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. N Engl J
543 Med. 2016 Jun 23;374(25):2453–64.

- 544 40. Taylor SM, Parobek CM, DeConti DK, Kayentao K, Coulibaly SO, Greenwood BM,
545 et al. Absence of putative artemisinin resistance mutations among *Plasmodium*
546 *falciparum* in Sub-Saharan Africa: a molecular epidemiologic study. *J Infect Dis.*
547 2015 Mar 1;211(5):680–8.
- 548 41. Schmedes SE, Patel D, Dhal S, Kelley J, Svigel SS, Dimbu PR, et al. *Plasmodium*
549 *falciparum* kelch 13 Mutations, 9 Countries in Africa, 2014-2018. *Emerg Infect Dis.*
550 2021 Jul;27(7):1902–8.
- 551 42. Kanya MR, Nankabirwa JI, Arinaitwe E, Rek J, Zedi M, Maiteki-Sebuguzi C, et al.
552 Dramatic resurgence of malaria after 7 years of intensive vector control
553 interventions in Eastern Uganda. *PLOS Glob Public Health.* 2024 Aug
554 29;4(8):e0003254.
- 555 43. Namuganga JF, Epstein A, Nankabirwa JI, Mpimbaza A, Kiggundu M, Sserwanga
556 A, et al. The impact of stopping and starting indoor residual spraying on malaria
557 burden in Uganda. *Nat Commun.* 2021 May 11;12(1):2635.
- 558 44. Takyi A, Carrara VI, Dahal P, Przybylska M, Harriss E, Insaïdoo G, et al.
559 Characterisation of populations at risk of sub-optimal dosing of artemisinin-based
560 combination therapy in Africa. *PLOS Glob Public Health.* 2023 Dec
561 1;3(12):e0002059.
- 562 45. White NJ. Antimalarial drug resistance. *J Clin Invest.* 2004 Apr 15;113(8):1084–92.
- 563 46. Ndong Ngomo JM, Mawili-Mboumba DP, M'Bondoukwé NP, Ditombi BM, Koumba
564 Lengongo JV, Batchy Ognagosso FB, et al. Drug resistance molecular markers of
565 *Plasmodium falciparum* and severity of malaria in febrile children in the sentinel site
566 for malaria surveillance of Melen in Gabon: Additional data from the *Plasmodium*
567 diversity network African network. *Trop Med Infect Dis.* 2023 Mar 23;8(4):184.
- 568 47. Sisowath C, Petersen I, Veiga MI, Mårtensson A, Premji Z, Björkman A, et al. In
569 vivo selection of *Plasmodium falciparum* parasites carrying the
570 chloroquine-susceptible pfcrt K76 allele after treatment with
571 artemether-lumefantrine in Africa. *J Infect Dis.* 2009 Mar 1;199(5):750–7.
- 572 48. Venkatesan M, Gadalla NB, Stepniewska K, Dahal P, Nsanzabana C, Moriera C, et
573 al. Polymorphisms in *Plasmodium falciparum* chloroquine resistance transporter
574 and multidrug resistance 1 genes: parasite risk factors that affect treatment
575 outcomes for *P. falciparum* malaria after artemether-lumefantrine and
576 artesunate-amodiaquine. *Am J Trop Med Hyg.* 2014 Oct;91(4):833–43.
- 577 49. Okell LC, Reiter LM, Ebbe LS, Baraka V, Bisanzio D, Watson OJ, et al. Emerging
578 implications of policies on malaria treatment: genetic changes in the *Pfmdr-1* gene
579 affecting susceptibility to artemether-lumefantrine and artesunate-amodiaquine in
580 Africa. *BMJ Glob Health.* 2018 Oct 19;3(5):e000999.
- 581 50. Wang P, Read M, Sims PF, Hyde JE. Sulfadoxine resistance in the human malaria

- 582 parasite *Plasmodium falciparum* is determined by mutations in dihydropteroate
583 synthetase and an additional factor associated with folate utilization. *Mol Microbiol.*
584 1997 Mar;23(5):979–86.
- 585 51. Omar SA, Adagu IS, Warhurst DC. Can pretreatment screening for dhps and dhfr
586 point mutations in *Plasmodium falciparum* infections be used to predict
587 sulfadoxine-pyrimethamine treatment failure? *Trans R Soc Trop Med Hyg.* 2001
588 May;95(3):315–9.
- 589 52. Desai M, Gutman J, Taylor SM, Wiegand RE, Khairallah C, Kayentao K, et al.
590 Impact of sulfadoxine-pyrimethamine resistance on effectiveness of intermittent
591 preventive therapy for malaria in pregnancy at clearing infections and preventing
592 low birth weight. *Clin Infect Dis.* 2016 Feb 1;62(3):323–33.
- 593 53. Esu E, Tacoli C, Gai P, Berens-Riha N, Pritsch M, Loescher T, et al. Prevalence of
594 the Pfdhfr and Pfdhps mutations among asymptomatic pregnant women in
595 Southeast Nigeria. *Parasitol Res.* 2018 Mar;117(3):801–7.
- 596 54. Tchuente PVK, Ngum LN, Ali IM, Chedjou JPK, Nji AM, Netongo PM, et al.
597 *Plasmodium falciparum* dhps and dhfr markers of resistance to
598 sulfadoxine–pyrimethamine five years (2016–2020) after the implementation of
599 seasonal malaria chemoprevention in Cameroon. *Wellcome Open Res.* 2024 Jun
600 20;9:323.
- 601 55. Djontu JC, Baina MT, Mbama Ntahi JD, Lissom A, Umuhoza DM, Assiomo Doulamu
602 NV, et al. Profile of molecular markers of Sulfadoxine-Pyrimethamine-resistant
603 *Plasmodium falciparum* in individuals living in southern area of Brazzaville,
604 Republic of Congo. *Int J Parasitol Drugs Drug Resist.* 2024
605 Dec;26(100569):100569.
- 606 56. Nana RRD, Bayengue SSB, Mogtomo MLK, Ngane ARN, Singh V. Anti-folate
607 quintuple mutations in *Plasmodium falciparum* asymptomatic infections in Yaoundé,
608 Cameroon. *Parasitol Int.* 2023 Feb;92(102657):102657.
- 609 57. Nkemngbo FN, Raissa LW, Nguete DN, Ndo C, Fru-Cho J, Njiokou F, et al.
610 Geographical emergence of sulfadoxine-pyrimethamine drug resistance-associated
611 *P. falciparum* and *P. malariae* alleles in co-existing *Anopheles* mosquito and
612 asymptomatic human populations across Cameroon. *Antimicrob Agents*
613 *Chemother.* 2023 Dec 14;67(12):e0058823.
- 614 58. L'Episcopia M, Doderer-Lang C, Perrotti E, Priuli GB, Cavallari S, Guidetti C, et al.
615 Polymorphism analysis of drug resistance markers in *Plasmodium falciparum*
616 isolates from Benin. *Acta Trop.* 2023 Sep;245(106975):106975.
- 617 59. Apinjoh TO, Mugri RN, Miotto O, Chi HF, Tata RB, Anchang-Kimbi JK, et al.
618 Molecular markers for artemisinin and partner drug resistance in natural
619 *Plasmodium falciparum* populations following increased insecticide treated net
620 coverage along the slope of mount Cameroon: cross-sectional study. *Infect Dis*

- Poverty. 2017 Nov 6;6(1):136.
60. Mbacham WF, Evehe MSB, Netongo PM, Ateh IA, Mimche PN, Ajua A, et al. Efficacy of amodiaquine, sulphadoxine-pyrimethamine and their combination for the treatment of uncomplicated *Plasmodium falciparum* malaria in children in Cameroon at the time of policy change to artemisinin-based combination therapy. *Malar J*. 2010 Jan 27;9(1):34.
61. Guémas E, Coppée R, Ménard S, du Manoir M, Nsango S, Makaba Mvumbi D, et al. Evolution and spread of *Plasmodium falciparum* mutations associated with resistance to sulfadoxine-pyrimethamine in central Africa: a cross-sectional study. *Lancet Microbe*. 2023 Dec;4(12):e983–93.
62. Popkin-Hall ZR, Niaré K, Crudale R, Simkin A, Fola AA, Sanchez JF, et al. High-Throughput Genotyping of *Plasmodium vivax* in the Peruvian Amazon via Molecular Inversion Probes. *medRxiv* [Internet]. 2024 Jun 28; Available from: <http://dx.doi.org/10.1101/2024.06.27.24309599>
63. Chang HH, Worby CJ, Yeka A, Nankabirwa J, Kamya MR, Staedke SG, et al. THE REAL McCOIL: A method for the concurrent estimation of the complexity of infection and SNP allele frequency for malaria parasites. *PLoS Comput Biol*. 2017 Jan;13(1):e1005348.
64. Paschalidis A, Watson OJ, Aydemir O, Verity R, Bailey JA. coiaf: directly estimating complexity of infection with allele frequencies [Internet]. *bioRxiv*. 2022 [cited 2022 Sep 7]. p. 2022.05.26.493561. Available from: <https://www.biorxiv.org/content/10.1101/2022.05.26.493561v1>