

Circulation of an Artemisinin-Resistant Malaria Lineage in a Traveler Returning from East Africa to France

Romain Coppée,^{1,a,b} Justine Bailly,^{1,2,b} Véronique Sarrasin,^{1,2} Bertin Vianou,³ Boris-Enock Zinsou,³ Edith Mazars,⁴ Hugues Georges,⁵ Samia Hamane,⁶ Rose-Anne Lavergne,⁷ Eric Dannaoui,⁸ Betty Balikagala,⁹ Naoyuki Fukuda,⁹ Emmanuel I. Odongo-Aginya,¹⁰ Toshihiro Mita,⁹ Sandrine Houzé,^{1,2} and Jérôme Clain^{1,2}, on behalf of the French National Malaria Reference Center Study Group

¹Université de Paris, IRD, MERIT, Paris, France; ²Centre National de Référence du Paludisme, AP-HP, Hôpital Bichat-Claude Bernard, Paris, France; ³Institut de Recherche Clinique du Bénin, Cotonou, Bénin; ⁴Laboratoire de microbiologie, Pôle de Biologie-hygiène, Centre Hospitalier de Valenciennes, Valenciennes, France; ⁵Service de Réanimation Médicale et Maladies Infectieuses, Hôpital Chatiliez, Tourcoing, France; ⁶Laboratoire de Parasitologie-Mycologie, AP-HP, Groupe Hospitalier Saint-Louis Lariboisière Fernand-Widal, Paris, France; ⁷Laboratoire de Parasitologie-Mycologie, Institut de Biologie, Centre Hospitalier Universitaire de Nantes, Nantes, France; ⁸Laboratoire de Parasitologie-Mycologie, AP-HP, Département de Microbiologie, Hôpital Européen Georges Pompidou, Paris, France; ⁹Department of Tropical Medicine and Parasitology, School of Medicine, Juntendo University, Hongo, Bunkyo-ku, Tokyo, Japan; and ¹⁰Faculty of Medicine, Gulu University, Gulu, Uganda

A returned traveler to Uganda presented with a *Plasmodium falciparum* *kelch13* A675V mutant infection that exhibited delayed clearance under artesunate therapy. Parasites were genetically related to recently reported Ugandan artemisinin-resistant A675V parasites. Adequate malaria prevention measures and clinical and genotypic surveillance are important tools to avoid and track artemisinin resistance.

Keywords. *Plasmodium falciparum*; *kelch13*; artemisinin resistance; East Africa; whole-genome sequencing

Artemisinin (ART)-based combination therapies (ACTs) are the current first-line treatments for uncomplicated *Plasmodium falciparum* malaria [1]. However, ART-resistant *P. falciparum* parasites that display delayed clearance have emerged and largely spread in the Greater Mekong Subregion of Southeast Asia during the past 15 years [2, 3]. ART resistance is mainly conferred by nonsynonymous mutations in the *kelch13* gene of the parasite (PlasmoDB accession: PF3D7_1343700) [4, 5], and about 10 of those mutations are validated markers of clinical ART resistance in Southeast Asia [6, 7].

ART-resistant parasites harboring the *kelch13* mutations R561H, A675V, and C469Y are now independently emerging in Rwanda and Northern Uganda, East Africa [8, 9]. Because no alternative to ART derivatives is currently available, the spread of ART-resistant parasites in African settings could threaten the long-term use of ACTs to treat malaria. A continual screening of *kelch13* gene diversity in Africa is mandatory to track the presence of ART-resistant parasites lineages.

Here we surveyed *kelch13* mutations in *P. falciparum* isolates from 125 malaria cases imported in France after a travel to Eastern Africa (Burundi, Djibouti, Eritrea, Ethiopia, Kenya, Rwanda, Sudan, Tanzania, and Uganda) between January 2006 and December 2020 (Supplementary Figure 1). Malaria was diagnosed in French metropolitan hospitals participating to the national malaria survey network (French National Malaria Reference Center, France) using thin or thick blood smear or both. Thick blood smears were considered positive if at least 1 trophozoite was visualized after examination of 1 000 leukocytes. The *Plasmodium* species identification was confirmed microscopically or with polymerase chain reaction (PCR) in case of low parasitemia. Patients were treated according to French recommendations [9].

Using this biobank, we identified by Sanger sequencing 1 adult patient with parasites carrying the *kelch13* A675V mutation (Supplementary Figure 1). The patient first traveled to Northern Thailand (Southeast Asia) and, then, about 1 month later, visited several regions of Uganda (East Africa) in December 2019/January 2020. The patient reported to sleep under insecticide-treated bed net but not to take antimalarial medicines for prophylaxis. At hospital admission in France, 2 weeks after the return from Uganda, *P. falciparum* malaria was diagnosed with hyperparasitemia (765 000 parasites/μL) and other biological criteria of severity. Four intravenous artesunate doses (2.4 mg per kilogram of body weight) were administered at hours 0, 12, 24, and 48, followed by a full course of dihydroartemisinin-piperaquine on day 3 [10]. Although a day-3 follow-up parasitemia is recommended, a day-2 plus 8 hours (56 hours) was done instead for logistical reasons within the intensive care unit. At 56 hours post-treatment, the parasitemia was still positive (20 250 parasites/μL; 37.8-fold reduction compared to pretreatment). No other follow-up parasitemia was available between day 0 and day 7. On days 7 and 36 follow-ups, blood smears were negative, indicating that the overall parasitological response was adequate. As the positive parasitemia at 56 hours was suggestive of a delayed parasite clearance in the traveler, we looked for follow-up parasitemia at a similar timepoint in a reference set of Ugandan infections (n = 229) similarly treated and in which the *kelch13* A675V mutation was frequent [9]. In

Received 11 January 2022; editorial decision 18 February 2022; published online 25 February 2022.

^aCurrent affiliation: Université de Paris, Infection Modélisation Antimicrobienne Evolution (IAMÉ), Inserm UMR1137, F-75018 Paris, France.

^bR. C. and J. B. contributed equally to this work.

Correspondence: J. Clain, Université de Paris, IRD, MERIT, Centre National de Référence du Paludisme, AP-HP, Hôpital Bichat-Claude Bernard, F-75018 Paris, France (jerome.clain@u-paris.fr).

Clinical Infectious Diseases® 2022;75(7):1242–4

© The Author(s) 2022. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. <https://doi.org/10.1093/cid/ciac162>

this trial in Uganda, artesunate was administered intravenously at a dose of 3 mg or 2.4 mg per kilogram of body weight (in patients weighing <20 kg or ≥20 kg, respectively) at 0, 12, and 24 hours, followed by a standard 3-day course of artemether-lumefantrine. If parasites were still observed after the initial 3 doses, additional doses of intravenous artesunate were administered every 24 hours until parasite clearance was observed. Follow-up parasitemia were collected at multiple timepoints: 0, 4, and 6 hours and then at 6-hour intervals until 2 consecutive negative blood smears had been obtained. Therefore, we extracted the parasitemia collected at 54 hours in the trial in Uganda as the closest timepoint to the one collected at 56 hours in the traveler. Of the 28 Ugandan patients carrying parasites with the *kelch13* A675V mutation, 10 (35.7%) showed parasite positive at 54 hours, whereas of the 201 patients with the *kelch13* wild-type allele, one only (0.5%) showed parasite positive at 54 hours ($P < .001$ by the Fisher exact test). In this *kelch13* A675V population ($n = 28$), the largest ratio of parasitemia at 54 hours to pretreatment parasitemia was 0.0098, compared to a ratio of 0.0260 at 56 hours for the *kelch13* A675V from the traveler (Figure 1A). Altogether, despite different epidemiological and demographic contexts, the data are suggesting a delayed parasite clearance in the traveler under artesunate therapy.

We determined the whole genome of the *kelch13* A675V mutant parasites from the traveler by Illumina sequencing using the

selective whole-genome amplification strategy (Supplementary Method 1 and Supplementary Tables 1 and 2; next-generation sequence files are accessible on the European Nucleotide Archive under the accession number ERS7266441). This new *kelch13* A675V genome was compared with 131 published African and Asian parasite genomes, including 14 ART-resistant A675V mutants from Uganda recently reported (Supplementary Table 3) [9, 11–13]. Using 20 266 genome-wide, high-quality single-nucleotide polymorphisms (SNPs) (Supplementary Method 2), the phylogenetic tree showed that the parasite isolate from the traveler was genetically highly related to parasites from Uganda but not from Southeast Asia (Figure 1B). This indicated that the traveler got infected in Uganda, rather than during the recent previous trip in Thailand. Also, the A675V mutant from the traveler formed a monophyletic group with previously reported Ugandan A675V mutants, within the East African clade (Figure 1B). The high genetic relatedness of all Ugandan A675V mutants was also evidenced at the *kelch13* locus using a subset of 238 SNPs flanking the *kelch13* gene within ±100 kb (Supplementary Figure 2). We conclude that the traveler got infected by ART-resistant *kelch13* A675V parasites related to those already circulating in Uganda [9].

In conclusion, this study shows that people traveling to East Africa are at risk of being infected with ART-resistant malaria parasites associated with delayed parasite clearance under




Figure 1. In vivo clearance phenotype and geographic origin of the *kelch13* A675V mutant. A, Ratio of parasitemia at 54 hours post-treatment to pretreatment parasitemia by *kelch13* genotype (blue: wild-type [WT], red: A675V mutation). Data are from Ugandan patients reported in [9]. For the A675V mutant isolate from this study, we estimated the ratio of parasitemia at 56 hours post-treatment to pretreatment parasitemia. Statistical comparison was done using Mann-Whitney U test and did not include the data from the traveler. B, Genome-wide phylogenetic tree of the 15 *kelch13* A675V parasite isolates (one from this study, and 14 from [9] and [11]) and 117 parasite isolates collected in Sub-Saharan Africa and Southeast Asia [12, 13]. Countries of collection of the samples are Benin, Burkina-Faso, Nigeria for West Africa (blue); Congo and Gabon for Central Africa (yellow); Kenya, Tanzania, and Uganda for East Africa (red); and Cambodia, Myanmar, Thailand, and Vietnam for Southeast Asia (green). Tree was inferred by neighbor-joining using 20 266 SNPs. Isolates marked with a green or red star are A675V mutants from Southeast Asia or Uganda, respectively. Abbreviation: SNP, single-nucleotide polymorphism.

treatment with artemisinin derivatives. As highly mobile individuals, travelers may also participate in spreading drug resistance [14]. Here, the traveler visited several regions of Uganda, although we do not know the precise itinerary, and exhibited malaria symptoms (which appear when parasites are circulating in blood) only after returning to France. The traveler therefore likely did not spread this A675V resistant strain in some other parts of Uganda. However, vigilance is mandatory to avoid such a scenario in the future, especially with mid- or long-term travelers. Adequate malaria prevention measures are warranted in travelers to protect from both the disease and drug resistance spread. Also, monitoring of ACT therapeutic efficacy and genotypic surveillance of drug resistance should be carefully conducted for imported malaria.

Supplementary Data

Supplementary materials are available at *Clinical Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. The authors thank the patients and the staff of the hospitals of the network of the French National Malaria Reference Center. The members of the French National Malaria Reference Center Study Group are as follows: A. Delaval (Centre Hospitalier Intercommunal Robert Ballanger, Aulnay-sous-bois), H. Broutier (Centre Hospitalier Intercommunal Robert Ballanger, Aulnay-sous-bois), C. Leprince (Centre Hospitalier Intercommunal Robert Ballanger, Aulnay-sous-bois), P. Agnamey (Centre Hospitalier Universitaire Amiens, Amiens), C. Damiani (Centre Hospitalier Universitaire Amiens, Amiens), L. De Gentile (Centre hospitalier Universitaire Angers, Angers), S. Houze (Hôpital Bichat-Claude Bernard, Paris), N. Argy (Hôpital Bichat-Claude Bernard, Paris), A. Enache Angoulvant (Centre Hospitalier Universitaire Kremlin-Bicêtre, Kremlin-Bicêtre), T. Pistone (Groupe Hospitalier Pellegrin, Bordeaux), C. Rapp (Hôpital d'Instruction des Armées Bégin, Saint Mandé), J.-C. Pilo (Hôpital d'Instruction des Armées Bégin, Saint Mandé), G. Carroger (Centre Hospitalier Jacques Cœur, Bourges), L. Million (Centre Hospitalier Régional Universitaire Besançon, Besançon), D. Quinio (Centre Hospitalier Universitaire de Brest, Brest), D. Raffenot (Centre Hospitalier de Chambéry, Chambéry), N. Dahane (Hôpital Cochin, Paris), T. Mergey (Centre Hospitalier de Cholet, Cholet), D. Pons (Centre Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand), B. Cuisenier (Centre Hospitalier Universitaire de Dijon, Dijon), C. Garnaud (Centre Hospitalier Universitaire de Grenoble, Grenoble), E. Dannaoui (Hôpital Européen Georges Pompidou, Paris), F. Botterel (Hôpital Henri Mondor, Créteil), A. Durand (Institut Pasteur, Paris), D. Bemba (Hôpital Jean Verdier, Bondy), A.-S. Deleplancque (Centre Hospitalier Universitaire de Lille), P. Penn (Centre Hospitalier Le Mans, Le Mans), G. Pean-de-Ponfilly (Hôpital Lariboisière, Paris), A. Mendes-Moreira (Centre Hospitalier de La Rochelle, La Rochelle), M.-E. Bougnoux (Hôpital Necker, Paris), C. Pomares-Estran (Centre Hospitalier Universitaire de l'Archet, Nice), M. Sasso (Hôpital Universitaire Caremeau, Nîmes), B. Buret (Centre Hospitalier de Niort, Niort), R.-A. Lavergne (Centre Hospitalier Universitaire de Nantes, Nantes), L. Bret (Centre Hospitalier Universitaire d'Orléans, Orléans), M. Thellier (Hôpital de la Pitié-Salpêtrière, Paris), A. Taieb (Hôpital de la Pitié-Salpêtrière, Paris), J.-Y. Peltier (Centre Hospitalier Intercommunal Poissy/Saint-Germain-en-Laye, Poissy), E. Perraud-Cateau (Centre Hospitalier Universitaire de Poitiers, Poitiers), D. Toubas (Hôpital Maison Blanche, Reims), O. Albert

(Hôpital Maison Blanche, Reims), S. Belaz (Hôpital Pontchaillou, Rennes), G. Gargala (Centre Hospitalier Universitaire de Rouen, Rouen), L. Favennec (Centre Hospitalier Universitaire de Rouen, Rouen), G. Belkadi (Hôpital Saint-Antoine, Paris), A. Abou-Bacar (Centre Hospitalier Universitaire de Strasbourg, Strasbourg), N. Godineau (Centre Hospitalier Delafontaine, Saint Denis), S. Hamane (Hôpital Saint-Louis, Paris), G. Ménard (Hôpital d'Instruction des Armées Saint-Anne, Toulon), M. Develoux (Hôpital Tenon, Paris), E. Mazars (Centre Hospitalier de Valenciennes, Valenciennes), O. Eloy (Centre Hospitalier de Versailles, Versailles).

Ethics statement. Samples received at the French National Malaria Reference Center were registered and declared for research purposes as a biobank for both the Assistance Publique des Hôpitaux de Paris and the French National Institute of Health Survey. No institutional review board approval was required according to French legislation (article L. 1111-7 du Code de la Santé Publique, article L. 1211-2 du Code de Santé Publique, articles 39 et suivants de la loi 78-17 du 6 janvier 1978 modifiée en 2004 relative à l'informatique, aux fichiers, et aux libertés). The online reporting system for malaria cases of the French National Malaria Reference Center has been declared to the "Commission Nationale de l'Informatique et des Libertés" (CNIL) under number 1223103.

Financial support. This work was partly supported by Santé Publique France (to S. H.), the Agence Nationale de la Recherche (grant number ANR-17-CE15-0013-03 to J. C.). T. M. reports Grants-in-aid for scientific research (grant numbers 26460515, 26305015, and 17H04074) from Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan.

Potential conflicts of interest. All authors: No reported conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

References

- World Malaria Report 2019. Available at: <https://www.who.int/publications-detail-redirect/9789241565721>. Accessed 8 March 2022.
- Ashley EA, Dhorda M, Fairhurst RM, et al. Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 2014; 371:411–23.
- Dondorp AM, Nosten F, Yi P, et al. Artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 2009; 361:455–67.
- Ariey F, Witkowski B, Amaratunga C, et al. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature* 2014; 505:50–5.
- Straimer J, Gnädig NF, Witkowski B, et al. Drug resistance. K13-propeller mutations confer artemisinin resistance in *Plasmodium falciparum* clinical isolates. *Science* 2015; 347:428–31.
- Report on antimalarial drug efficacy, resistance and response: 10 years of surveillance (2010–2019). Available at: <https://www.who.int/publications-detail-redirect/9789240012813>. Accessed 8 March 2022.
- WWARN K13 Genotype-Phenotype Study Group. Association of mutations in the *Plasmodium falciparum* Kelch13 gene (Pf3D7_1343700) with parasite clearance rates after artemisinin-based treatments—a WWARN individual patient data meta-analysis. *BMC Med* 2019; 17:1.
- Uwimana A, Legrand E, Stokes BH, et al. Emergence and clonal expansion of in vitro artemisinin-resistant *Plasmodium falciparum* kelch13 R561H mutant parasites in Rwanda. *Nat Med* 2020; 26:1602–8.
- Balikagala B, Fukuda N, Ikeda M, et al. Evidence of artemisinin-resistant malaria in Africa. *N Engl J Med* 2021; 385:1163–71.
- Bouchaud O, Bruneel F, Caumes E, et al. Management and prevention of imported malaria: 2018 update of the 2007 French clinical guidelines. *Med Mal Infect* 2020; 50:161–93.
- Ikeda M, Kaneko M, Tachibana S-I, et al. Artemisinin-resistant *Plasmodium falciparum* with high survival rates, Uganda, 2014–2016. *Emerg Infect Dis* 2018; 24:718–26.
- Malaria GEN, Ahouidi A, Ali M, et al. An open dataset of *Plasmodium falciparum* genome variation in 7000 worldwide samples. *Wellcome Open Res* 2021; 6:42.
- Amambua-Ngwa A, Amenga-Etego L, Kamau E, et al. Major subpopulations of *Plasmodium falciparum* in sub-Saharan Africa. *Science* 2019; 365:813–6.
- Tatem AJ, Jia P, Ordanovich D, et al. The geography of imported malaria to non-endemic countries: a meta-analysis of nationally reported statistics. *Lancet Infect Dis* 2017; 17:98–107.