

ColumbiaX: Machine Learning

Prof. John Paisley

Department of Electrical Engineering
& Data Science Institute

Columbia University

ColumbiaX: Machine Learning

Lecture 1

Prof. John Paisley

Department of Electrical Engineering
& Data Science Institute

Columbia University

OVERVIEW

This class will cover model-based techniques for extracting information from data with an end-task in mind. Such tasks include:


- ▶ predicting an unknown “output” given its corresponding “input”
- ▶ uncovering information within the data to better understand it
- ▶ data-driven recommendation, grouping, classification, ranking, etc.

There are a few ways we can divide up the material as we go along, e.g.,


supervised learning		unsupervised learning
probabilistic models		non-probabilistic models
modeling approach		optimization techniques

We'll adopt the first method and work in the second two along the way.

OVERVIEW: SUPERVISED LEARNING



(a) Regression



(b) Classification

Regression: Using set of inputs, predict real-valued output.

Classification: Using set of inputs, predict a discrete label (aka class).

EXAMPLE CLASSIFICATION PROBLEM

Given a set of inputs characterizing an item, assign it a label.

Is this spam?


hi everyone,

i saw that close to my hotel there is a pub with bowling
(it's on market between 9th and 10th avenue). meet
there at 8:30?


What about this?

Enter for a chance to win a trip to Universal Orlando to
celebrate the arrival of Dr. Seuss's The Lorax on Movies
On Demand on August 21st! [Click here now!](#)

OVERVIEW: UNSUPERVISED LEARNING



(c) topic modeling



(d) recommendations¹

With unsupervised learning our goal is often to uncover structure in the data. This helps with predictions, recommendations, efficient data exploration.

¹ Figure from Koren, Y., Robert B., and Volinsky, C.. "Matrix factorization techniques for recommender systems." Computer 42.8 (2009): 30-37.


EXAMPLE UNSUPERVISED PROBLEM

Goal: Learn the dominant topics from a set of news articles.

The New York Times

music band songs rock album jazz pop song singer night	book life novel story books man stories love children family	art museum show exhibition artist artists paintings painting century works	game knicks nets points team season play games night coach	show film television movie series says life man character know
theater play production show stage street broadway director musical directed	clinton bush campaign gore political republican dole presidential senator house	stock market percent fund investors funds companies stocks investment trading	restaurant sauce menu food dishes street dining dinner chicken served	budget tax governor county mayor billion taxes plan legislature fiscal

DATA MODELING




- ▶ Supervised vs. unsupervised: Blocks #1 and #4
- ▶ Probabilistic vs. non-probabilistic: Primarily Block #2 (Some Block #3)
- ▶ Model development (Block #2) vs. Optimization techniques (Block #3)

GAUSSIAN DISTRIBUTION (MULTIVARIATE)

Gaussian density in d dimensions

- ▶ Block #1: Data x_1, \dots, x_n . Each $x_i \in \mathbb{R}^d$
- ▶ Block #2: An i.i.d. Gaussian model
- ▶ Block #3: Maximum likelihood
- ▶ Block #4: Leave undefined



The density function is

$$p(x|\mu, \Sigma) := \frac{1}{(2\pi)^{\frac{d}{2}} \sqrt{\det(\Sigma)}} \exp\left(-\frac{1}{2}(x-\mu)^T \Sigma^{-1}(x-\mu)\right)$$

The central moments are:

$$\mathbb{E}[x] = \int_{\mathbb{R}^d} x p(x|\mu, \Sigma) dx = \mu,$$

$$\text{Cov}(x) = \mathbb{E}[(x - \mathbb{E}[x])(x - \mathbb{E}[x])^T] = \mathbb{E}[xx^T] - \mathbb{E}[x]\mathbb{E}[x]^T = \Sigma.$$

BLOCK #2: A PROBABILISTIC MODEL

Probabilistic Models

- ▶ A *probabilistic model* is a set of probability distributions, $p(x|\theta)$.
- ▶ We pick the *distribution family* $p(\cdot)$, but don't know the parameter θ .

Example: Model data with a Gaussian distribution $p(x|\theta)$, $\theta = \{\mu, \Sigma\}$.

The i.i.d. assumption

Assume data is *independent and identically distributed (iid)*. This is written

$$x_i \stackrel{iid}{\sim} p(x|\theta), \quad i = 1, \dots, n.$$

Writing the density as $p(x|\theta)$, then the *joint* density decomposes as

$$p(x_1, \dots, x_n|\theta) = \prod_{i=1}^n p(x_i|\theta).$$

BLOCK #3: MAXIMUM LIKELIHOOD ESTIMATION

Maximum Likelihood approach

We now need to find θ . *Maximum likelihood* seeks the value of θ that maximizes the likelihood function:

$$\hat{\theta}_{\text{ML}} := \arg \max_{\theta} p(x_1, \dots, x_n | \theta),$$

This value best explains the data according to the chosen distribution family.

Maximum Likelihood equation

The analytic criterion for this maximum likelihood estimator is:

$$\nabla_{\theta} \prod_{i=1}^n p(x_i | \theta) = 0.$$

Simply put, the maximum is at a peak. There is no “upward” direction.

BLOCK #3: LOGARITHM TRICK

Logarithm trick

Calculating $\nabla_{\theta} \prod_{i=1}^n p(x_i|\theta)$ can be complicated. We use the fact that the logarithm is monotonically increasing on \mathbb{R}_+ , and the equality

$$\ln\left(\prod_i f_i\right) = \sum_i \ln(f_i).$$

Consequence: Taking the logarithm does not change the *location* of a maximum or minimum:

$$\max_y \ln g(y) \neq \max_y g(y)$$

The *value* changes.

$$\arg \max_y \ln g(y) = \arg \max_y g(y)$$

The *location* does not change.

BLOCK #3: ANALYTIC MAXIMUM LIKELIHOOD

Maximum likelihood and the logarithm trick

$$\hat{\theta}_{\text{ML}} = \arg \max_{\theta} \prod_{i=1}^n p(x_i|\theta) = \arg \max_{\theta} \ln \left(\prod_{i=1}^n p(x_i|\theta) \right) = \arg \max_{\theta} \sum_{i=1}^n \ln p(x_i|\theta)$$

To then solve for $\hat{\theta}_{\text{ML}}$, find

$$\nabla_{\theta} \sum_{i=1}^n \ln p(x_i|\theta) = \sum_{i=1}^n \nabla_{\theta} \ln p(x_i|\theta) = 0.$$

Depending on the choice of the model, we will be able to solve this

1. analytically (via a simple set of equations)
2. numerically (via an iterative algorithm using different equations)
3. approximately (typically when #2 converges to a local optimal solution)

EXAMPLE: MULTIVARIATE GAUSSIAN MLE

Block #2: Multivariate Gaussian data model

Model: Set of all Gaussians on \mathbb{R}^d with unknown mean $\mu \in \mathbb{R}^d$ and covariance $\Sigma \in \mathbb{S}_{++}^d$ (positive definite $d \times d$ matrix).

We assume that x_1, \dots, x_n are i.i.d. $p(x|\mu, \Sigma)$, written $x_i \stackrel{iid}{\sim} p(x|\mu, \Sigma)$.

Block #3: Maximum likelihood solution

We have to solve the equation

$$\sum_{i=1}^n \nabla_{(\mu, \Sigma)} \ln p(x_i|\mu, \Sigma) = 0$$

for μ and Σ . (Try doing this without the log to appreciate it's usefulness.)

EXAMPLE: GAUSSIAN MEAN MLE

First take the gradient with respect to μ .

$$\begin{aligned} 0 &= \nabla_{\mu} \sum_{i=1}^n \ln \frac{1}{\sqrt{(2\pi)^d |\Sigma|}} \exp\left(-\frac{1}{2}(x_i - \mu)^T \Sigma^{-1}(x_i - \mu)\right) \\ &= \nabla_{\mu} \sum_{i=1}^n -\frac{1}{2} \ln(2\pi)^d |\Sigma| - \frac{1}{2}(x_i - \mu)^T \Sigma^{-1}(x_i - \mu) \\ &= -\frac{1}{2} \sum_{i=1}^n \nabla_{\mu} \left(x_i^T \Sigma^{-1} x_i - 2\mu^T \Sigma^{-1} x_i + \mu^T \Sigma^{-1} \mu \right) = -\Sigma^{-1} \sum_{i=1}^n (x_i - \mu) \end{aligned}$$

Since Σ is positive definite, the only solution is

$$\sum_{i=1}^n (x_i - \mu) = 0 \quad \Rightarrow \quad \hat{\mu}_{\text{ML}} = \frac{1}{n} \sum_{i=1}^n x_i$$

Since this solution is independent of Σ , it doesn't depend on $\hat{\Sigma}_{\text{ML}}$.

EXAMPLE: GAUSSIAN COVARIANCE MLE

Now take the gradient with respect to Σ .

$$\begin{aligned} 0 &= \nabla_{\Sigma} \sum_{i=1}^n -\frac{1}{2} \ln(2\pi)^d |\Sigma| - \frac{1}{2} (x_i - \mu)^T \Sigma^{-1} (x_i - \mu) \\ &= -\frac{n}{2} \nabla_{\Sigma} \ln |\Sigma| - \frac{1}{2} \nabla_{\Sigma} \text{trace} \left(\Sigma^{-1} \sum_{i=1}^n (x_i - \mu)(x_i - \mu)^T \right) \\ &= -\frac{n}{2} \Sigma^{-1} + \frac{1}{2} \Sigma^{-2} \sum_{i=1}^n (x_i - \mu)(x_i - \mu)^T \end{aligned}$$

Solving for Σ and plugging in $\mu = \hat{\mu}_{\text{ML}}$,

$$\hat{\Sigma}_{\text{ML}} = \frac{1}{n} \sum_{i=1}^n (x_i - \hat{\mu}_{\text{ML}})(x_i - \hat{\mu}_{\text{ML}})^T.$$

EXAMPLE: GAUSSIAN MLE (SUMMARY)

So if we have data x_1, \dots, x_n in \mathbb{R}^d that we hypothesize is i.i.d. Gaussian, the maximum likelihood values of the mean and covariance matrix are

$$\hat{\mu}_{\text{ML}} = \frac{1}{n} \sum_{i=1}^n x_i, \quad \hat{\Sigma}_{\text{ML}} = \frac{1}{n} \sum_{i=1}^n (x_i - \hat{\mu}_{\text{ML}})(x_i - \hat{\mu}_{\text{ML}})^T.$$

Are we done? There are many assumptions/issues with this approach that makes finding the “best” parameter values not a complete victory.

- ▶ We made a model assumption (multivariate Gaussian).
- ▶ We made an i.i.d. assumption.
- ▶ We assumed that maximizing the likelihood is the best thing to do.

Comment: We often use θ_{ML} to make predictions about x_{new} (Block #4).

How does θ_{ML} generalize to x_{new} ?

If $x_{1:n}$ don't “capture the space” well, θ_{ML} can *overfit* the data.

ColumbiaX: Machine Learning

Lecture 4

Prof. John Paisley

Department of Electrical Engineering
& Data Science Institute


Columbia University

LINEAR REGRESSION

EXAMPLE: OLD FAITHFUL




EXAMPLE: OLD FAITHFUL



Can we meaningfully predict the time between eruptions only using the duration of the last eruption?

EXAMPLE: OLD FAITHFUL



Can we meaningfully predict the time between eruptions only using the duration of the last eruption?

EXAMPLE: OLD FAITHFUL


One model for this

$$(\text{wait time}) \approx w_0 + (\text{last duration}) \times w_1$$

- ▶ w_0 and w_1 are to be learned.
- ▶ This is an example of linear regression.

Refresher

w_1 is the slope, w_0 is called the intercept, bias, shift, offset.




HIGHER DIMENSIONS

Two inputs

$$(\text{output}) \approx w_0 + (\text{input 1}) \times w_1 + (\text{input 2}) \times w_2$$

With two inputs the intuition
is the same \longrightarrow



REGRESSION: PROBLEM DEFINITION

Data

Input: $x \in \mathbb{R}^d$ (i.e., measurements, covariates, features, indepen. variables)

Output: $y \in \mathbb{R}$ (i.e., response, dependent variable)

Goal

Find a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $y \approx f(x; w)$ for the data pair (x, y) .
 $f(x; w)$ is called a *regression function*. Its free parameters are w .

Definition of linear regression

A regression method is called *linear* if the prediction f is a linear function of the unknown parameters w .

LEAST SQUARES LINEAR REGRESSION MODEL

Model

The linear regression model we focus on now has the form

$$y_i \approx f(x_i; w) = w_0 + \sum_{j=1}^d x_{ij} w_j.$$

Model learning

We have the set of *training data* $(x_1, y_1) \dots (x_n, y_n)$. We want to use this data to learn a w such that $y_i \approx f(x_i; w)$. But we first need an *objective function* to tell us what a “good” value of w is.

Least squares

The *least squares* objective tells us to pick the w that minimizes the sum of squared errors

$$w_{\text{LS}} = \arg \min_w \sum_{i=1}^n (y_i - f(x_i; w))^2 \equiv \arg \min_w \mathcal{L}.$$

LEAST SQUARES IN PICTURES


Observations:

Vertical length is error.

The objective function \mathcal{L} is the sum of all the squared lengths.

Find weights (w_1, w_2) plus an offset w_0 to minimize \mathcal{L} .

(w_0, w_1, w_2) defines this plane.



EXAMPLE: EDUCATION, SENIORITY AND INCOME

2-dimensional problem

Input: (education, seniority) $\in \mathbb{R}^2$.


Output: (income) $\in \mathbb{R}$

Model: (income) $\approx w_0 + (\text{education})w_1 + (\text{seniority})w_2$

Question: Both $w_1, w_2 > 0$. What does this tell us?

Answer: As education and/or seniority goes up, income tends to go up.

(Caveat: This is a statement about correlation, not causation.)



LEAST SQUARES LINEAR REGRESSION MODEL

Thus far

We have data pairs (x_i, y_i) of measurements $x_i \in \mathbb{R}^d$ and a response $y_i \in \mathbb{R}$.
We believe there is a linear relationship between x_i and y_i ,

$$y_i = w_0 + \sum_{j=1}^d x_{ij} w_j + \epsilon_i$$

and we want to minimize the objective function

$$\mathcal{L} = \sum_{i=1}^n \epsilon_i^2 = \sum_{i=1}^n (y_i - w_0 - \sum_{j=1}^d x_{ij} w_j)^2$$

with respect to (w_0, w_1, \dots, w_d) .

Can math notation make this easier to look at/work with?

NOTATION: VECTORS AND MATRICES

We think of data with d dimensions as a *column* vector:

$$x_i = \begin{bmatrix} x_{i1} \\ x_{i2} \\ \vdots \\ x_{id} \end{bmatrix} \quad (\text{e.g.}) \Rightarrow \begin{bmatrix} \text{age} \\ \text{height} \\ \vdots \\ \text{income} \end{bmatrix}$$

A set of n vectors can be stacked into a matrix:

$$\mathbf{X} = \begin{bmatrix} x_{11} & \dots & x_{1d} \\ x_{21} & \dots & x_{2d} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nd} \end{bmatrix} = \begin{bmatrix} -x_1^T- \\ -x_2^T- \\ \vdots \\ -x_n^T- \end{bmatrix}$$

Assumptions for now:

- ▶ All features are treated as continuous-valued ($x \in \mathbb{R}^d$)
- ▶ We have more observations than dimensions ($d < n$)

NOTATION: REGRESSION (AND CLASSIFICATION)

Usually, for linear regression (and classification) we include an intercept term w_0 that doesn't interact with any element in the vector $x \in \mathbb{R}^d$.

It will be convenient to attach a 1 to the first dimension of each vector x_i (which we indicate by $x_i \in \mathbb{R}^{d+1}$) and in the first column of the matrix X :

$$x_i = \begin{bmatrix} 1 \\ x_{i1} \\ x_{i2} \\ \vdots \\ x_{id} \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & x_{11} & \dots & x_{1d} \\ 1 & x_{21} & \dots & x_{2d} \\ \vdots & & \vdots & \\ 1 & x_{n1} & \dots & x_{nd} \end{bmatrix} = \begin{bmatrix} 1 - x_1^T - \\ 1 - x_2^T - \\ \vdots \\ 1 - x_n^T - \end{bmatrix}.$$

We also now view $w = [w_0, w_1, \dots, w_d]^T$ as $w \in \mathbb{R}^{d+1}$.

LEAST SQUARES IN VECTOR FORM

Original least squares objective function: $\mathcal{L} = \sum_{i=1}^n (y_i - w_0 - \sum_{j=1}^d x_{ij}w_j)^2$

Using vectors, this can now be written: $\mathcal{L} = \sum_{i=1}^n (y_i - x_i^T w)^2$

Least squares solution (vector version)

We can find w by setting,

$$\nabla_w \mathcal{L} = 0 \quad \Rightarrow \quad \sum_{i=1}^n \nabla_w (y_i^2 - 2w^T x_i y_i + w^T x_i x_i^T w) = 0.$$

Solving gives,

$$-\sum_{i=1}^n 2y_i x_i + \left(\sum_{i=1}^n 2x_i x_i^T \right) w = 0 \quad \Rightarrow \quad w_{\text{LS}} = \left(\sum_{i=1}^n x_i x_i^T \right)^{-1} \left(\sum_{i=1}^n y_i x_i \right).$$

LEAST SQUARES IN MATRIX FORM

Least squares solution (matrix version)

Least squares in matrix form is even cleaner.

Start by organizing the y_i in a column vector, $y = [y_1, \dots, y_n]^T$. Then

$$\mathcal{L} = \sum_{i=1}^n (y_i - x_i^T w)^2 = \|y - Xw\|^2 = (y - Xw)^T (y - Xw).$$

If we take the gradient with respect to w , we find that

$$\nabla_w \mathcal{L} = 2X^T Xw - 2X^T y = 0 \quad \Rightarrow \quad w_{\text{LS}} = (X^T X)^{-1} X^T y.$$

RECALL FROM LINEAR ALGEBRA

Recall: Matrix \times vector ($X^T y = \sum_{i=1}^n y_i x_i$)

$$\begin{bmatrix} | & | & \dots & | \\ x_1 & x_2 & \dots & x_n \\ | & | & & | \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = y_1 \begin{bmatrix} | \\ x_1 \\ | \end{bmatrix} + y_2 \begin{bmatrix} | \\ x_2 \\ | \end{bmatrix} + \dots + y_n \begin{bmatrix} | \\ x_n \\ | \end{bmatrix}$$

Recall: Matrix \times matrix ($X^T X = \sum_{i=1}^n x_i x_i^T$)

$$\begin{bmatrix} | & | & \dots & | \\ x_1 & x_2 & \dots & x_n \\ | & | & & | \end{bmatrix} \begin{bmatrix} -x_1^T - \\ -x_2^T - \\ \vdots \\ -x_n^T - \end{bmatrix} = x_1 x_1^T + \dots + x_n x_n^T.$$

LEAST SQUARES LINEAR REGRESSION: KEY EQUATIONS

Two notations for the *key equation*

$$w_{\text{LS}} = \left(\sum_{i=1}^n x_i x_i^T \right)^{-1} \left(\sum_{i=1}^n y_i x_i \right) \iff w_{\text{LS}} = (X^T X)^{-1} X^T y.$$

Making Predictions

We use w_{LS} to make predictions.

Given x_{new} , the least squares prediction for y_{new} is

$$y_{\text{new}} \approx x_{\text{new}}^T w_{\text{LS}}$$

LEAST SQUARES SOLUTION

Potential issues

Calculating $w_{\text{LS}} = (X^T X)^{-1} X^T y$ assumes $(X^T X)^{-1}$ exists.

When doesn't it exist?

Answer: When $X^T X$ is not a full rank matrix.


When is $X^T X$ full rank?

Answer: When the $n \times (d + 1)$ matrix X has at least $d + 1$ *linearly independent* rows. This means that any point in \mathbb{R}^{d+1} can be reached by a weighted combination of $d + 1$ rows of X .

Obviously if $n < d + 1$, we can't do least squares. If $(X^T X)^{-1}$ doesn't exist, there are an infinite number of possible solutions.


Takeaway: We want $n \gg d$ (i.e., X is “tall and skinny”).

BROADENING LINEAR REGRESSION




BROADENING LINEAR REGRESSION

$$y = w_0 + w_1x$$



BROADENING LINEAR REGRESSION

$$y = w_0 + w_1x + w_2x^2 + w_3x^3$$



POLYNOMIAL REGRESSION IN \mathbb{R}

Recall: Definition of linear regression


A regression method is called *linear* if the prediction f is a linear function of the unknown parameters w .

- ▶ Therefore, a function such as $y = w_0 + w_1x + w_2x^2$ is *linear* in w . The LS solution is the same, only the preprocessing is different.
- ▶ E.g., Let $(x_1, y_1) \dots (x_n, y_n)$ be the data, $x \in \mathbb{R}$, $y \in \mathbb{R}$. For a p th-order polynomial approximation, construct the matrix


$$\mathbf{X} = \begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^p \\ 1 & x_2 & x_2^2 & \dots & x_2^p \\ \vdots & & & \ddots & \\ 1 & x_n & x_n^2 & \dots & x_n^p \end{bmatrix}$$

- ▶ Then solve exactly as before: $w_{\text{LS}} = (X^T X)^{-1} X^T y$.


POLYNOMIAL REGRESSION (M TH ORDER)




POLYNOMIAL REGRESSION (M TH ORDER)



POLYNOMIAL REGRESSION (M TH ORDER)



POLYNOMIAL REGRESSION (M TH ORDER)




POLYNOMIAL REGRESSION IN TWO DIMENSIONS

Example: 2nd and 3rd order polynomial regression in \mathbb{R}^2


The width of X grows as $(\text{order}) \times (\text{dimensions}) + 1$.

2nd order: $y_i = w_0 + w_1x_{i1} + w_2x_{i2} + w_3x_{i1}^2 + w_4x_{i2}^2$

3rd order: $y_i = w_0 + w_1x_{i1} + w_2x_{i2} + w_3x_{i1}^2 + w_4x_{i2}^2 + w_5x_{i1}^3 + w_6x_{i2}^3$



(a) 1st order



(b) 3rd order

FURTHER EXTENSIONS

More generally, for $x_i \in \mathbb{R}^{d+1}$ least squares linear regression can be performed on functions $f(x_i; w)$ of the form

$$y_i \approx f(x_i, w) = \sum_{s=1}^S g_s(x_i) w_s.$$

For example,

$$\begin{aligned} g_s(x_i) &= x_{ij}^2 \\ g_s(x_i) &= \log x_{ij} \\ g_s(x_i) &= \mathbb{I}(x_{ij} < a) \\ g_s(x_i) &= \mathbb{I}(x_{ij} < x_{ij'}) \end{aligned}$$

As long as the function is *linear* in w_1, \dots, w_S , we can construct the matrix X by putting the transformed x_i on row i , and solve $w_{\text{LS}} = (X^T X)^{-1} X^T y$.

One caveat is that, as the number of functions increases, we need more data to avoid overfitting.

GEOMETRY OF LEAST SQUARES REGRESSION

Thinking geometrically about least squares regression helps a lot.

- ▶ We want to minimize $\|y - Xw\|^2$. Think of the vector y as a point in \mathbb{R}^n . We want to find w in order to get the product Xw close to y .
- ▶ If X_j is the j th *column* of X , then $Xw = \sum_{j=1}^{d+1} w_j X_j$.
- ▶ That is, we weight the columns in X by values in w to approximate y .
- ▶ The LS solutions returns w such that Xw is as close to y as possible in the Euclidean sense (i.e., intuitive “direct-line” distance).

GEOMETRY OF LEAST SQUARES REGRESSION

$$\arg \min_w \|y - Xw\|^2 \Rightarrow w_{\text{LS}} = (X^T X)^{-1} X^T y.$$


The columns of X define a $d + 1$ -dimensional subspace in the higher dimensional \mathbb{R}^n .

The closest point in that subspace is the *orthonormal projection* of y into the *column space* of X .


Right: $y \in \mathbb{R}^3$ and data $x_i \in \mathbb{R}$.

$$X_1 = [1, 1, 1]^T \text{ and } X_2 = [x_1, x_2, x_3]^T$$


The approximation is $\hat{y} = Xw_{\text{LS}} = X(X^T X)^{-1} X^T y$.



GEOMETRY OF LEAST SQUARES REGRESSION



(a) $y_i \approx w_0 + x_i^T w$ for $i = 1, \dots, n$



(b) $y \approx Xw$

There are some key difference between (a) and (b) worth highlighting as you try to develop the corresponding intuitions.

(a) Can be shown for all n , but only for $x_i \in \mathbb{R}^2$ (not counting the added 1).

(b) This corresponds to $n = 3$ and one-dimensional data: $X = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \end{bmatrix}$.