

Altered Deoxyribonucleotide Pools in P2 Eductants of *Escherichia coli* K-12 due to Deletion of the *dcd* Gene

JAN NEUHARD* AND ELISABETH THOMASSEN

Enzyme Division, University Institute of Biological Chemistry B, DK-1307 Copenhagen K, Denmark

Received for publication 18 December 1975

Deletions of the *Escherichia coli* K-12 chromosome associated with P2-mediated eduction extend through the structural gene for uridine kinase, *udk*, and the *dcd* gene encoding 2'-deoxycytidine 5'-triphosphate deaminase. The lack of uridine kinase makes a positive selection possible for these strains. Due to the *dcd* mutation, P2 eductants show large alterations in their deoxyribonucleoside triphosphate pools.

Escherichia coli K-12 lysogenic for P2 in the H site segregates histidine-requiring eductants, which contain deletions extending from the *attP2_H* gene through the *sbcB*, *his*, *gnd*, and *rfb* genes into the *mglP* gene (13).

Recently we isolated mutants of *Salmonella typhimurium* lacking 2'-deoxycytidine 5'-triphosphate (dCTP) deaminase (EC 3.5.4..) activity, i.e., *dcd* mutants. Transduction studies, using P22, showed that *dcd* cotransduces 95% with the structural gene *udk*, coding for uridine kinase (EC 2.7.1.48) (C.F. Beck, J. Neuhard, and E. Thomassen, unpublished data). On the *S. typhimurium* chromosome, *udk* is located between *metG* and *hisE*. With phage P1, *udk* cotransduces 23% with *metG* and about 12% with *hisE* (3). In *E. coli*, P1 transduction has indicated about 10% cotransduction between *udk* and *his* (7). These results suggest that the host deletions obtained by P2-mediated eduction in *E. coli* might extend through the *udk* and *dcd* genes. Mutants of *E. coli* defective in dCTP deaminase have been described; however, they were never characterized genetically (10).

Mutants defective in dCTP deaminase have no easily detectable phenotype (10). However, *udk* mutants are resistant to 5-fluorouridine, provided that the strain is already resistant to 5-fluorouracil, i.e., is defective in uracil phosphoribosyltransferase (EC 2.4.2.9) (*upp*) (4).

E. coli K-12 strain S0864 (F⁻, *trp*, *lacZ*, *strA*, *thi*, *upp*) was infected with P2 and plated in LB soft agar (0.7% agar) on LB agar plates (1.5% agar). Cells from the center of six plaques were combined and grown overnight in LB. Aliquots of this culture were spread on minimal agar plates containing basal salts (5), 0.2% glucose, tryptophan (40 µg/ml), histidine (40 µg/ml), thiamine (5 µg/ml), 5-fluorouridine (10 µg/ml) (F. Hoffman-La Roche & Co., Ltd., Basel) and

1.5% agar (Difco). The plates were incubated for 36 h at 37 C, and the resistant colonies appearing were tested for histidine requirement. Most of the resistant colonies had simultaneously acquired a histidine requirement. Five were kept for further studies (S0874, S0875, S0876, S0877, and S0878). All five had become sensitive to P2 and did not produce P2 by spotting on susceptible indicator strains.

Table 1 shows the results of enzyme assays

TABLE 1. Uridine kinase and dCTP deaminase activities in extracts^a of different *E. coli* K-12 strains

| Strain | His phenotype | Sp act ^b | |
|-----------------|------------------|-----------------------------|-----------------------------|
| | | Uridine kinase ^c | dCTP deaminase ^d |
| S0864 | His ⁺ | 12.2 | 2.03 |
| S0874 | His ⁻ | <0.1 | <0.05 |
| S0875 | His ⁻ | <0.1 | <0.05 |
| S0876 | His ⁻ | <0.1 | <0.05 |
| S0877 | His ⁻ | <0.1 | <0.05 |
| S0878 | His ⁻ | <0.1 | <0.05 |
| HfrH <i>thi</i> | His ⁺ | 19.5 | 2.10 |
| QE1 | His ⁻ | <0.1 | <0.05 |

^a Cells were grown overnight at 37 C with aeration in AB medium (5) containing 0.2% glucose, 0.2% Casamino Acids (Difco), thiamine (5 µg/ml), and tryptophan (40 µg/ml). They were harvested by centrifugation, washed once with 0.9% NaCl, and kept frozen until assayed. For uridine kinase assays, cells were suspended in a solution of 0.1 M tris(hydroxymethyl)aminomethane (Tris)-chloride (pH 7.8) and 1 mM ethylenediaminetetraacetate (EDTA), sonically disrupted, and centrifuged. The supernatants were dialyzed for 2 h against 100 volumes of the sonication buffer and assayed for uridine kinase activity. For dCTP deaminase assays, cells were suspended in a solution of 0.05 M potassium phosphate buffer (pH 6.8) and 2 mM EDTA, sonically disrupted, and centrifuged. The supernatants were treated with streptomycin sulfate at 4 C (final concentration, 3%), left standing for 30 min,

TABLE 1—Continued

and centrifuged. The supernatants were made 55% saturated with solid ammonium sulfate, left standing for 30 min at 4 C, and centrifuged again. The precipitates were dissolved in a solution of 0.05 M potassium phosphate buffer (pH 6.8), 2 mM EDTA, and 2 mM mercaptoethanol and dialyzed for 2 h against 100 volumes of the same buffer.

^b Nanomoles of substrate converted per minute per milligram of protein. Proteins were determined by the method of Lowry (8).

^c Uridine kinase assay. Reaction mixtures of 40 μ l contained 0.05 M Tris-chloride (pH 7.8), 2.5 mM MgCl₂, 3 mM guanosine 5'-triphosphate, 1 mM [2-¹⁴C]uridine (specific activity, 0.5 μ Ci/ μ mol), and extract. Reactions were started by the addition of extracts and were performed at 37 C. At 5, 10, and 15 min, 10- μ l samples were applied to the "starting" spots of poly(ethyleneimine)-impregnated cellulose thin-layer plates (12) and dried with hot air. The chromatograms were developed for 12 cm with water. After the chromatograms were dry, the starting spots containing phosphorylated uridine compounds were cut out and counted in a Packard Tri-Carb liquid scintillation spectrometer.

^d dCTP deaminase activity was determined by the spectrophotometric assay described by Beck et al. (2).

performed on S0864 and the five eductants S0874 through 878. All five mutants are deficient in both uridine kinase and dCTP deaminase. Similarly, QE1, which is a P2 eductant of *E. coli* HfrH *thi*, isolated by Sunshine and Kelly (13), is devoid of both activities (Table 1). Since QE1 was isolated as a *his*⁻ eductant with-




FIG. 1. Pyrimidine deoxyribonucleotide interconversion in *E. coli*. Symbols: \rightarrow , main pathway for dUMP synthesis; \cdots , secondary pathway for dUMP synthesis. The enzymes are identified by numbers as follows: 1, ribonucleoside diphosphate reductase (EC 1.8.); 2, nucleoside diphosphate kinase (EC 2.7.4.6); 3, dCTP deaminase (EC 3.5.4.); 4, deoxyuridine 5'-triphosphatase (EC 3.1.); 5, thymidylate synthetase (EC 2.1.1.b); 6, thymidylate kinase (EC 2.7.4.9). The gene designations for the corresponding structural genes, where known, are underlined.

out employing 5-fluorouridine, we conclude that the *udk* and *dcd* genes are located on that piece of deoxyribonucleic acid which is lost concomitant to the exit of the P2 phage from its H site.

dCTP deaminase is the key enzyme of the main pathway for the synthesis of 2'-deoxyuridine 5'-monophosphate (dUMP), the ultimate precursor for thymidine nucleotide synthesis (2) (Fig. 1). Thus, *dcd* mutants can only synthesize dUMP via the second pathway, which in-

TABLE 2. Nucleoside triphosphate pools^a in different *E. coli* K-12 strains

| Strain | Addition to the medium | Pool sizes ^b | | | | | | | |
|--------|-----------------------------------|-------------------------|------|------|------|------|------|------|------|
| | | ATP | GTP | CTP | UTP | dATP | dGTP | dCTP | dTTP |
| S0864 | None | 4.84 | 2.59 | 1.07 | 1.24 | 0.23 | 0.08 | 0.17 | 0.20 |
| | UdR ^b (100 μ g/ml) | 6.93 | 3.51 | 1.81 | 1.72 | 0.30 | 0.20 | 0.25 | 0.54 |
| | Thymidine (100 μ g/ml) | 5.87 | 2.97 | 1.47 | 1.74 | 0.24 | 0.08 | 0.22 | 0.44 |
| | Thymine (50 μ g/ml) | 7.34 | 3.43 | 1.87 | 2.14 | 0.30 | 0.12 | 0.30 | 0.30 |
| S0874 | None | 4.97 | 2.25 | 1.00 | 1.56 | 0.22 | 0.07 | 7.43 | 0.07 |
| | UdR ^b (100 μ g/ml) | 3.87 | 2.06 | 1.27 | 1.13 | 0.17 | 0.11 | 0.79 | 0.72 |
| | Thymidine (100 μ g/ml) | 4.28 | 2.10 | 1.23 | 1.35 | 0.11 | 0.05 | 1.07 | 0.89 |
| | Thymine (50 μ g/ml) | 2.90 | 1.23 | 0.69 | 1.03 | 0.12 | 0.06 | 4.39 | 0.08 |

^a S0864 and S0874 were grown at 37 C in low-phosphate medium (6) containing 0.2% glucose, tryptophan (40 μ g/ml), histidine (40 μ g/ml), thiamine (5 μ g/ml), [³²P]orthophosphate (specific activity, 10 μ Ci/ μ mol), and the additions mentioned in the table. After two generations of exponential growth, 5 ml of each culture was filtered on membrane filters (Millipore Corp.), and the filters were extracted with ice-cold 0.3 N perchloric acid. The quantitative thin-layer chromatographic determination of individual nucleoside triphosphates in the extracts was described previously (9).

^b Pools are given in micromoles per gram of bacterial dry weight. ATP, Adenosine 5'-triphosphate; UTP, uridine 5'-triphosphate; dATP, 2'-deoxyadenosine 5'-triphosphate; dGTP, 2'-deoxyguanosine 5'-triphosphate; UdR, 2'-deoxyuridine.

volves the reduction of uridine 5'-diphosphate (UDP) to dUDP (Fig. 1), and they are therefore stressed in their ability to synthesize thymidine nucleotides (10). The P2 eductant S0874 is indeed limited in its 2'-deoxythymidine 5'-triphosphate (dTTP) pool (Table 2). Concomitantly with the low-dTTP pool the dCTP pool is increased about 20-fold. This agrees with the previous studies on the effect of thymine starvation on the deoxyribonucleoside triphosphate pools of *thy* mutants of *E. coli* (9).

Since the low-dTTP pool of the eductants is due to a decrease in the availability of the precursor dUMP, one would expect that the addition of either thymidine or deoxyuridine to the growth medium of the mutants would increase the dTTP pool and, as a result, restore the dCTP pool to normal values. This is, indeed, what is observed (Table 2). However, the addition of even high concentrations of thymine is not capable of normalizing the deoxyribonucleotide pools of S0874 (Table 2). This latter result is to be expected, since the deoxyribose-1-phosphate required for the utilization of exogenous thymine by *dcd*⁺ strains is derived from dUMP (1) which, in the eductants, is the limiting compound for thymidine nucleotide synthesis.

The somewhat lower growth rate observed with P2 eductants as compared with the parental strains is not normalized by the addition of either thymidine or deoxyuridine to the growth medium.

The present work has extended the number of genes known to be deleted from the *E. coli* K-12 chromosome after eduction of P2. Of particular interest is the finding that the deletion covers the *dcd* gene, resulting in partial thymidine nucleotide starvation of the eductants. This, in turn, causes alterations in the deoxyribonucleotide pools (Table 2), elongations of the cells (unpublished data), and, eventually, a decrease in the replication time of the chromosome (11).

We would like to thank R. Buxton, who supplied us with the P2 phage, and O. Karlström for strain QE1.

LITERATURE CITED

1. Beacham, I. R., and R. H. Pritchard. 1971. The role of nucleoside phosphorylases in the degradation of deoxyribonucleosides by thymine requiring mutants of *E. coli*. *Mol. Gen. Genet.* 110:289-298.
2. Beck, C. F., A. R. Eisenhardt, and J. Neuhaard. 1975. Deoxycytidine triphosphate deaminase of *Salmonella typhimurium*. *J. Biol. Chem.* 250:609-616.
3. Beck, C. F., J. L. Ingraham, and J. Neuhaard. 1972. Location on the chromosome of *Salmonella typhimurium* of genes governing pyrimidine metabolism. II. Uridine kinase, cytosine deaminase and thymidine kinase. *Mol. Gen. Genet.* 115:208-215.
4. Beck, C. F., J. L. Ingraham, J. Neuhaard, and E. Thomassen. 1972. Metabolism of pyrimidines and pyrimidine nucleosides by *Salmonella typhimurium*. *J. Bacteriol.* 110:219-228.
5. Clark, D. J., and O. Maaløe. 1967. DNA replication and the division cycle in *Escherichia coli*. *J. Mol. Biol.* 23:99-112.
6. Edlin, G., and O. Maaløe. 1966. Synthesis and breakdown of messenger RNA without protein synthesis. *J. Mol. Biol.* 15:428-434.
7. Hammer-Jespersen, K., and A. Munch-Petersen. 1973. Mutants of *Escherichia coli* unable to metabolize cytidine: isolation and characterization. *Mol. Gen. Genet.* 126:177-186.
8. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.
9. Neuhaard, J., and E. Thomassen. 1971. Turnover of the deoxyribonucleoside triphosphates in *Escherichia coli* 15 T⁻ during thymine starvation. *Eur. J. Biochem.* 20:36-43.
10. O'Donovan, G. A., G. Edlin, J. A. Fuchs, J. Neuhaard, and E. Thomassen. 1971. Deoxycytidine triphosphate deaminase: characterization of an *Escherichia coli* mutant deficient in the enzyme. *J. Bacteriol.* 105:666-672.
11. Pritchard, R. H., and A. Zaritsky. 1970. Effect of thymine concentration on the replication velocity of DNA in a thymineless mutant of *Escherichia coli*. *Nature (London)* 226:126-131.
12. Randerath, K., and E. Randerath. 1966. Ion-exchange thin-layer chromatography. XV. Preparation, properties and applications of paper-like PEI-cellulose sheets. *J. Chromatogr.* 22:110-117.
13. Sunshine, M. G., and B. Kelly. 1971. Extent of host deletions associated with bacteriophage P2-mediated eduction. *J. Bacteriol.* 108:695-704.