COMP 576 - Fall 2017 Assignment 1

September 30, 2017

Backpropagation in a Simple Neural Network

- 1a) Dataset
- 1b) Derivatives of Activation Functions

Sigmoid:

$$f(x) = \frac{1}{1+e^{-x}} = (1+e^{-x})^{-1}$$

$$\frac{d(1+e^{-x})^{-1}}{dx} = (1+e^{-x})^{-2}(e^{-x})$$

$$= \frac{e^{-x}}{(1+e^{-x})^2} = \frac{1}{1+e^{-x}} \frac{e^{-x}}{1+e^{-x}}$$

$$= \frac{1}{1+e^{-x}} \frac{(1+e^{-x})-1}{1+e^{-x}}$$

$$= \frac{1}{1+e^{-x}} \left(1 - \frac{1}{1+e^{-x}}\right)$$

$$\frac{d(1+e^{-x})^{-1}}{dx} = f(x)(1-f(x))$$

Tanh:

$$f(x) = tanh(x) = \frac{sinh(x)}{cosh(x)}$$

$$\frac{d\left(\frac{sinh(x)}{cosh(x)}\right)}{dx} = \frac{cosh(x)cosh(x) - sinh(x)sinh(x)}{cosh^2(x)}$$

$$= \frac{cosh^2(x) - sinh^2(x)}{cosh^2(x)} = \frac{cosh^2(x)}{cosh^2(x)} - \frac{sinh^2(x)}{cosh^2(x)}$$

$$\frac{d\left(\frac{sinh(x)}{cosh(x)}\right)}{dx} = 1 - tanh^2(x)$$

ReLu:

$$f(x) = max(0, x)$$

$$f(x) = \begin{cases} x, & x > 0\\ 0, & \text{otherwise} \end{cases}$$

$$f'(x) = \begin{cases} 1, & x > 0 \\ 0, & \text{otherwise} \end{cases}$$

1c) Building the Neural Network

Three Layer Network

$$z^1 = W^1 x + b^1 (1)$$

$$a^1 = actFun(z^1) (2)$$

$$z^2 = W^2 a^1 + b^2 (3)$$

$$a^2 = \hat{y} = softmax(z^2) \tag{4}$$

Mean Cross Entropy Loss of Batch

$$L(y, \hat{y}) = -\frac{1}{N} \sum_{n=1}^{N} \sum_{i=1}^{C} y_{n,i} \log \hat{y}_{n,i}$$
 (5)

1d) Backward Pass - Backpropagation

Gradients: $\frac{\partial L}{\partial W^2}$, $\frac{\partial L}{\partial b^2}$, $\frac{\partial L}{\partial W^1}$, $\frac{\partial L}{\partial b^1}$

$$\frac{\partial L}{\partial W^2} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z^2} \frac{\partial z^2}{\partial W^2}$$

$$\frac{\partial z^2}{\partial W^2} = \frac{\partial (W^2 a^2 + b^2)}{\partial W^2}$$

$$\frac{\partial z^2}{\partial W^2} == a^2$$

$$\frac{\partial \hat{y_i}}{\partial z_i^2} = \frac{\partial \left(\frac{e^{z_i^2}}{\sum_{j=1}^C e^{z_j^2}}\right)}{\partial z_i^2}$$

if
$$j = i$$

$$=\frac{e^{z_i^2}\sum_{j=1}^C e^{z_j^2}-e^{z_i^2}e^{z_j^2}}{\left(\sum_{j=1}^C e^{z_j^2}\right)^2}=\frac{e^{z_i^2}(\sum_{j=1}^C e^{z_j^2}-e^{z_j^2})}{\sum_{j=1}^C e^{z_j^2}\sum_{j=1}^C e^{z_j^2}}$$

$$rac{\partial \hat{y_i}}{\partial z_i^2} = \hat{y_i}(1 - \hat{y_i})$$

if
$$j \neq i$$

$$\frac{\partial \hat{y}_i}{\partial z_j^2} = \frac{0 - e^{z_i^2} e^{z_j^2}}{(\sum_{i=1}^C e^{z_i^2})^2}$$

$$rac{\partial \hat{y_i}}{\partial z_j^2} = -\hat{y_i}\hat{y_j}$$

$$\frac{\partial L}{\partial \hat{y}_i} = -\frac{1}{N} \sum_{n=1}^{N} \sum_{i=1}^{C} \frac{\partial (y_i \log \hat{y}_i)}{\partial \hat{y}_i} = -\frac{1}{N} \sum_{n=1}^{N} \sum_{i=1}^{C} y_i \left(\frac{1}{\hat{y}_i}\right) \frac{\partial \hat{y}_i}{\partial z_i^2}$$

 $\forall i$

$$= -\frac{1}{N} \sum_{n=1}^{N} \left[\frac{y_i}{\hat{y_i}} (1 - \hat{y_i}) + \sum_{j \neq i}^{C} \frac{y_j}{\hat{y_j}} (-\hat{y_j} \hat{y_i}) \right]$$

$$= -\frac{1}{N} \sum_{n=1}^{N} \left[y_i (1 - \hat{y_i}) - \sum_{j \neq i}^{C} y_j \hat{y_i} \right]$$

$$= \frac{1}{N} \sum_{n=1}^{N} \left[-y_i + y_i \hat{y_i} + \sum_{j \neq i}^{C} y_j \hat{y_i} \right]$$

$$= \frac{1}{N} \sum_{n=1}^{N} \left[-y_i + \sum_{j=1}^{C} y_j \hat{y}_i \right]$$

= since y_j is one-hot encoded

$$\implies \sum_{j=1}^{C} y_j = 1$$

$$\therefore rac{\partial L}{\partial z_i^2} = rac{1}{N} \sum_{n=1}^N (\hat{y_i} - y_i)$$

Let
$$\delta^3 = (\hat{y_i} - y_i)$$

$$\frac{\partial L}{\partial W^2} = \frac{1}{N} \sum_{n=1}^{N} \delta^3 a^3$$

$$\frac{\partial L}{\partial W^2} = \frac{1}{N} a^{1T} \delta^3$$

$$\frac{\partial L}{\partial b^2} = \frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z^2} \frac{\partial z^2}{\partial b^2}$$

$$\frac{\partial z^2}{\partial b^2} = 1$$

$$rac{\partial L}{\partial b^2} = rac{1}{N} \sum_{n=1}^C \delta^3$$

$$\frac{\partial L}{\partial W^1} = \underbrace{\frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z^2}}_{\frac{1}{N} \sum_{i=1}^{N} \delta^3} \underbrace{\frac{\partial z^2}{\partial a^1}}_{f'(z^1)} \underbrace{\frac{\partial a^1}{\partial z^1}}_{f'(z^1)} \underbrace{\frac{\partial z^1}{\partial W^1}}_{x}$$

$$=\frac{1}{N}W^{2T}\delta^3f'(z^1)$$

Let
$$\delta^2 = W^{2T} \delta^3 f'(z^1)$$

$$=\frac{1}{N}\delta^2x$$

$$rac{\partial L}{\partial W^1} = rac{1}{N} x^T \delta^2$$

$$\frac{\partial L}{\partial b^1} = \underbrace{\frac{\partial L}{\partial \hat{y}} \frac{\partial \hat{y}}{\partial z^2} \frac{\partial z^2}{\partial a^1} \frac{\partial a^1}{\partial z^1}}_{\frac{1}{N} \delta^2} \underbrace{\frac{\partial z^1}{\partial b^1}}_{1}$$

$$rac{\partial L}{\partial b^1} = rac{1}{N} \delta^2$$