



ATLAS Note

GROUP-2017-XX

8th November 2020



Draft version 0.1

1

2

3

4

A Deep Learning Approach to Differential Measurements of $t\bar{t}H$ Production in Multilepton Final States

5

The ATLAS Collaboration

6

7

8

9

10

11

12

The possibility of using the kinematic properties of the Higgs boson to search for new physics is investigated using $t\bar{t}H$ events with multiple leptons in the final state. Because of the challenges inherent to reconstructing the Higgs in multilepton final states, a deep learning approach is used to reconstruct the momentum spectrum of the Higgs, which would be altered by the presence of new physics without affecting the overall rate of $t\bar{t}H$ production. Simulations representing 139 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$ are used to provide estimates of the sensitivity to variations in the Higgs p_T spectrum.

Contents

16	1	Changes and outstanding items	3
17	1.1	Changelog	3
18	2	Introduction	4
19	3	Data and Monte Carlo Samples	4
20	3.1	Data Samples	5
21	3.2	Monte Carlo Samples	5
22	4	Object Reconstruction	6
23	4.1	Trigger Requirements	6
24	4.2	Light Leptons	6
25	4.3	Jets	7
26	4.4	Missing Transverse Energy	8
27	5	Higgs Momentum Reconstruction	8
28	5.1	Decay Candidate Reconstruction	8
29	5.2	b-jet Identification	9
30	5.3	Higgs Reconstruction	12
31	5.4	p_T Prediction	20
32	5.5	3l Decay Mode	25
33	6	Signal Region Definitions	27
34	6.1	Pre-MVA Event Selection	27
35	6.2	Event MVA	28
36	6.3	Signal Region Definitions	29
37	7	Systematic Uncertainties	30
38	8	Results	32
39	9	Conclusion	32
40		Appendices	34
41	A	Machine Learning Models	34
42	A.1	Background Rejection MVAs	34

43 **1 Changes and outstanding items**

44 **1.1 Changelog**

45 This is version 1

2 Introduction

Since the discovery of a Higgs boson compatible with the Standard Model (SM) in 2012 [], its interactions with other particles have been studied using proton-proton collision data produced by the Large Hadron Collider (LHC). The strongest of these interactions is the coupling of the Higgs to the top quark, making the Yukawa coupling between these two particles of particular interest for study.

These interactions can be measured directly by studying the production of a Higgs Boson in association with a pair of Top Quarks ($t\bar{t}H$) []. While this process has been observed by both the ATLAS [] and CMS [] collaborations, these analyses have focused on measuring the overall rate of $t\bar{t}H$ production. There are several theories of physics Beyond the Standard Model (BSM), however, that would affect the kinematics of $t\bar{t}H$ production without altering its overall rate [].

An Effective Field Theory approach can be used to model the low energy effects of new, high energy physics, by parameterizing BSM effects as dimension-six operators. The addition of these operators can be shown to modify the transverse momentum (p_T) spectrum of the Higgs Boson []. Therefore, reconstructing the momentum spectrum of the Higgs provides a means to observe new physics in the Higgs sector.

This note reports on the feasibility of measuring the impact of dimension-six operators in $t\bar{t}H$ events with multiple leptons in the final state, using Monte Carlo (MC) simulations scaled to 139 fb^{-1} at an energy $\sqrt{s} = 13 \text{ TeV}$. Events are separated into channels based on the number of light leptons (electrons and muons) in the final state - either two same-sign leptons (2LSS), or three leptons (3L). A deep neural network is used to identify which objects originate from the decay of the Higgs, and reconstruct the momentum of the Higgs Boson in each event. This reconstructed momentum spectrum is used to place limits on BSM effects, and on the parameters of dimension-six operators.

This note is organized as follows: Section ?? describes the LHC and the ATLAS detector. The dataset and Monte Carlo (MC) simulations used in the analysis is outlined in section 3. Section 4 describes the identification and reconstruction of the various physics objects. The models used to reconstruct the momentum spectrum of the Higgs is discussed in section 5. The selection and categorisation of events comprises section 6, and the theoretical and experimental systematic uncertainties considered are described in section 7. Finally, the results of the study are summarized in section 8.

3 Data and Monte Carlo Samples

For both data and Monte Carlo (MC) simulations, samples were prepared in the xAOD format, which was used to produce a xAOD based on the HIGG8D1 derivation framework. This framework was designed for the main $t\bar{t}H$ multi-lepton analysis. Because this analysis targets events with

multiple light leptons, as well as tau hadrons, this framework skims the dataset of any events that do not meet at least one of the following requirements:

- at least two light leptons within a range $|\eta| < 2.6$, with leading lepton $p_T > 15$ GeV and subleading lepton $p_T > 5$ GeV
- at least one light lepton with $p_T > 15$ GeV within a range $|\eta| < 2.6$, and at least two hadronic taus with $p_T > 15$ GeV.

Samples were then generated from these HIGG8D1 derivations using a modified version of AnalysisBase version 21.2.127.

3.1 Data Samples

The study uses proton-proton collision data collected by the ATLAS detector from 2015 through 2018, which represents an integrated luminosity of 139 fb^{-1} and an energy of $\sqrt{s} = 13$ TeV. All data used in this analysis was included in one the following Good Run Lists:

- data15_13TeV.periodAllYear_DetStatus-v79-repro20-02_DQDefects-00-02-02_PHYS_StandardGRL_All_Good_25ns.xml
- data16_13TeV.periodAllYear_DetStatus-v88-pro20-21_DQDefects-00-02-04_PHYS_StandardGRL_All_Good_25ns.xml
- data17_13TeV.periodAllYear_DetStatus-v97-pro21-13_Unknown_PHYS_StandardGRL_All_Good_25ns_TriggerNo17e33prim.xml
- data18_13TeV.periodAllYear_DetStatus-v102-pro22-04_Unknown_PHYS_StandardGRL_All_Good_25ns_TriggerNo17e33prim.xml

3.2 Monte Carlo Samples

Several Monte Carlo (MC) generators were used to simulate both signal and background processes. For all of these, the effects of the ATLAS detector are simulated in Geant4. The specific event generator used for each of these MC samples is listed in table 1.

Table 1: The configurations used for event generation of signal and background processes, including the event generator, matrix element (ME) order, parton shower algorithm, and parton distribution function (PDF).

Process	Event generator	ME order	Parton Shower	PDF
$t\bar{t}H$	MG5_AMC (MG5_AMC)	NLO (NLO)	PYTHIA 8 (HERWIG++)	NNPDF 3.0 NLO [Ball:2014uwa] (CT10 [ct10])
$t\bar{t}W$	MG5_AMC (SHERPA 2.1.1)	NLO (LO multileg)	PYTHIA 8 (SHERPA)	NNPDF 3.0 NLO (NNPDF 3.0 NLO)
$t\bar{t}(Z/\gamma^* \rightarrow ll)$	MG5_AMC	NLO	PYTHIA 8	NNPDF 3.0 NLO
VV	SHERPA 2.2.2	MEPS NLO	SHERPA	CT10
$t\bar{t}$	POWHEG-BOX v2 [powheggt]	NLO	PYTHIA 8	NNPDF 3.0 NLO
$t\bar{t}\gamma$	MG5_AMC	LO	PYTHIA 8	NNPDF 2.3 LO
tZ	MG5_AMC	LO	PYTHIA 6	CTEQ6L1
$tHqb$	MG5_AMC	LO	PYTHIA 8	CT10
tHW	MG5_AMC (SHERPA 2.1.1)	NLO (LO multileg)	HERWIG++ (SHERPA)	CT10 (NNPDF 3.0 NLO)
tWZ	MG5_AMC	NLO	PYTHIA 8	NNPDF 2.3 LO
$t\bar{t}t, t\bar{t}t\bar{t}$	MG5_AMC	LO	PYTHIA 8	NNPDF 2.3 LO
$t\bar{t}W^+W^-$	MG5_AMC	LO	PYTHIA 8	NNPDF 2.3 LO
$s-, t\text{-channel},$ Wt single top	POWHEG-BOX v1 [powhegstp]	NLO	PYTHIA 6	CT10
$q\bar{q}VV, VVV$ $Z \rightarrow l^+l^-$	SHERPA 2.2.1	MEPS NLO	SHERPA	NNPDF 3.0 NLO

4 Object Reconstruction

All analysis channels considered in this note share a common object selection for leptons and jets, as well as a shared trigger selection.

4.1 Trigger Requirements

Events are required to be selected by dilepton triggers, as summarized in table 2.

4.2 Light Leptons

Electron candidates are reconstructed from energy clusters in the electromagnetic calorimeter that are associated with charged particle tracks reconstructed in the inner detector [ATLAS-CONF-2016-024]. Electron candidates are required to have $p_T > 10$ GeV and $|\eta_{\text{cluster}}| < 2.47$. Candidates in the transition region between different electromagnetic calorimeter components, $1.37 < |\eta_{\text{cluster}}| < 1.52$, are rejected. A multivariate likelihood discriminant combining shower shape and track information is used to distinguish prompt electrons from nonprompt leptons, such as those originating from hadronic showers.

Dilepton triggers (2015)	
$\mu\mu$ (asymm.)	HLT_mu18_mu8noL1
ee (symm.)	HLT_2e12_lhloose_L12EM10VH
$e\mu, \mu e$ (~symm.)	HLT_e17_lhloose_mu14
Dilepton triggers (2016)	
$\mu\mu$ (asymm.)	HLT_mu22_mu8noL1
ee (symm.)	HLT_2e17_lhvloose_nod0
$e\mu, \mu e$ (~symm.)	HLT_e17_lhloose_nod0_mu14
Dilepton triggers (2017)	
$\mu\mu$ (asymm.)	HLT_mu22_mu8noL1
ee (symm.)	HLT_2e24_lhvloose_nod0
$e\mu, \mu e$ (~symm.)	HLT_e17_lhloose_nod0_mu14
Dilepton triggers (2018)	
$\mu\mu$ (asymm.)	HLT_mu22_mu8noL1
ee (symm.)	HLT_2e24_lhvloose_nod0
$e\mu, \mu e$ (~symm.)	HLT_e17_lhloose_nod0_mu14

Table 2: List of lowest p_T -threshold, un-prescaled dilepton triggers used for 2015-2018 data taking.

To further reduce the non-prompt contribution, the track of each electron is required to originate from the primary vertex; requirements are imposed on the transverse impact parameter significance ($|d_0|/\sigma_{d_0}$) and the longitudinal impact parameter ($|\Delta z_0 \sin \theta_\ell|$), as shown in table ??.

Muon candidates are reconstructed by combining inner detector tracks with track segments or full tracks in the muon spectrometer [PERF-2014-05]. Muon candidates are required to have $p_T > 10$ GeV and $|\eta| < 2.5$. All leptons are required to be isolated, and pass a non-prompt BDT selection described in detail in [ttH_paper].

4.3 Jets

Jets are reconstructed from calibrated topological clusters built from energy deposits in the calorimeters [ATL-PHYS-PUB-2015-015], using the anti- k_t algorithm with a radius parameter $R = 0.4$. Jets with energy contributions likely arising from noise or detector effects are removed from consideration [ATLAS-CONF-2015-029], and only jets satisfying $p_T > 25$ GeV and $|\eta| < 2.5$ are used in this analysis. For jets with $p_T < 60$ GeV and $|\eta| < 2.4$, a jet-track association algorithm is used to confirm that the jet originates from the selected primary vertex, in order to reject jets arising from pileup collisions [PERF-2014-03].

4.4 Missing Transverse Energy

Because all $t\bar{t}H$ — ML channels considered include multiple neutrinos, missing transverse energy (E_T^{miss}) is present in each event. The missing transverse momentum vector is defined as the inverse of the sum of the transverse momenta of all reconstructed physics objects as well as remaining unclustered energy, the latter of which is estimated from low- p_T tracks associated with the primary vertex but not assigned to a hard object [ATL-PHYS-PUB-2015-027].

5 Higgs Momentum Reconstruction

Reconstructing the momentum of the Higgs boson is a particular challenge for channels with leptons in the final state: Because all channels include at least two neutrinos in the final state, the Higgs can never be fully reconstructed. However, the momentum spectrum can be well predicted by a neural network when provided with the four-vectors of the Higgs Boson decay products, as shown in section 5.1. With this in mind, several layers of MVAs are used to reconstruction the Higgs momentum.

The first layer is a model designed to select which jets are most likely to be the b-jets that came from the top decay, detailed in section 5.2. As described in section 5.3, the kinematics of these jets are fed into the second layer, which is designed to identify the decay products of the Higgs Boson itself. The kinematics of these particles are then fed into yet another neural-network, which predicts the momentum of the Higgs (5.4). MVAs are also used in the analysis to determine the decay of the Higgs boson in the 3l channel (5.5).

For all of these models, the Keras neural network framework, with Tensorflow as the backend, is used, and the number of hidden layers and nodes are determined using grid search optimization. Each neural network uses the LeakyReLU activation function, a learning rate of 0.01, and the Adam optimization algorithm, as alternatives are found to either decrease or have no impact on performance. Batch normalization is applied after each layer. For the classification algorithms (b-jet matching, Higgs reconstruction, and 3l decay identification) binary-cross entropy is used as the loss function, while the p_T reconstruction algorithm uses MSE.

The specific inputs features used for each model are arrived at through a process of trial and error - features considered potentially useful are tried, and those that are found to increase performance are included. While each model includes a relatively large number of features, some using upwards of 30, this inclusive approach is found to maximize the performance of each model while decreasing the variance compared to a reduced number of inputs. Each input feature is validated by comparing MC simulations to 80 fb^{-1} of data, as shown in the sections below.

5.1 Decay Candidate Reconstruction

Machine Learning algorithms are trained to identify the decay products of the Higgs Boson using MC simulations of $t\bar{t}H$ events. These include light leptons and jets. Reconstructed physics

objects are matched to truth level particles, in order to identify the parents of these reconstructed objects. The kinematics of the decay product candidates as well as event level variables are used as inputs.

Leptons considered as possible Higgs and top decay candidates are required to pass the selection described in section 4.2. For jets, however, it is found that a large fraction that originate from either the top decay or the Higgs decay fall outside the selection described in section 4.3. Specifically, jets from the Higgs decay tend to be soft, with 32% having $p_T < 25$ GeV. Therefore jets with $p_T < 15$ GeV are considered as possible candidates in the models described below. By contrast, less than 5% of the jets originating from the Higgs fall below this p_T . The jets are found to be well modeled even down to this low p_T threshold, as shown in section 6.1. The impact of using different p_T selection for the jet candidates is considered in detail in section ???. As they are expected to originate from the primary vertex, jets are also required to pass a JVT cut.

5.2 b-jet Identification

Including the kinematics of the b-jets that originate from the top decay is found to improve the identification of the Higgs decay products, and improve the accuracy with which the Higgs momentum can be reconstructed. Because these b-jets are reconstructed by the detector with high efficiency (just over 90% of the time), and can be identified relatively consistently, the first step in reconstructing the Higgs is selecting the b-jets from the top decay.

Exactly two b-jets are expected in the final state of $t\bar{t}H$ – ML events. However, in both the 3l and 2lSS channels, only one or more b-tagged jets are required (where the 70% DL1r b-tag working point is used). Therefore, for events which have exactly one, or more than two, b-tagged jets, deciding which combination of jets correspond to the top decay is non-trivial. Further, events with 1 b-tagged jet represent just over half of all $t\bar{t}H$ – ML events. Of those, both b-jets are reconstructed by the detector 75% of the time. Therefore, rather than adjusting the selection to require exactly 2 b-tagged jets, and losing more than half of the signal events, a neural network is used to predict which pair of jets is most likely to correspond to truth b-jets.

Once the network is trained, all possible pairings of jets are fed into the model, and the pair of jets with the highest output score are taken to be b-jets in successive steps of the analysis.

5.2.1 2lSS Channel

For the 2lSS channel, the input features shown in table 3 are used for training. Here j_0 and j_1 are the two jet candidates, while l_0 and l_1 are the two leptons in the event, ordered by p_T . jet DL1r is an integer corresponding to the calibrated b-tagging working points reached by each jet, where 5 represents the tightest working point and 1 represents the loosest. The variables nJets DL1r 60% and nJets DL1r 85% represent the number of jets in the event passing the 60% and 85% b-tag working points, respectively.

jet p_T 0	jet p_T 1	Lepton p_T 0
Lepton p_T 1	jet η 0	jet η 1
$\Delta R(j_0)(j_1)$	$M(j_0 j_1)$	$\Delta R(l_0)(j_0)$
$\Delta R(l_0)(j_1)$	$\Delta R(l_1)(j_0)$	$\Delta R(l_1)(j_1)$
$M(l_0 j_0)$	$M(l_0 j_1)$	$M(l_1 j_0)$
$M(l_1 j_1)$	jet DL1r 0	jet DL1r 1
nJets OR DL1r 85	nJets OR DL1r 60	$\Delta R(j_0 l_0)(j_1 l_1)$
$\Delta R(j_0 l_1)(j_1 l_0)$	$p_T(j_0 j_1 l_0 l_1 E_T^{\text{miss}})$	$M(j_0 j_1 l_0 l_1 E_T^{\text{miss}})$
$\Delta\phi(j_0)(E_T^{\text{miss}})$	$\Delta\phi(j_1)(E_T^{\text{miss}})$	HT jets
nJets	E_T^{miss}	

Table 3: Input features used in the 2ISS b-jet identification algorithm

As there are far more incorrect combinations than correct ones, by a factor of more than 20:1, the training set is resampled to reduce the fraction of incorrect combinations. A random sample of 5 million incorrect entries are used for training, along with close 1 million correct entries. 10% of the dataset is set aside for testing, leaving around 5 million datapoints for training.

The difference between the distributions for a few of these features for the correct (i.e. both jets are truth b-jets), and incorrect combinations are shown in figure 5.1. The correct and incorrect contributions are scaled to the same integral, so as to better demonstrate the differences in the distributions.

The modeling of these inputs is validated against data, with figure 5.2 showing good general agreement between data and MC. Plots for the complete list of features can be found in section A.

Based on the results of grid search evaluation, the optimal architecture is found to include 5 hidden layers with 40 nodes each. No regularizer or dropout is added to the network, as overfitting is found to not be an issue. The output score distribution as well as the ROC curve for the trained model are shown in figure 5.2.1. The model is found to identify the correct pairing of jets for 73% of 2ISS signal events on test data.

For point of comparison, a naïve approach to identify b-jets is used as well: The two jets which pass the highest DL1r b-tag working point are assumed to be the b-jets from the top decay. In the case that multiple jets meet the same b-tag working point, the jet with higher p_T is used. This method identifies the correct jet pair 65% of the time.

The accuracy of the model for different values of n-bjets, compared to this naive approach, is shown in table 4.

5.2.2 3l Channel

The input features used in the 3l channel are listed in table 5, with the same naming convention as the 2ISS channel.




Figure 5.1: Input features for top2ISS training. The signal in blue represents events where both jet candidates are truth b-jets from top decays, and the orange is all other combinations. Scaled to the same number of events.

Table 4: Accuracy of the NN in identifying b-jets from tops in 2ISS events for, compared to the accuracy of taking the two highest b-tagged jets.

b-jet Selection	Neural Network	Naive
1 b-jet	58.6%	42.1%
2 b-jets	88.4%	87.1%
≥ 3 b-jets	61.7%	53.3%
Overall	73.9%	67.2%

A few of these features are shown in figure 5.4, comparing the distributions for correct and incorrect combinations of jets.

The modeling of these inputs is validated against data, with figure 5.5 showing good general agreement between data and MC. Plots for the complete list of features can be found in section A.

Again, the dataset is downsized to reduce the ratio of correct and incorrect combination from 20:1, to 5:1. Around 7 million events are used for training, with 10% set aside for testing. Based on the results of grid search evaluation, the optimal architecture is found to include 5 hidden layers with 60 nodes each. The output score distribution as well as the ROC curve for the trained model are shown in figure 5.2.2.

This procedure is found to identify the correct pairing of jets for nearly 80% of 3l signal events. The accuracy of the model is summarized in table 6.




Figure 5.2: Data/MC comparisons of input features for top2ISS training for 80 fb^{-1} of data.

5.3 Higgs Reconstruction

Techniques similar to the b-jet identification algorithms are employed to select the decay products of the Higgs: kinematics of all possible combinations of reconstructed objects are fed into a neural network to determine which of those is most mostly to be the decay products of the Higgs.

Again separate models are used for the 2ISS and 3l channels, while the 3l channel has now been split into two: $t\bar{t}H$ events with three leptons in the final state include both instances where the Higgs decays into a lepton (and a neutrino) and a pair of jets, and instances where the Higgs decays to two leptons.

3l events are therefore categorized as either semi-leptonic or fully-leptonic. In the semi-leptonic case the reconstructed decay products consist of two jets and a single leptons. For the fully-leptonic case, the decay products include 2 of the three leptons associated with the event. For training the models, events are separated into these two categories using truth level information. A separate MVA, described in section 5.5, is used to make this distinction at reco level and determine which model to use.




Figure 5.3: (a) the output score of the NN for correct and incorrect combinations of jets. (b) the ROC curve of the output, showing background rejection as a function of signal efficiency

jet p_T 0	jet p_T 1	jet η 0
jet η 1	Lepton p_T 0	Lepton p_T 1
Lepton p_T 2	$\Delta R(j_0)(j_1)$	$M(j_0j_1)$
$\Delta R(l_0)(j_0)$	$\Delta R(l_1)(j_0)$	$\Delta R(l_2)(j_0)$
$\Delta R(l_0)(j_1)$	$\Delta R(l_1)(j_1)$	$\Delta R(l_2)(j_1)$
$M(l_0j_0)$	$M(l_1j_0)$	$M(l_2j_0)$
$M(l_0j_1)$	$M(l_1j_1)$	$M(l_2j_1)$
$\Delta R(j_0l_0)(j_1l_1)$	$\Delta R(j_0l_0)(j_1l_2)$	$\Delta R(j_0l_1)(j_1l_0)$
$\Delta R(j_0l_2)(j_1l_0)$	jet DL1r 0	jet DL1r 1
$p_T(j_0j_1l_0l_1l_2E_T^{\text{miss}})$	$M(tj_0j_1l_0l_1l_2E_T^{\text{miss}})$	$\Delta\phi(j_0)(E_T^{\text{miss}})$
$\Delta\phi(j_1)(E_T^{\text{miss}})$	HT Lepton	HT jets
nJets	E_T^{miss}	nJets OR DL1r 85
nJets OR DL1r 60		

Table 5: Input features

Table 6: Accuracy of the NN in identifying b-jets from tops, compared to the naive method of taking the highest b-tagged jets.

	NN	Naive
1 b-jet	69.0%	48.9%
2 b-jets	89.6%	88.3%
≥ 3 b-jets	55.7%	52.3%
Overall	79.8%	70.2%




Figure 5.4: Input features for top3l training. The signal in blue represents events where both jet candidates are truth b-jets from top decays, and the orange is all other combinations. Scaled to the same number of events.

For all channels, the models described in section 5.2 are used to identify b-jet candidates, whose kinematics are used to identify the Higgs decay products. These jets are not considered as possible candidates for the Higgs decay, justified by the fact that these models are found to misidentify jets from the Higgs decay as jets from the top decay less than 1% of the time.

5.3.1 2lSS Channel

For the 2lSS channel, the Higgs decay products include one light lepton and two jets. The neural network is trained on the kinematics of different combinations of leptons and jets, as well as the b-jets identified in section 5.2, with the specific input features listed in table ??.




Figure 5.5: Data/MC comparisons of input features for top31 training for 80 fb^{-1} of data.




Figure 5.6: tmp




Figure 5.7: Input features for higgs2lSS training. The signal in blue represents events where both jet candidates are truth b-jets from top decays, and the orange is all other combinations. Scaled to the same number of events.

260 The modeling of these inputs is validated against data, with figure 5.2 showing good general
 261 agreement between data and MC. Plots for the complete list of features can found in section A.

262 The neural network identifies the correct combination 55% of the time. It identifies the correct
 263 lepton 85% of the time, and selects the correct lepton and at least one of the correct jets 81% of
 264 the time.

265 5.3.2 3l Semi-leptonic Channel

266 For 3l $t\bar{t}H$ where the Higgs decay semi-leptonically, the decay products include one of the three
 267 leptons and two jets. In this case, the other two leptons originated from the decay of the tops,
 268 meaning the opposite-sign (OS) lepton cannot have come the Higgs. This leave only the two
 269 same-sign (SS) leptons as possible Higgs decay products.




Figure 5.8: Data/MC comparisons of input features for higgs2lSS training for 80 fb⁻¹ of data.




Figure 5.9: (a) the output score of the NN for correct and incorrect combinations of jets. (a) the ROC curve of the output, showing background rejection as a function of signal efficiency




Figure 5.10: Input features for higgs3lS training. The signal in blue represents events where both jet candidates are truth b-jets from top decays, and the orange is all other combinations. Scaled to the same number of events.

270 The modeling of these inputs is validated against data, with figure 5.11 showing good general
 271 agreement between data and MC. Plots for the complete list of features can found in section A.

272 The neural network identifies the correct combination 65% of the time. It identifies the correct
 273 lepton 85% of the time, anselects the correct lepton and at least one of the correct jets 83% of the
 274 time.

275 5.3.3 3l Fully-leptonic Channel

276 In the fully-leptonic 3l case, the goal is identify which two of the three leptons originated from
 277 the Higgs decay. Since one of these two must be the OS lepton, this problem is reduced to
 278 determining which of the two SS leptons originated from the Higgs. The kinematics of both
 279 possibilities are used for training, one where the SS lepton from the Higgs is correctly labeled,
 280 and one where it is not.




Figure 5.11: Data/MC comparisons of input features for higgs3lS training for 80 fb^{-1} of data.




Figure 5.12: (a) the output score of the NN for correct and incorrect combinations of jets. (a) the ROC curve of the output, showing background rejection as a function of signal efficiency




Figure 5.13: Input features for higgs3lF training. The signal in blue represents events where both jet candidates are truth b-jets from top decays, and the orange is all other combinations. Scaled to the same number of events.

281 The modeling of these inputs is validated against data, with figure 5.14 showing good general
 282 agreement between data and MC. Plots for the complete list of features can found in section A.

283 The correct lepton is identified 80% of the time.

284 5.4 p_T Prediction

285 Once the most probable decay products have been identified, their kinematics are used as inputs
 286 to a regression model which attempts to predict the momentum of the Higgs Boson. Once again,
 287 a DNN is used. Input variables representing the b-jets and leptons not from the Higgs decay
 288 are included as well, as these are found to improve performance. The truth p_T of the Higgs,
 289 as predicted by MC, are used as labels. Separate models are built for each channel - 2lSS, 3l
 290 Semi-leptonic and 3l Fully-leptonic.

291 As a two-bin fit is targeted for the final result, some metrics evaluating the performance of the
 292 models aim to show how well it distinguished between "high p_T " and "low p_T " events. A cutoff
 293 point of 150 GeV is used to define these two categories.

294 Because the analysis uses a two bin fit of the Higgs p_T , the momentum reconstruction could be
 295 treated as a binary classification problem, rather than a regression problem. This approach is
 296 explored in detail in section ??, and is found not to provide any significant increase in sensitivity.
 297 The regression approach is used because it provides more flexibility for future analyses, as it is




Figure 5.15: (a) the output score of the NN for correct and incorrect combinations of jets. (a) the ROC curve of the output, showing background rejection as a function of signal efficiency

307 The optimal neural network architecture for this channel is found to consist of 5 hidden layers with
 308 40 nodes each. The inputdata set includes 1.2 million events, 10% of which is used for testing,
 309 the other 90% for training. Training is found to converge after around 150 epochs.




Figure 5.16:

310 As the analysis strategy targets a two p_T bin fit, the problem ultimately reduces to a binary
 311 classification problem.




Figure 5.17:

312 5.4.2 3l Semi-leptonic Channel

313 The optimal neural network architecture for this channel is found to consist of 5 hidden layers
 314 with 40 nodes each. The inputdata set includes one million events, 10% of which is used for
 315 testing, the other 90% for training. Training is found to converge after around 150 epochs.




Figure 5.18:




Figure 5.19:

316 5.4.3 3l Fully-leptonic Channel

317 The optimal neural network architecture for this channel is found to consist of 5 hidden layers
 318 with 40 nodes each. The input data set includes 400,000 events, 10% of which is used for testing,
 319 the other 90% for training. Training is found to converge after around 150 epochs.

320 The predicted transverse momentum, as a function of the truth p_T , is shown in figure ??.




Figure 5.20:

321 When split into high and low p_T , based on a cutoff of 150 GeV, the




Figure 5.21:

322 5.5 3l Decay Mode

323 In the 3l channel, there are two possible ways for the Higgs to decay, both involving intermediate
 324 W boson pairs: Either both W bosons decay leptonically, in which case the reconstructed decay

325 consists of two leptons (referred as the fully-leptonic 3l channel), or one W decays leptonically
326 and the other hadronically, giving two jets and one lepton in the final state (referred to as the
327 semi-leptonic 3l channel). In order to accurately reconstruct the Higgs, it is necessary to identify
328 which of these decays took place for each 3l event.

329 The kinematics of each event, along with the output scores of the Higgs and top reconstruction
330 algorithms, are used to distinguish these two possible decay modes. The particular inputs used
331 are listed in table ??.

A neural network with 5 hidden layers, each with 50 nodes, is trained to distinguish these two decay modes. The output of the model is summarized in figure 5.22.




Figure 5.22:

6 Signal Region Definitions

Events are divided into two channels based on the number of leptons in the final state: one with two same-sign leptons, the other with three leptons. The 3l channel includes events where both leptons originated from the Higgs boson as well as events where only one of the leptons

6.1 Pre-MVA Event Selection

A preselection is applied to define orthogonal analysis channels based on the number of leptons in each event. For the 2lSS channel, the following preselection is used:

- Two very tight, same-charge, light leptons with $p_T > 20$ GeV
- ≥ 4 reconstructed jets, ≥ 1 b-tagged jets
- No reconstructed tau candidates

For the 3l channel, the following selection is applied:

- Three light leptons with total charge ± 1
- Same charge leptons are required to be very tight, with $p_T > 20$ GeV
- Opposite charge lepton must be loose, with $p_T > 10$ GeV




Figure 6.1:

- ≥ 2 reconstructed jets, ≥ 1 b-tagged jets
- No reconstructed tau candidates
- $|M(l^+l^-) - 91.2 \text{ GeV}| > 10 \text{ GeV}$ for all opposite-charge, same-flavor lepton pairs

6.2 Event MVA

Separate multi-variate analysis techniques (MVAs) are used in order to distinguish signal events from background for each analysis channel - 2lSS, 3l semi-leptonic, and 3l fully leptonic. In particular, Neural Networks produced with Tensorflow are trained using the kinematics of signal and background events derived from Monte Carlo simulations. Further, because the background composition differs for events with a high reconstructed Higgs p_T compared to events with low reconstructed Higgs p_T , separate MVAs are produced for high and low p_T regions.

Output distributions of each MVA are shown in figure 6.2. Detailed explanations of each of the models can be found in section A.




Figure 6.2:

6.3 Signal Region Definitions

Once pre-selection has been applied, channels are further refined based on the MVAs described above. The output of the model described in section 5.5 is used to separate the three channel into two - Semi-leptonic and Fully-leptonic - based on the predicted decay mode of the Higgs boson.

For each event, depending on the channel as well as the predicted p_T of the Higgs derived from the algorithm described in section 5.4, a cut on the appropriate background rejection algorithm is applied. The specific selection used, and the event yield in each channel after this selection has been applied, is summarized below.




Figure 6.3: scores

6.3.1 2LSS

6.3.2 3l — Semi — leptonic

6.3.3 3l — Fully — leptonic

7 Systematic Uncertainties

The systematic uncertainties that are considered are summarized in table ?? . These are implemented in the fit either as a normalization factors or as a shape variation or both in the signal and background estimations. The numerical impact of each of these uncertainties is outlined in section 8 .

The uncertainty in the combined 2015+2016 integrated luminosity is derived from a calibration of the luminosity scale using x-y beam-separation scans performed in August 2015 and May 2016

Table 7: Sources of systematic uncertainty considered in the analysis. Some of the systematic uncertainties are split into several components, as indicated by the number in the rightmost column.

Systematic uncertainty	Components
Luminosity	1
Pileup reweighting	1
Physics Objects	
Electron	6
Muon	15
Jet energy scale and resolution	28
Jet vertex fraction	1
Jet flavor tagging	131
E_T^{miss}	3
Total (Experimental)	186
Background Modeling	
Cross section	24
Renormalization and factorization scales	10
Parton shower and hadronization model	2
Shower tune	4
Total (Signal and background modeling)	40
Background Modeling	
Cross section	24
Renormalization and factorization scales	10
Parton shower and hadronization model	2
Shower tune	4
Total (Signal and background modeling)	40
Total (Overall)	226

[lumi].

The experimental uncertainties are related to the reconstruction and identification of light leptons and b-tagging of jets, and to the reconstruction of E_T^{miss} . The sources which contribute to the uncertainty in the jet energy scale [jes] are decomposed into uncorrelated components and treated as independent sources in the analysis.

The uncertainties in the b-tagging efficiencies measured in dedicated calibration analyses [btag_cal] are also decomposed into uncorrelated components. The large number of components for b-tagging is due to the calibration of the distribution of the BDT discriminant.

The systematic uncertainties associated with the signal and background processes are accounted for by varying the cross-section of each process within its uncertainty.

8 Results

A maximum likelihood fit is performed simultaneously over the regions described in section ??.

9 Conclusion

As search for the effects of dimension-six operators on $t\bar{t}H$ production is performed. An effective field theory approach is used to parameterize the effects of high energy physics on the Higgs momentum spectrum. The momentum spectrum is reconstructed using various MVA techniques, and the limits on dimension-six operators are limited to X.

396 **List of contributions**

397

Appendices

A Machine Learning Models

A.1 Background Rejection MVAs

Separate models are used in order to distinguish signal events from background for each analysis channel - 2ISS, 3l semi-leptonic, and 3l fully leptonic. In particular, Neural Networks produced with Tensorflow are trained using the kinematics of signal and background events derived from Monte Carlo simulations. Further, because the background composition differs for events with a high reconstructed Higgs p_T compared to events with low reconstructed Higgs p_T , separate MVAs are produced for high and low p_T regions.

A.1.1 2ISS - High p_T




Figure A.1:

408 **A.1.2 2lSS - Low p_T**

409 **A.1.3 3l Semi-Leptonic - High p_T**

410 **A.1.4 3l Semi-Leptonic - Low p_T**

411 **A.1.5 3l Fully Leptonic - High p_T**

412 **A.1.6 3l Fully Leptonic - Low p_T**