

Study of WZ + Heavy Flavor Production in the Fully Leptonic Channel



Aaron Webb
University of Texas at Austin



Abstract

A measurement of WZ produced with an associated heavy flavor jet is performed using 140 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13 \text{ TeV}$ from the ATLAS experiment at the LHC. The measurement is performed in the fully leptonic decay mode, $WZ \rightarrow l\nu ll$. Events are separated into inclusive 1-jet and 2-jet categories, and regions formed based on pseudo-continuous b-tag spectrum of the associated jets are fit to data.

Introduction




Figure 1: WZ + b and WZ + c production

- Many major analyses include WZ + b as a background, motivating a measurement of this process, which is difficult to simulate accurately
- The continuous b-tagging spectrum of the jets is used to separate out WZ + b from WZ + light, forming regions which are fit to data.
- Currently blinded to data, showing MC only Asimov fits




Figure 2: Distribution of DL1r score for b, charm, and light jets

Event Preselection

- Exactly three, tight, isolated leptons with $p_T > 20 \text{ GeV}$, $|\eta| < 2.47$
- M_{l+l-} of a pair of oppositely charged, same flavor leptons within 10 GeV of 91.2 GeV
- Require 1-2 jets, with $p_T > 25 \text{ GeV}$, $|\eta| < 2.5$
- $E_{miss}^T > 20 \text{ GeV}$
- Data/MC Yields after preselection has been applied are shown in figure 1

Fit Procedure

- MC predictions are fit to the full Run-2 dataset, 140 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ data
- Truth jets are binned based on the b-jet efficiency working points of the DL1r algorithm - 85%, 77%, 70% and 60%
- Events meeting the highest working point are further separated into a signal like region and a tZ CR based on an MVA
- The WZ events are separated by truth flavor into three templates, which are fit to data




Figure 3: Distribution of (left) the p_T of the opposite sign lepton, (center) the invariant mass of a lepton pair, and (right) the p_T of the leading jet.

tZ BDT

- Primary background in the high b-tag region is tZ, introduces large uncertainty
- Lepton, jet kinematics, and reconstructed top mass are used as inputs to a BDT to distinguish WZ from tZ events
- Output score is used to form a tZ CR




Figure 4: Output of the BDT for WZ and tZ events

Results




Figure 5: Summary of the fit regions for (left) 1-jet events and (right) 2-jet events.

1-jet Asimov Fit Results:

$$\mu_{WZ+b} = 1.00^{+0.54}_{-0.49}$$

$$\mu_{WZ+c} = 1.00 \pm 0.21$$

$$\mu_{WZ+l} = 1.00 \pm 0.06$$

- The impact of the systematic uncertainties considered in the fit on the measured value of WZ+b is shown in figure 1




Figure 6: Ranking of the impact of nuisance parameters on (left) the 1-jet fit and (right) the 2-jet fit

Conclusions

- Blinded fits show a measurement of WZ + heavy flavor can be made
- Nearly ready to unblind, present final results