

Lecture 14

Optimization

Ashley Gannon

ISC3313 Fall 2021



What is optimization?



What is optimization?

- The determination of a function's maximum or minimum (or optimal) value is referred to as *optimization*.




What is optimization?

- The determination of a function's maximum or minimum (or optimal) values is referred to as *optimization*.
- As you learned in calculus, such solutions can be obtained analytically by determining the value at which the function is flat; $f'(x) = 0$.



What is optimization?

- The determination of a function's maximum or minimum (or optimal) value is referred to as *optimization*.
- As you learned in calculus, such solutions can be obtained analytically by determining the value at which the function is flat; $f'(x) = 0$.
- The fundamental difference is that finding the location of a root involves searching for x where the $f(x) = 0$ while optimization involves searching for x where $f'(x) = 0$.



Optimization Example

Let's consider the following problem:

An object like a bungee jumper can be projected upward at a specified velocity. If it is subject to linear drag, its altitude as a function of time can be computed as

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

Where z is the distance from the Earth's surface, z_0 is the initial distance from the earth's surface, v_0 is the initial velocity, m is the mass of the jumper, c_d is the drag coefficient, and t is time.




Optimization Example

Let's consider the following problem:

An object like a bungee jumper can be projected upward at a specified velocity. If it is subject to linear drag, its altitude as a function of time can be computed as

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

If we let $z_0 = 100\text{m}$, $m = 80\text{kg}$, $c_d = 15\text{kg/s}$, $v_0 = 55\text{m/s}$, and $g = 9.81\text{m/s}^2$, and plot z for $t = 0 \dots 12$



Optimization Example

Let's consider the following problem:

An object like a bungee jumper can be projected upward at a specified velocity. If it is subject to linear drag, its altitude as a function of time can be computed as

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

To find the maximum altitude analytically, we have to find the time where $z'(t) = 0$.

$$\frac{dz}{dt} = v_0 e^{\frac{-c_d t}{m}} - \frac{mg}{c_d} \left(1 - e^{\frac{-c_d t}{m}} \right) = 0$$



Optimization Example

Let's consider the following problem:

An object like a bungee jumper can be projected upward at a specified velocity. If it is subject to linear drag, its altitude as a function of time can be computed as

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

To find the maximum altitude analytically, we have to find the time where $z'(t) = 0$.

$$\frac{dz}{dt} = v_0 e^{\frac{-c_d t}{m}} - \frac{mg}{c_d} \left(1 - e^{\frac{-c_d t}{m}} \right) = 0$$

Solving analytically for t we get

$$t = \frac{m}{c_d} \ln \left(1 + \frac{c_d v_0}{mg} \right)$$



Optimization Example

Let's consider the following problem:

An object like a bungee jumper can be projected upward at a specified velocity. If it is subject to linear drag, its altitude as a function of time can be computed as

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

To find the maximum altitude analytically, we have to find the time where $z'(t) = 0$.

$$\frac{dz}{dt} = v_0 e^{\frac{-c_d t}{m}} - \frac{mg}{c_d} \left(1 - e^{\frac{-c_d t}{m}} \right) = 0$$

Solving analytically for t we get

$$t = \frac{m}{c_d} \ln \left(1 + \frac{c_d v_0}{mg} \right)$$

Again, letting $z_0 = 100\text{m}$, $m = 80\text{kg}$, $c_d = 15\text{kg/s}$, $v_0 = 55\text{m/s}$, and $g = 9.81\text{m/s}^2$

$$t \approx 3.83166\text{s}$$



Optimization Example

Now to find the maximum position, we plug this value for t and our other parameters back into

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

to find that

$$z \approx 192.8609\text{m}$$



Optimization Example

Now to find the maximum position, we plug this value for t and our other parameters back into

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

to find that

$$z \approx 192.8609\text{m}$$

Now if we want to verify that this is a maximum value, we need to evaluate $z''(x)$

$$\frac{dz^2}{d^2z} = -\frac{c_d v_0}{m} e^{\frac{-c_d t}{m}} - g e^{\frac{-c_d t}{m}}$$



Optimization Example

Now to find the maximum position, we plug this value for t and our other parameters back into

$$z = z_0 + \frac{m}{c_d} \left(v_0 + \frac{mg}{c_d} \right) \left(1 - e^{\frac{-c_d t}{m}} \right) - \frac{mg}{c_d} t$$

to find that

$$z \approx 192.8609\text{m}$$

Now if we want to verify that this is a maximum value, we need to evaluate $z''(x)$

$$\frac{dz^2}{d^2z} = -\frac{c_d v_0}{m} e^{\frac{-c_d t}{m}} - g e^{\frac{-c_d t}{m}}$$

plugging in our variable values,


$$\frac{dz^2}{d^2z} \approx -9.81\text{m/s}$$

The fact that the second derivative is negative tells us that we have a maximum.



What is optimization?

- The determination of a function's maximum or minimum (or optimal) value is referred to as *optimization*.
- As you learned in calculus, such solutions can be obtained analytically by determining the value at which the function is flat; $f'(x) = 0$.
- The fundamental difference is that finding the location of a root involves searching for x where the $f(x) = 0$ while optimization involves searching for x where $f'(x) = 0$.




- Although such analytical solutions are sometimes feasible, most practical optimization problems require numerical, computer solutions.



What is optimization?

- The determination of a function's maximum or minimum (or optimal) value is referred to as *optimization*.
- As you learned in calculus, such solutions can be obtained analytically by determining the value at which the tangent of the function is flat; $f'(x) = 0$.
- The fundamental difference is that finding the location of a root involves searching for x where the $f(x) = 0$ while optimization involves searching for x where $f'(x) = 0$.



- Although such analytical solutions are sometimes feasible, most practical optimization problems require numerical, computer solutions.
- From a numerical standpoint, such optimization methods are similar in spirit to the root-location methods we discussed in the last section.
 - both involve guessing and searching for a location on a continuous function.



One Dimensional Optimization




1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.




1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.
- Cases that include local optima are called *multimodal*.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.
- Cases that include local optima are called *multimodal*.
- In such cases, we will almost always be interested in finding the global optimum.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.
- Cases that include local optima are called *multimodal*.
- In such cases, we will almost always be interested in finding the global optimum.
- Therefore, we must be concerned about mistaking a local result for the global optimum.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.




- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.
- Cases that include local optima are called *multimodal*.
- In such cases, we will almost always be interested in finding the global optimum.
- Therefore, we must be concerned about mistaking a local result for the global optimum.
- We will cover two methods for optimization - The *golden section-search method* and the *parabolic method*.



1D Optimization

- In this unit we will cover techniques that find the minimum or maximum of a function of a single variable, i.e. $f(x)$.
- Just as finding the root location was complicated by the fact that several roots can occur for a single function, both local and global optima can occur in optimization.



- A *global* optimum represents the very best solution.
- A *local* optimum, though not the very best, is better than its immediate neighbors.
- Cases that include local optima are called *multimodal*.
- In such cases, we will almost always be interested in finding the global optimum.
- Therefore, we must be concerned about mistaking a local result for the global optimum.
- We will cover two methods for optimization - The *golden section-search method* and the *parabolic method*.
- Both of these methods can be considered *Bracketing methods*.



The Golden-Section Search Method



The golden-section search method

- Like with Bisection, we need to define the interval that contains the optima.



The golden-section search method

- Like with Bisection, we need to define the interval that contains the optima.
- We can use the same nomenclature we did with the Bisection method, letting x_l be the lower bound and x_u being the upper bound.



The golden-section search method

- Like with Bisection, we need to define the interval that contains the optima.
- We can use the same nomenclature we did with the Bisection method, letting x_l be the lower bound and x_u being the upper bound.
- We cannot use a single intermediate value, x_r , like we did for Bisection, instead we need two intermediate values to detect whether a optima occurred.



The golden-section search method

- Like with Bisection, we need to define the interval that contains the optima.
- We can use the same nomenclature we did with the Bisection method, letting x_l be the lower bound and x_u being the upper bound.
- We cannot use a single intermediate value, x_r , like we did for Bisection, instead we need two intermediate values to detect whether an optima occurred.
 - The key to making this approach efficient is the wise choice of the intermediate points.
 - For the golden-section search, the two intermediate points are chosen according to the golden ratio:


$$x_1 = x_l + d$$

$$x_2 = x_u - d$$

Where

$$d = (\Phi - 1)(x_u - x_l)$$

and the golden ratio
 $\Phi = 1.61803398874989$.



The golden-section search method

- For the golden-section search, the two intermediate points are chosen according to the golden ratio:

$$x_1 = x_l + d$$


$$x_2 = x_u - d$$

Where

$$d = (\Phi - 1)(x_u - x_l)$$

and the golden ratio $\Phi = 1.61803398874989$.

- The function is evaluated at these two interior points. Two results can occur:
 - If $f(x_1) < f(x_2)$, then $f(x_1)$ is the optima, and the domain of x to the left of x_2 , from x_l to x_2 , can be eliminated because it does not contain the optima. For this case, x_2 becomes the new x_l for the next round.
 - If $f(x_2) < f(x_1)$, then $f(x_2)$ is the optima and the domain of x to the right of x_1 , from x_1 to x_u would be eliminated. For this case, x_1 becomes the new x_u for the next round.



Example

Let's find the minimum of

$$f(x) = \frac{x^2}{10} - 2\sin(x)$$


With and initial $x_l = 0$ and $x_u = 4$.

The first thing we need to do is calculate the golden ratio that we use to create the two interior points:

$$\begin{aligned} d &= (1.61803 - 1)(4 - 0) \\ &= 2.4721 \end{aligned}$$

Then our two interior points are

$$\begin{aligned} x_1 &= x_l + d = 0 + 2.4721 \\ &= 2.4721 \\ x_2 &= x_u - d \\ &= 4 - 2.4721 \\ &= 1.5279 \end{aligned}$$



Example

Now we need to compute our function values and compare them to find our new domain

$$\begin{aligned} f(x_1) &= \frac{2.4721^2}{10} - 2 \sin(2.4721) \\ &= -0.6300 \end{aligned}$$


$$\begin{aligned} f(x_2) &= \frac{1.5279^2}{10} - 2 \sin(1.5279) \\ &= -1.7647 \end{aligned}$$

Now since $f(x_2) < f(x_1)$,

$$x_l = x_l$$

$$x_u \leftarrow x_1$$

$$x_1 \leftarrow x_2$$




Example

We now need to recalculate d

$$\begin{aligned} d &= 0.61803(x_u - x_l) \\ &= 0.61803(2.4721 - 0) \\ &= 1.5279 \end{aligned}$$

And define a new x_2

$$\begin{aligned} x_2 &= x_u - d \\ &= 2.4721 - 1.5279 \\ &= 0.9443 \end{aligned}$$



Example

Next iteration: We need to compute our function values and compare them to find our new domain

$$\begin{aligned} f(x_1) &= \frac{1.5279^2}{10} - 2 \sin(1.5279) \\ &= -1.7647 \end{aligned}$$


$$\begin{aligned} f(x_2) &= \frac{0.9443^2}{10} - 2 \sin(0.9443) \\ &= -1.5310 \end{aligned}$$

Now since $f(x_1) < f(x_2)$,

$$x_u = x_u$$

$$x_l \leftarrow x_2$$

$$x_2 \leftarrow x_1$$




Example

We now need to recalculate d

$$\begin{aligned} d &= 0.61803(x_u - x_l) \\ &= 0.61803(2.4721 - 0.9443) \\ &= 0.9456 \end{aligned}$$

And define a new x_2

$$\begin{aligned} x_1 &= x_l + d \\ &= 0.9443 + 0.9456 \\ &= 1.8899 \end{aligned}$$



Keep iterating until

$$\epsilon_a = (2 - \Phi) \left| \frac{x_u - x_l}{x_{opt}} \right| < tol$$

Where x_{opt} is the new value of x_1 or x_2 .



Sign-Off Activity

Now that you have an idea how this method works, write a pseudo code and post it to the **Golden-section search method** discussion board. You may sign-off once you've finished.

