

Integration

Richardson Extrapolation and Romberg Integration

Ashley Gannon

Fall 2020



Richardson Extrapolation



Richardson Extrapolation

Richardson extrapolation methods improve the results of numerical integration by using two estimates of an integral to compute a third, more accurate approximation.



Richardson Extrapolation

Richardson extrapolation methods improve the results of numerical integration by using two estimates of an integral to compute a third, more accurate approximation.

We'll do examples involving successive iterates of the Trapezoidal Rule.



Richardson Extrapolation

Richardson extrapolation methods improve the results of numerical integration by using two estimates of an integral to compute a third, more accurate approximation.

We'll do examples involving successive iterates of the Trapezoidal Rule.

We can represent the error of the Trapezoidal rule by

$$I = I(h) + E(h)$$

where I is the exact integral, $I(h)$ is the integral approximation using the Trapezoidal rule, and $E(h)$ is the truncation error

$$E \cong -\frac{b-a}{12} h^2 \bar{f}''$$

when $h = (b-a)/n$.



Richardson Extrapolation

If we make two separate estimates using segment sizes h_1 and h_2 , we can rewrite

$$I = I(h) + E(h)$$

as

$$I(h_1) + E(h_1) = I(h_2) + E(h_2)$$



Richardson Extrapolation

If we make two separate estimates using segment sizes h_1 and h_2 , we can rewrite

$$I = I(h) + E(h)$$

as

$$I(h_1) + E(h_1) = I(h_2) + E(h_2)$$

If we assume that \bar{f}'' is constant for any h , we have the relation

$$\frac{E(h_1)}{E(h_2)} \cong \frac{-\frac{b-a}{12} h_1^2 \bar{f}''}{-\frac{b-a}{12} h_2^2 \bar{f}''} \cong \frac{h_1^2}{h_2^2}$$



Richardson Extrapolation

If we make two separate estimates using segment sizes h_1 and h_2 , we can rewrite

$$I = I(h) + E(h)$$

as

$$I(h_1) + E(h_1) = I(h_2) + E(h_2)$$

If we assume that \bar{f}'' is constant for any h , we have the relation

$$\frac{E(h_1)}{E(h_2)} \cong \frac{-\frac{b-a}{12} h_1^2 \bar{f}''}{-\frac{b-a}{12} h_2^2 \bar{f}''} \cong \frac{h_1^2}{h_2^2}$$

Solving for $E(h_1)$,

$$E(h_1) = E(h_2) \frac{h_1^2}{h_2^2}$$



Richardson Extrapolation

If we make two separate estimates using segment sizes h_1 and h_2 , we can rewrite

$$I = I(h) + E(h)$$

as

$$I(h_1) + E(h_1) = I(h_2) + E(h_2)$$

If we assume that \bar{f}'' is constant for any h , we have the relation

$$\frac{E(h_1)}{E(h_2)} \cong \frac{-\frac{b-a}{12} h_1^2 \bar{f}''}{-\frac{b-a}{12} h_2^2 \bar{f}''} \cong \frac{h_1^2}{h_2^2}$$

Solving for $E(h_1)$,

$$E(h_1) = E(h_2) \frac{h_1^2}{h_2^2}$$

Which we can plug back into the above equation so that the integral approximate is only dependent on $E(h_2)$.

$$I(h_1) + E(h_2) \frac{h_1^2}{h_2^2} = I(h_2) + E(h_2)$$



Richardson Extrapolation

If we make two separate estimates using segment sizes h_1 and h_2 , we can rewrite

$$I = I(h) + E(h)$$

as

$$I(h_1) + E(h_1) = I(h_2) + E(h_2)$$

If we assume that \tilde{f}'' is constant for any h , we have the relation

$$\frac{E(h_1)}{E(h_2)} \cong \frac{-\frac{b-a}{12} h_1^2 \tilde{f}''}{-\frac{b-a}{12} h_2^2 \tilde{f}''} \cong \frac{h_1^2}{h_2^2}$$

Solving for $E(h_1)$,

$$E(h_1) = E(h_2) \frac{h_1^2}{h_2^2}$$

Which we can plug back into the above equation so that the integral approximate is only dependent on $E(h_2)$.

$$I(h_1) + E(h_2) \frac{h_1^2}{h_2^2} = I(h_2) + E(h_2)$$

Solving for $E(h_2)$ we have

$$E(h_2) = \frac{I(h_1) - I(h_2)}{1 - \frac{h_1^2}{h_2^2}}$$



Richardson Extrapolation

Since we are interested in the best approximate for the integral, we will use

$$I = I(h_2) + E(h_2)$$



Richardson Extrapolation

Since we are interested in the best approximate for the integral, we will use

$$I = I(h_2) + E(h_2)$$

We can substitute the error we found in the previous slide into this equation to yield

$$I = I(h_2) + \frac{I(h_1) - I(h_2)}{1 - \frac{h_1^2}{h_2^2}}$$



Richardson Extrapolation

Since we are interested in the best approximate for the integral, we will use

$$I = I(h_2) + E(h_2)$$

We can substitute the error we found in the previous slide into this equation to yield

$$I = I(h_2) + \frac{I(h_1) - I(h_2)}{1 - \frac{h_1^2}{h_2^2}}$$

The error of this approximate is now $\mathcal{O}(h^4)$, compared to each iteration of the Trapezoidal rule $\mathcal{O}(h^2)$. (Raltson and Rabinowitz, 1978)



Richardson Extrapolation

Since we are interested in the best approximate for the integral, we will use

$$I = I(h_2) + E(h_2)$$

We can substitute the error we found in the previous slide into this equation to yield

$$I = I(h_2) + \frac{I(h_1) - I(h_2)}{1 - \frac{h_1^2}{h_2^2}}$$

The error of this approximate is now $\mathcal{O}(h^4)$, compared to each iteration of the Trapezoidal rule $\mathcal{O}(h^2)$. (Ralston and Rabinowitz, 1978)

For the special case where the interval is halved, $h_2 = h_1/2 = \frac{b-a}{2n}$, we can simplify our approximate to be

$$\begin{aligned} I &= I(h_2) + \frac{I(h_1) - I(h_2)}{1 - \frac{h_1^2}{(h_1/2)^2}} = I(h_2) + \frac{I(h_1) - I(h_2)}{1 - 4} = I(h_2) + \frac{I(h_2)}{3} - \frac{I(h_1)}{3} \\ &= \frac{4}{3}I(h_2) - \frac{1}{3}I(h_1) \end{aligned}$$



Richardson Extrapolation Example

Use Richardson Extrapolation to evaluate the integral of

$f(x) = 0.2 + 25x - 200x^2 + 675x^3 - 900x^4 + 400x^5$ from $a = 0$ to $b = 0.8$.



Richardson Extrapolation Example

Use Richardson Extrapolation to evaluate the integral of
 $f(x) = 0.2 + 25x - 200x^2 + 675x^3 - 900x^4 + 400x^5$ from $a = 0$ to $b = 0.8$.

First we will use the trapezoid rule on one segment

$$h = (0.8 - 0)/1 = 0.8$$

$$f(0.8) = 0.232, \quad f(0) = 0.2$$

$$\begin{aligned} I &\approx h \frac{f(a) + f(b)}{2} = 0.8 * \frac{0.232 + 0.2}{2} \\ &= 0.1728 \end{aligned}$$



Richardson Extrapolation Example

Use Richardson Extrapolation to evaluate the integral of
 $f(x) = 0.2 + 25x - 200x^2 + 675x^3 - 900x^4 + 400x^5$ from $a = 0$ to $b = 0.8$.

First we will use the trapezoid rule on one segment

$$h = (0.8 - 0)/1 = 0.8$$

$$f(0.8) = 0.232, \quad f(0) = 0.2$$

$$\begin{aligned} I &\approx h \frac{f(a) + f(b)}{2} = 0.8 * \frac{0.232 + 0.2}{2} \\ &= 0.1728 \end{aligned}$$

Next we will halve the interval $h = (0.8 - 0)/2 = 0.4$, and apply the Trapezoidal rule

$$\begin{aligned} I &\approx h \frac{f(x_0) + f(x_1)}{2} + h \frac{f(x_1) + f(x_2)}{2} \\ &= 0.4 \frac{0.2 + 2.456}{2} + 0.4 \frac{2.456 + f(0.232)}{2} \\ &= 1.0688 \end{aligned}$$



Richardson Extrapolation Example

Use Richardson Extrapolation to evaluate the integral of $f(x) = 0.2 + 25x - 200x^2 + 675x^3 - 900x^4 + 400x^5$ from $a = 0$ to $b = 0.8$.

Now we can plug these values into our integral approximate

$$\begin{aligned} I &= \frac{4}{3}I(h_2) - \frac{1}{3}I(h_1) \\ &= \frac{4}{3}1.0688 - \frac{1}{3}0.1728 \\ &\approx 1.367467 \end{aligned}$$

Which is closer to the true value, 1.640533 than either Trapezoidal rule approximate.



Richardson Extrapolation Example

In the same manner, we can use estimates for 2 and 4 segments.

For 4 segments

$$h = (0.8 - 0)/4 = 0.2$$

$$x_0 = 0, \quad x_1 = 0.2, \quad x_2 = 0.4,$$

$$x_3 = 0.6, \quad x_4 = 0.8$$

Plugging these values into our trapezoid rule,

$$I \approx \frac{h}{2}(f(x_0) + f(x_4)) + h \sum_{i=2}^4 f(x_i) = 1.4848$$



Richardson Extrapolation Example

In the same manner, we can use estimates for 2 and 4 segments.

For 4 segments

$$h = (0.8 - 0)/4 = 0.2$$

$$x_0 = 0, \quad x_1 = 0.2, \quad x_2 = 0.4,$$

$$x_3 = 0.6, \quad x_4 = 0.8$$

Plugging these values into our trapezoid rule,

$$I \approx \frac{h}{2}(f(x_0) + f(x_4)) + h \sum_{i=2}^4 f(x_i) = 1.4848$$

Substituting this and the two segment approximate into

$$\begin{aligned} I &= \frac{4}{3}1.4848 - \frac{1}{3}1.0688 \\ &\approx 1.623467 \end{aligned}$$

Which is even more accurate than the last approximation.



Richardson Extrapolation Example

These two improved integrals can, in turn, be combined to yield an even better value with $\mathcal{O}(h^6)$. For the special case where the original trapezoidal estimates are based on successive halving of the step size, the equation used for $\mathcal{O}(h^6)$ accuracy is

$$I = \frac{16}{15} I_m - \frac{1}{15} I_l$$

where the I_m is the more accurate estimate, I_l is the less accurate estimate.



Richardson Extrapolation Example

These two improved integrals can, in turn, be combined to yield an even better value with $\mathcal{O}(h^6)$. For the special case where the original trapezoidal estimates are based on successive halving of the step size, the equation used for $\mathcal{O}(h^6)$ accuracy is

$$I = \frac{16}{15} I_m - \frac{1}{15} I_l$$

where the I_m is the more accurate estimate, I_l is the less accurate estimate.

For our example, we have

$$\begin{aligned} I &= \frac{16}{15} (1.623467) - \frac{1}{15} (1.367467) \\ &= 1.640533 \end{aligned}$$

Which is the exact integral.



Romberg Integration



Romberg Integration

Notice the coefficients for

$$I = \frac{4}{3} I_m - \frac{1}{3} I_l$$

and

$$I = \frac{16}{15} I_m - \frac{1}{15} I_l$$

add to 1.

You may think of these coefficients as weights. We place a larger weight on the more accurate integral approximation.



Romberg Integration

Notice the coefficients for

$$I = \frac{4}{3}I_m - \frac{1}{3}I_l$$

and

$$I = \frac{16}{15}I_m - \frac{1}{15}I_l$$

add to 1.

You may think of these coefficients as weights. We place a larger weight on the more accurate integral approximation.

If we let $k = 1, 2, \dots, n$ correspond to the order of accuracy where

k	L
1	$\mathcal{O}(h^2)$
2	$\mathcal{O}(h^4)$
3	$\mathcal{O}(h^6)$
\vdots	\vdots
n	$\mathcal{O}(h^{2n})$



Romberg Integration

Notice the coefficients for

$$I = \frac{4}{3} I_m - \frac{1}{3} I_l$$

and

$$I = \frac{16}{15} I_m - \frac{1}{15} I_l$$

add to 1.

You may think of these coefficients as weights. We place a larger weight on the more accurate integral approximation.

If we let $k = 1, 2, \dots, n$ correspond to the order of accuracy where

k	L
1	$\mathcal{O}(h^2)$
2	$\mathcal{O}(h^4)$
3	$\mathcal{O}(h^6)$
\vdots	\vdots
n	$\mathcal{O}(h^{2n})$

We can generalize the form of our integral approximate to be

$$I_k = \frac{4^{k-1} I_{m,k-1} - I_{l,k-1}}{4^{k-1} - 1}$$



Romberg Integration

Notice the coefficients for

$$I = \frac{4}{3} I_m - \frac{1}{3} I_l$$

and

$$I = \frac{16}{15} I_m - \frac{1}{15} I_l$$

add to 1.

You may think of these coefficients as weights. We place a larger weight on the more accurate integral approximation.

If we let $k = 1, 2, \dots, n$ correspond to the order of accuracy where

k	L
1	$\mathcal{O}(h^2)$
2	$\mathcal{O}(h^4)$
3	$\mathcal{O}(h^6)$
\vdots	\vdots
n	$\mathcal{O}(h^{2n})$

We can generalize the form of our integral approximate to be

$$I_k = \frac{4^{k-1} I_{m,k-1} - I_{l,k-1}}{4^{k-1} - 1}$$

Applying this function iteratively is known as *Romberg Integration*. This method has the error

$$E_a = \left| \frac{I_k - I_{m,k-1}}{I_k} \right|$$




Romberg Integration

We can generalize the form of our integral approximate to be

$$I_k = \frac{4^{k-1} I_{m,k-1} - I_{l,k-1}}{4^{k-1} - 1}$$

Applying this function iteratively is known as *Romberg Integration*

Graphical depiction of the sequence of integral estimates generated using Romberg integration. (a) First iteration, (b) second iteration, (c) third iteration.



$O(h^4)$ Romberg Method Pseudocode

Download the `TrapezoidalRule.cpp` file from Canvas. We are going to modify it to contain the Romberg Method pseudocode.

```
define function that takes in a, b, and tol
declare/define error
declare/define n = 1
declare/define I1 = TrapezoidalRule(a,b,n)
declare/define Ik = 0, Im = 0
declare/define k = 2

loop while error is greater than the tolerance
    update n for next trapezoid rule call, n*=2
    Find Im by calling Trapezoidal rule for new n,
    Im = TrapezoidalRule(a,b,n)
    Compute Ik = (pow(4,k-1)*Im-I1)/(pow(4,k-1)-1)
    Compute the error = abs((Ik-Im)/Ik)
    Update I1 = Im
```

