

COMS21103: Integer Programming

Dima Damen

`Dima.Damen@bristol.ac.uk`

Bristol University, Department of Computer Science
Bristol BS8 1UB, UK

December 9, 2015

Integer Program

- ▶ An integer program is encountered when the optimum output is expected as integer values
- ▶ The book shipment example from the last lecture is an integer program
- ▶ This is because the output is not meaningful unless it is in integers
- ▶ In the book shipment example, even though the solution was actually given as integers (200,0,200,600), solving a linear program cannot guarantee that the solution is in integers

Integer Program

- ▶ An integer program is encountered when the optimum output is expected as integer values
- ▶ The book shipment example from the last lecture is an integer program
- ▶ This is because the output is not meaningful unless it is in integers
- ▶ In the book shipment example, even though the solution was actually given as integers (200,0,200,600), solving a linear program cannot guarantee that the solution is in integers
- ▶ In formulating this as a linear program, we ignored the 'integer' constraint
- ▶ This is referred to as **linear program relaxation of an integer program**

Integer Program

- ▶ An integer program is encountered when the optimum output is expected as integer values
- ▶ The book shipment example from the last lecture is an integer program
- ▶ This is because the output is not meaningful unless it is in integers
- ▶ In the book shipment example, even though the solution was actually given as integers (200,0,200,600), solving a linear program cannot guarantee that the solution is in integers
- ▶ In formulating this as a linear program, we ignored the 'integer' constraint
- ▶ This is referred to as **linear program relaxation of an integer program**
- ▶ One way around this is to round numbers... this can cause serious problems

Integer Program

An integer program thus has the standard form:

$$\begin{array}{ll} \text{maximise} & \sum_{j=1}^n c_j x_j \\ \text{subject to} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad \text{for } i = 1, 2, \dots, m \\ & x_j \geq 0 \quad \text{for } j = 1, 2, \dots, n \\ & x_j \in \mathbb{Z}(\text{integer}) \quad \text{for } j = 1, 2, \dots, n \end{array}$$

Integer Program

- ▶ An integer linear program is a linear program further constrained by the integrality restrictions
- ▶ In integer programs all variables are restricted to be integers

Integer Program

- ▶ An integer linear program is a linear program further constrained by the integrality restrictions
- ▶ In integer programs all variables are restricted to be integers
- ▶ In mixed-integer programs, some variables only are restricted to be integers
- ▶ Mixed-integer programs are out of the scope in these slides

Integer Program

- ▶ An integer linear program is a linear program further constrained by the integrality restrictions
- ▶ In integer programs all variables are restricted to be integers
- ▶ In mixed-integer programs, some variables only are restricted to be integers
- ▶ Mixed-integer programs are out of the scope in these slides
- ▶ Binary programs are programs where variables are restricted to the binary $\in \{0, 1\}$.
- ▶ These are special cases of integer programs and could be solved in the same way by adding inequalities $x_j \leq 1$ for $j = 1, 2, \dots, n$

Integer Program

- ▶ An integer linear program is a linear program further constrained by the integrality restrictions
- ▶ In integer programs all variables are restricted to be integers
- ▶ In mixed-integer programs, some variables only are restricted to be integers
- ▶ Mixed-integer programs are out of the scope in these slides
- ▶ Binary programs are programs where variables are restricted to the binary $\in \{0, 1\}$.
- ▶ These are special cases of integer programs and could be solved in the same way by adding inequalities $x_j \leq 1$ for $j = 1, 2, \dots, n$
- ▶ Integer programming has applications in planning, scheduling and networking

Integer Program - Case Study

Work of Raymond Kwan, University of Leeds
Scheduling crew shifts for train companies




Fig.1 Train work and crew shifts

R. Kwan (2009). Case Studies of Successful Train Crew Scheduling Optimization. Multidisciplinary International Conference on Scheduling: Theory and Applications.

Integer Program - Case Study

Work of Raymond Kwan, University of Leeds Scheduling crew shifts for train companies

$$\begin{aligned} &\text{Minimise} && W_1 \sum_{j=1}^n c_j x_j + W_2 \sum_{j=1}^n x_j \\ &\text{Subject to} && \end{aligned}$$

$$\sum_{j=1}^n a_{ij} x_j \geq 1 \quad i = 1, 2, \dots, m$$

$$x_j = 0 \text{ or } 1 \quad j = 1, 2, \dots, n$$

where

n is the number of candidate shifts

m is the number of work pieces

x_j is a shift variable, $x_j = \begin{cases} 1 & \text{if shift } j \text{ is selected} \\ 0 & \text{otherwise} \end{cases}$

c_j is the cost of shift j

$a_{ij} = \begin{cases} 1 & \text{if work piece } i \text{ is covered by shift } j \\ 0 & \text{otherwise} \end{cases}$

W_1 and W_2 are weight constants





Fig.1 Train work and crew shifts

Fig.2 Set covering ILP for train crew scheduling

R. Kwan (2009). Case Studies of Successful Train Crew Scheduling Optimization. Multidisciplinary International Conference on Scheduling: Theory and Applications.


Integer Programming and Linear Programming

- ▶ P_I : the space of feasible solutions of the integer program
- ▶ P : the space of feasible solutions of the *relaxed* linear program
- ▶ $P_I \subseteq P$



Integer Programming and Linear Programming

- ▶ P_I : the space of feasible solutions of the integer program
- ▶ P : the space of feasible solutions of the *relaxed* linear program
- ▶ $P_I \subseteq P$
- ▶ If rounding is used, the integer solution could be infeasible.



Integer Programming and Linear Programming

- Whereas the simplex method is effective for solving linear programs, there is no single technique for solving integer programs.

Integer Programming and Linear Programming

- ▶ Whereas the simplex method is effective for solving linear programs, there is no single technique for solving integer programs.
- ▶ We thus first relax the integer program by removing integer constraints on the variables
- ▶ The *relaxed* linear program is then solved using the simplex method

Integer Programming and Linear Programming

- ▶ Whereas the simplex method is effective for solving linear programs, there is no single technique for solving integer programs.
- ▶ We thus first relax the integer program by removing integer constraints on the variables
- ▶ The *relaxed* linear program is then solved using the simplex method
- ▶ The integer solution is then found using one of three strategies:
 - ▶ Enumeration techniques (e.g. Branch-and-bound)
 - ▶ Cutting plane techniques
 - ▶ Group-theoretic techniques
- ▶ We will discuss the first two strategies next,

Integer Programming and Linear Programming


- ▶ Whereas the simplex method is effective for solving linear programs, there is no single technique for solving integer programs.
- ▶ We thus first relax the integer program by removing integer constraints on the variables
- ▶ The *relaxed* linear program is then solved using the simplex method
- ▶ The integer solution is then found using one of three strategies:
 - ▶ Enumeration techniques (e.g. Branch-and-bound)
 - ▶ Cutting plane techniques
 - ▶ Group-theoretic techniques
- ▶ We will discuss the first two strategies next,
- ▶ These approaches assume a solution to the *relaxed* linear program has been found
- ▶ If the relaxed linear program is infeasible, the integer program is definitely infeasible

Integer Programming - Ex.

maximise	$5x_1 + 8x_2$
subject to	$x_1 + x_2 \leq 6$
	$5x_1 + 9x_2 \leq 45$
	$x_1, x_2 \geq 0$ and integer

Integer Programming - Ex.

maximise $5x_1 + 8x_2$
subject to $x_1 + x_2 \leq 6$
 $5x_1 + 9x_2 \leq 45$
 $x_1, x_2 \geq 0$ and integer



Integer Programming - Ex.

$$\begin{array}{ll}
 \text{maximise} & 5x_1 + 8x_2 \\
 \text{subject to} & x_1 + x_2 \leq 6 \\
 & 5x_1 + 9x_2 \leq 45 \\
 & x_1, x_2 \geq 0 \text{ and integer}
 \end{array}$$




Table 9.1 Problem features.

	<i>Continuous optimum</i>	<i>Round off</i>	<i>Nearest feasible point</i>	<i>Integer optimum</i>
x_1	$\frac{9}{4} = 2.25$	2	2	0
x_2	$\frac{15}{4} = 3.75$	4	3	5
z	41.25	Infeasible	34	40

Integer Programming - Ex.

$$\begin{array}{ll}
 \text{maximise} & 5x_1 + 8x_2 \\
 \text{subject to} & x_1 + x_2 \leq 6 \\
 & 5x_1 + 9x_2 \leq 45 \\
 & x_1, x_2 \geq 0 \text{ and integer}
 \end{array}$$




Table 9.1 Problem features.

	Continuous optimum	Round off	Nearest feasible point	Integer optimum
x_1	$\frac{9}{4} = 2.25$	2	2	0
x_2	$\frac{15}{4} = 3.75$	4	3	5
z	41.25	Infeasible	34	40

- ▶ the upper bound on $z^* \leq z^0 = 41\frac{1}{4}$, and since z^* must be integral then $z^* \leq 41$.

Method 1: Branch and Bound


- ▶ Branch and bound is another “divide and conquer” strategy.
- ▶ systematically subdivide the linear programming feasible region
- ▶ make assessments of the integer-programming problem based upon these subdivisions.

Method 1: Branch and Bound

- ▶ The linear programming solution is $x_1 = 2\frac{1}{4}$, $x_2 = 3\frac{3}{4}$
- ▶ We can divide the feasible region by *attempting* to make either x_1 or x_2 integer.
- ▶ Assume we attempt to make x_2 integer $\rightarrow x_2 \geq 4$ or $x_2 \leq 3$


Method 1: Branch and Bound

- ▶ The linear programming solution is $x_1 = 2\frac{1}{4}$, $x_2 = 3\frac{3}{4}$
- ▶ We can divide the feasible region by *attempting* to make either x_1 or x_2 integer.
- ▶ Assume we attempt to make x_2 integer $\rightarrow x_2 \geq 4$ or $x_2 \leq 3$





Method 1: Branch and Bound

- ▶ The linear programming solution is $x_1 = 2\frac{1}{4}$, $x_2 = 3\frac{3}{4}$
- ▶ We can divide the feasible region by *attempting* to make either x_1 or x_2 integer.
- ▶ Assume we attempt to make x_2 integer $\rightarrow x_2 \geq 4$ or $x_2 \leq 3$
- ▶ We can visualise this by an **enumeration tree**.




Method 1: Branch and Bound

- ▶ The linear programming solution is $x_1 = 2\frac{1}{4}, x_2 = 3\frac{3}{4}$
- ▶ We can divide the feasible region by *attempting* to make either x_1 or x_2 integer.
- ▶ Assume we attempt to make x_2 integer $\rightarrow x_2 \geq 4$ or $x_2 \leq 3$
- ▶ We can visualise this by an **enumeration tree**.





Method 1: Branch and Bound

- Follow the branch and solve for L_1





Method 1: Branch and Bound

- Follow the branch and solve for L_1



Method 1: Branch and Bound



- Follow the branch and solve for L_1



- L_1 is feasible and x_2 is integer.

Method 1: Branch and Bound

- Follow the branch and solve for L_1





- L_1 is feasible and x_2 is integer.
- Shift to explore changing x_1 to integer.

Method 1: Branch and Bound

- Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$



Method 1: Branch and Bound

- Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$





Method 1: Branch and Bound

- ▶ Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$
- ▶ L_3 is an infeasible problem





Method 1: Branch and Bound

- ▶ Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$
- ▶ L_3 is an infeasible problem
- ▶ The branch in the enumeration tree does not need to be explored further





Method 1: Branch and Bound

- ▶ Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$
- ▶ L_3 is an infeasible problem
- ▶ The branch in the enumeration tree does not need to be explored further
- ▶ At this point we can explore L_2 or L_4





Method 1: Branch and Bound

- ▶ Subdivide L_1 further into regions L_3 with $x_1 \geq 2$ and L_4 with $x_1 \leq 1$
- ▶ L_3 is an infeasible problem
- ▶ The branch in the enumeration tree does not need to be explored further
- ▶ At this point we can explore L_2 or L_4
- ▶ The decision can be done arbitrarily, or using some heuristics





Method 1: Branch and Bound

- Let's choose L_4 , solution to L_4 is $(x_1, x_2) = (1, 4\frac{4}{9})$.




Method 1: Branch and Bound



- ▶ Let's choose L_4 , solution to L_4 is $(x_1, x_2) = (1, 4\frac{4}{9})$.
- ▶ x_2 is not yet an integer \rightarrow explore $x_2 \leq 4$ and $x_2 \geq 5$



Method 1: Branch and Bound





Method 1: Branch and Bound





Method 1: Branch and Bound

- L_5 is a feasible problem with the optimal solution $(x_1, x_2) = (1, 4)$ and $z_{L_5} = 37$.





Method 1: Branch and Bound

- ▶ L_5 is a feasible problem with the optimal solution $(x_1, x_2) = (1, 4)$ and $z_{L_5} = 37$.
- ▶ We know that $37 \leq z^* \leq 41$.



Method 1: Branch and Bound

- ▶ L_5 is a feasible problem with the optimal solution $(x_1, x_2) = (1, 4)$ and $z_{L_5} = 37$.
- ▶ We know that $37 \leq z^* \leq 41$.
- ▶ L_2 or L_6 might contain a better answer




Method 1: Branch and Bound

- Solving for L_2 and L_6 accordingly
optimal answers are integers


Method 1: Branch and Bound

- ▶ Solving for L_2 and L_6 accordingly optimal answers are integers
- ▶ L_6 has the optimal solution
 $z^* = 40$



Method 1: Branch and Bound

- ▶ Solving for L_2 and L_6 accordingly optimal answers are integers
- ▶ L_6 has the optimal solution $z^* = 40$
- ▶ The optimum answer to the integer program is $(x_1, x_2) = (0, 5)$



Method 2: Cutting Planes

- ▶ Cutting-plane algorithm solves integer programs by modifying *relaxed* linear-programming solutions until the integer solution is obtained.
- ▶ It does not partition the feasible region, as in branch-and-bound
- ▶ It instead works with a single linear program, which is refined by adding new constraints.
- ▶ The new constraints successively reduce the feasible region until an integer optimal solution is found.
- ▶ In practice, the branch-and-bound procedures almost always outperform the cutting-plane algorithm.
- ▶ Cutting-plane was though the first algorithm developed for integer programming that could be proven to converge in a finite number of steps.

Method 2: Cutting Planes

$$\begin{array}{ll}\text{maximise} & 5x_1 + 8x_2 \\ \text{subject to} & x_1 + x_2 \leq 6 \\ & 5x_1 + 9x_2 \leq 45 \\ & x_1, x_2 \geq 0 \text{ and integer}\end{array}$$

change to slack form

$$\begin{array}{ll}z = & 5x_1 + 8x_2 \\ s_1 = & 6 - x_1 - x_2 \\ s_2 = & 45 - 5x_1 - 9x_2\end{array}$$

Method 2: Cutting Planes

$$\begin{array}{ll}\text{maximise} & 5x_1 + 8x_2 \\ \text{subject to} & x_1 + x_2 \leq 6 \\ & 5x_1 + 9x_2 \leq 45 \\ & x_1, x_2 \geq 0 \text{ and integer}\end{array}$$

change to slack form

$$\begin{array}{ll}z = & 5x_1 + 8x_2 \\ s_1 = & 6 - x_1 - x_2 \\ s_2 = & 45 - 5x_1 - 9x_2\end{array}$$

Solving using the Simplex method results in:

$$\begin{array}{llll}z = & 41\frac{1}{4} & - & \frac{5}{4}s_1 & - & \frac{3}{4}s_2 \\ x_1 = & \frac{9}{4} & - & \frac{9}{4}s_1 & + & \frac{1}{4}s_2 \\ x_2 = & \frac{15}{4} & + & \frac{5}{4}s_1 & - & \frac{1}{4}s_2\end{array}$$

Method 2: Cutting Planes

$$\begin{array}{rclclcl} z = & 41\frac{1}{4} & - & \frac{5}{4}s_1 & - & \frac{3}{4}s_2 \\ x_1 = & \frac{9}{4} & - & \frac{9}{4}s_1 & + & \frac{1}{4}s_2 \\ x_2 = & \frac{15}{4} & + & \frac{5}{4}s_1 & - & \frac{1}{4}s_2 \end{array}$$

Method 2: Cutting Planes

$$\begin{array}{rclclcl} Z & = & 41\frac{1}{4} & - & \frac{5}{4}S_1 & - & \frac{3}{4}S_2 \\ X_1 & = & \frac{9}{4} & - & \frac{9}{4}S_1 & + & \frac{1}{4}S_2 \\ X_2 & = & \frac{15}{4} & + & \frac{5}{4}S_1 & - & \frac{1}{4}S_2 \end{array}$$

Rewrite as

$$\begin{array}{rclclclcl} -Z & & & - & \frac{5}{4}S_1 & - & \frac{3}{4}S_2 & = & -41\frac{1}{4} \\ & X_1 & & + & \frac{9}{4}S_1 & - & \frac{1}{4}S_2 & = & \frac{9}{4} \\ & & X_2 & - & \frac{5}{4}S_1 & + & \frac{1}{4}S_2 & = & \frac{15}{4} \end{array}$$

Method 2: Cutting Planes

$$\begin{array}{rclclcl}
 Z & = & 41\frac{1}{4} & - & \frac{5}{4}S_1 & - & \frac{3}{4}S_2 \\
 X_1 & = & \frac{9}{4} & - & \frac{9}{4}S_1 & + & \frac{1}{4}S_2 \\
 X_2 & = & \frac{15}{4} & + & \frac{5}{4}S_1 & - & \frac{1}{4}S_2
 \end{array}$$

Rewrite as

$$\begin{array}{rclclcl}
 -Z & & & - & \frac{5}{4}S_1 & - & \frac{3}{4}S_2 & = & -41\frac{1}{4} \\
 & X_1 & & + & \frac{9}{4}S_1 & - & \frac{1}{4}S_2 & = & \frac{9}{4} \\
 & & X_2 & - & \frac{5}{4}S_1 & + & \frac{1}{4}S_2 & = & \frac{14}{4}
 \end{array}$$

and rewrite as

$$\begin{array}{rclclcl}
 -Z & & & - & 2S_1 & - & S_2 & + & 42 & = & \frac{3}{4} & - & \frac{3}{4}S_1 & - & \frac{1}{4}S_2 \\
 & X_1 & & + & 2S_1 & - & S_2 & - & 2 & = & \frac{1}{4} & - & \frac{1}{4}S_1 & - & \frac{3}{4}S_2 \\
 & & X_2 & - & 2S_1 & & & - & 3 & = & \frac{3}{4} & - & \frac{3}{4}S_1 & - & \frac{1}{4}S_2
 \end{array}$$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$		$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1	$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
			x_2	$-$	$2s_1$		3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

- ▶ These algebraic manipulations
 - ▶ isolate integer coefficients to one side of the equalities and fractions to the other
 - ▶ constant terms on the righthand side are all nonnegative
 - ▶ slack variable coefficients on the righthand side are all nonpositive

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$			$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1		$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$			$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$			$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1		$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$			$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

► $s_1 \geq 0$ and $s_2 \geq 0$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$			$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1		$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$			$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

- ▶ $s_1 \geq 0$ and $s_2 \geq 0$
- ▶ Since s_1 and s_2 appear to the right with negative coefficients, then

$$\frac{3}{4} \leq \frac{3}{4}s_1 + \frac{1}{4}s_2$$

$$\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 \leq 0$$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$			$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1		$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$			$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

► $s_1 \geq 0$ and $s_2 \geq 0$

► Since s_1 and s_2 appear to the right with negative coefficients, then

$$\frac{3}{4} \leq \frac{3}{4}s_1 + \frac{1}{4}s_2$$

$$\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 \leq 0$$

Add a new slack variable $s_3 \geq 0$ where

$$\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 = 0$$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$		$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1	$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
			x_2	$-$	$2s_1$		3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

And similarly for the remaining constraints

$$\begin{aligned} \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 &= 0, & s_3 &\geq 0 & \text{and} & \text{integer} \\ \frac{1}{4} - \frac{1}{4}s_1 - \frac{3}{4}s_2 + s_4 &= 0, & s_4 &\geq 0 & \text{and} & \text{integer} \\ \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_5 &= 0, & s_5 &\geq 0 & \text{and} & \text{integer} \end{aligned}$$

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$			$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1		$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$			$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

And similarly for the remaining constraints

$$\begin{aligned}\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 &= 0, & s_3 &\geq 0 & \text{and} & \text{integer} \\ \frac{1}{4} - \frac{1}{4}s_1 - \frac{3}{4}s_2 + s_4 &= 0, & s_4 &\geq 0 & \text{and} & \text{integer} \\ \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_5 &= 0, & s_5 &\geq 0 & \text{and} & \text{integer}\end{aligned}$$

- Notice that first and third constraints are identical.

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$		$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1	$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
			x_2	$-$	$2s_1$		3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

And similarly for the remaining constraints

$$\begin{aligned} \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 &= 0, & s_3 &\geq 0 & \text{and} & \text{integer} \\ \frac{1}{4} - \frac{1}{4}s_1 - \frac{3}{4}s_2 + s_4 &= 0, & s_4 &\geq 0 & \text{and} & \text{integer} \\ \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_5 &= 0, & s_5 &\geq 0 & \text{and} & \text{integer} \end{aligned}$$

- Notice that first and third constraints are identical.
- The new equations are called **cuts**, because their derivation did not exclude any integer solutions to the problem.

Method 2: Cutting Planes

Let's look at the right hand side,

$-Z$		$-$	$2s_1$	$-$	s_2	$+$	42	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$
	x_1	$+$	$2s_1$	$-$	s_2	$-$	2	$=$	$\frac{1}{4}$	$-$	$\frac{1}{4}s_1$	$-$	$\frac{3}{4}s_2$
		x_2	$-$	$2s_1$		$-$	3	$=$	$\frac{3}{4}$	$-$	$\frac{3}{4}s_1$	$-$	$\frac{1}{4}s_2$

And similarly for the remaining constraints

$$\begin{aligned}\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 &= 0, & s_3 &\geq 0 & \text{and} & \text{integer} \\ \frac{1}{4} - \frac{1}{4}s_1 - \frac{3}{4}s_2 + s_4 &= 0, & s_4 &\geq 0 & \text{and} & \text{integer} \\ \frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_5 &= 0, & s_5 &\geq 0 & \text{and} & \text{integer}\end{aligned}$$

- ▶ Notice that first and third constraints are identical.
- ▶ The new equations are called **cuts**, because their derivation did not exclude any integer solutions to the problem.
- ▶ Any integer feasible solution to the original problem must satisfy the cut constraints
- ▶ A cut is to cut away the optimal linear-programming solution from the feasible region without excluding any feasible integer solution.

Method 2: Cutting Planes

$$\begin{aligned}\frac{3}{4} - \frac{3}{4}s_1 - \frac{1}{4}s_2 + s_3 &= 0, & s_3 &\geq 0 & \text{and} & \text{integer} \\ \frac{1}{4} - \frac{1}{4}s_1 - \frac{3}{4}s_2 + s_4 &= 0, & s_4 &\geq 0 & \text{and} & \text{integer}\end{aligned}$$

Recall from the slack form that:

$$\begin{aligned}z &= 5x_1 + 8x_2 \\ s_1 &= 6 - x_1 - x_2 \\ s_2 &= 45 - 5x_1 - 9x_2\end{aligned}$$

By replacing, you achieve two cut inequalities:

$$\begin{aligned}2x_1 + 3x_2 &\leq 15 \\ 4x_1 + 7x_2 &\leq 35\end{aligned}$$


Method 2: Cutting Planes

- ▶ The usual strategy is to add cuts (usually only one) to the constraints, then solve the resulting linear program.
- ▶ If the optimal values are all integer, they are optimal
- ▶ Otherwise, a new cut is derived from the new optimal linear program, and is appended to the constraints

Method 2: Cutting Planes

$$C_1: 2x_1 + 3x_2 \leq 15$$

$$C_2: 4x_1 + 7x_2 \leq 35$$



- ▶ In the example before, the first cut leads directly to the optimal solution, but the second does not.
- ▶ Also notice that the first cut cuts deeper into the feasible region, but it is difficult to determine which cuts will be deep in this sense.
- ▶ If the algorithm chooses a cut that removes very little from the feasible region, the algorithm's performance will be poor

Method 2: Cutting Planes - Time Complexity

- ▶ In 1958, R. Gomory made a theoretical break-through
- ▶ He showed that integer programs can be solved by *some* linear program (i.e. the relaxed linear program plus added constraints)

Method 2: Cutting Planes - Time Complexity

- ▶ In 1958, R. Gomory made a theoretical break-through
- ▶ He showed that integer programs can be solved by *some* linear program (i.e. the relaxed linear program plus added constraints)
- ▶ Unfortunately, the number of cuts to be added, though finite, is usually large
- ▶ This result does not have important practical ramifications

Method 2: Cutting Planes - Time Complexity

- ▶ In 1958, R. Gomory made a theoretical break-through
- ▶ He showed that integer programs can be solved by *some* linear program (i.e. the relaxed linear program plus added constraints)
- ▶ Unfortunately, the number of cuts to be added, though finite, is usually large
- ▶ This result does not have important practical ramifications
- ▶ In practice, the branch-and-bound procedures almost always outperform the cutting-plane algorithm.

Further Reading

- ▶ **Applied Mathematical Programming** (available online)
Bradley, Hax and Magnanti.
Addison-Wesley, 1977.
 - ▶ Chapter 9 – Integer Programming