

# Class 10 Structural Bioinformatics

Ani A16647613

## Section 1 Intro to RCSB Protein Data Bank

Here we examine the size and composition of the main database of biomolecular structures- the PDB.

```
pdbstats <- read.csv("datasummary.csv", row.names=1)
head(pdbstats)
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	161,663	12,592	12,337	200	74	32
Protein/Oligosaccharide	9,348	2,167	34	8	2	0
Protein/NA	8,404	3,924	286	7	0	0
Nucleic acid (only)	2,758	125	1,477	14	3	1
Other	164	9	33	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	186,898					
Protein/Oligosaccharide	11,559					
Protein/NA	12,621					
Nucleic acid (only)	4,378					
Other	206					
Oligosaccharide (only)	22					

Q1. What percentage of structures in the PDB are solved by X-ray and electron microscopy?

My pdb stats data has numbers with commas in them which may cause problems. Testing it out:

```
pdbstats$X.ray
```

```
[1] "161,663" "9,348" "8,404" "2,758" "164" "11"
```

```
as.numeric(pdbstats$X.ray)
```

Warning: NAs introduced by coercion

```
[1] NA NA NA NA 164 11
```

Found a function called `gsub()` so now we figure out how it works

```
x <- "49,234"  
sum(as.numeric(gsub(",", "", x)))
```

```
[1] 49234
```

Can use this function for every column

```
commasum <- function(x) {  
  sum(as.numeric(gsub(",", "", x)))  
}
```

```
commasum(pdbstats$X.ray)
```

```
[1] 182348
```

```
totals <- apply(pdbstats, 2, commasum)
```

```
round(totals/totals["Total"] * 100, 2)
```

X.ray	EM	NMR	Multiple.methods
84.54	8.72	6.57	0.11
Neutron	Other	Total	
0.04	0.02	100.00	

With X.ray 84.54 percent and EM 8.72 percent

```
# Percentage of Xray and EM  
84.54 + 8.72
```

```
[1] 93.26
```

Q2. What proportion of structures in the PDB are protein?

```
proteintotal <- as.numeric(gsub(",", "", pdbstats[1,7]))  
  
round(proteintotal/totals["Total"] * 100, 2)
```

```
Total  
86.65
```


```
(215684/249751891 * 100)
```

```
[1] 0.08635931
```

## Section 2 Visualizing Protein Structure

learning the basis of Mol\* (mol-star)

PDB code with 1HSG



PDB with Asp 25's showing

## Predict the dynamics or flexibility of an important protein:

```
library(bio3d)  
  
hiv <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
hiv
```

Call: read.pdb(file = "1hsg")

Total Models#: 1

Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)




Figure 1: HIV-Pr with a bound inhibitor showing two crucial Asp 25 amino acids

Protein Atoms#: 1514 (residues/Calpha atoms#: 198)

Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 172 (residues: 128)

Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]

Protein sequence:

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
head(hiv$atom)
```

```
type eleno elety alt resid chain resno insert x y z o b
```

```

1 ATOM      1      N <NA>  PRO      A      1      <NA> 29.361 39.686 5.862 1 38.10
2 ATOM      2      CA <NA>  PRO      A      1      <NA> 30.307 38.663 5.319 1 40.62
3 ATOM      3      C  <NA>  PRO      A      1      <NA> 29.760 38.071 4.022 1 42.64
4 ATOM      4      O  <NA>  PRO      A      1      <NA> 28.600 38.302 3.676 1 43.40
5 ATOM      5      CB <NA>  PRO      A      1      <NA> 30.508 37.541 6.342 1 37.87
6 ATOM      6      CG <NA>  PRO      A      1      <NA> 29.296 37.591 7.162 1 38.40
      segid elesy charge
1  <NA>      N  <NA>
2  <NA>      C  <NA>
3  <NA>      C  <NA>
4  <NA>      O  <NA>
5  <NA>      C  <NA>
6  <NA>      C  <NA>

```

`pdbseq(hiv)`

```

  1   2   3   4   5   6   7   8   9  10  11  12  13  14  15  16  17  18  19  20
"P" "Q" "I" "T" "L" "W" "Q" "R" "P" "L" "V" "T" "I" "K" "I" "G" "G" "Q" "L" "K"
 21  22  23  24  25  26  27  28  29  30  31  32  33  34  35  36  37  38  39  40
"E" "A" "L" "L" "D" "T" "G" "A" "D" "D" "T" "V" "L" "E" "E" "M" "S" "L" "P" "G"
 41  42  43  44  45  46  47  48  49  50  51  52  53  54  55  56  57  58  59  60
"R" "W" "K" "P" "K" "M" "I" "G" "G" "I" "G" "G" "F" "I" "K" "V" "R" "Q" "Y" "D"
 61  62  63  64  65  66  67  68  69  70  71  72  73  74  75  76  77  78  79  80
"Q" "I" "L" "I" "E" "I" "C" "G" "H" "K" "A" "I" "G" "T" "V" "L" "V" "G" "P" "T"
 81  82  83  84  85  86  87  88  89  90  91  92  93  94  95  96  97  98  99   1
"P" "V" "N" "I" "I" "G" "R" "N" "L" "L" "T" "Q" "I" "G" "C" "T" "L" "N" "F" "P"
  2   3   4   5   6   7   8   9  10  11  12  13  14  15  16  17  18  19  20  21
"Q" "I" "T" "L" "W" "Q" "R" "P" "L" "V" "T" "I" "K" "I" "G" "G" "Q" "L" "K" "E"
 22  23  24  25  26  27  28  29  30  31  32  33  34  35  36  37  38  39  40  41
"A" "L" "L" "D" "T" "G" "A" "D" "D" "T" "V" "L" "E" "E" "M" "S" "L" "P" "G" "R"
 42  43  44  45  46  47  48  49  50  51  52  53  54  55  56  57  58  59  60  61
"W" "K" "P" "K" "M" "I" "G" "G" "I" "G" "G" "F" "I" "K" "V" "R" "Q" "Y" "D" "Q"
 62  63  64  65  66  67  68  69  70  71  72  73  74  75  76  77  78  79  80  81
"I" "L" "I" "E" "I" "C" "G" "H" "K" "A" "I" "G" "T" "V" "L" "V" "G" "P" "T" "P"
 82  83  84  85  86  87  88  89  90  91  92  93  94  95  96  97  98  99
"V" "N" "I" "I" "G" "R" "N" "L" "L" "T" "Q" "I" "G" "C" "T" "L" "N" "F"

```

Doing NMA:

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file  
PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLDGFRTIPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```


```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

```
modes <- nma(adk)
```

```
Building Hessian... Done in 0.04 seconds.
```

```
Diagonalizing Hessian... Done in 0.38 seconds.
```

```
plot(modes)
```



There are certain regions of the sequence that is more flexible.

Make a “movie” called a trajectory to predict motions:

```
mktrj(modes, file = "adk_m7.pdb")
```

Then we can open it in Mol\*