

Fixed freq. identities for the Green function and uniqueness results for passive imaging


Alexey Agaltsov

Max Planck Institute for Solar System Research, Germany
agaltsov@mps.mpg.de

Joint work with T. Hohage and R. G. Novikov
SIAM J. Appl. Math. 2018

QIPA2018, MIPT – December 3, 2018

I. Motivation – Classical inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Point sources of waves on $\partial\Omega$

Classical inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in medical diagnostics, ocean tomography, . . .

I. Motivation – Classical inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Point sources of waves on $\partial\Omega$

Classical inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in medical diagnostics, ocean tomography, . . .

I. Motivation – Classical inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Point sources of waves on $\partial\Omega$

Classical inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in medical diagnostics, ocean tomography, . . .

I. Motivation – Classical inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Point sources of waves on $\partial\Omega$

Classical inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in medical diagnostics, ocean tomography, . . .

I. Motivation – Classical inverse problem



- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Point sources of waves on $\partial\Omega$

Classical inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in medical diagnostics, ocean tomography, . . .

I. Motivation – Classical inverse problem

- Green's function for the acoustic equation:

$$\begin{aligned} -\nabla_x \left(\frac{1}{\rho} \nabla_x G(x, y) \right) - \frac{\omega^2}{\rho c^2} G(x, y) &= \delta_y(x) \quad x \in \Omega, \\ \frac{\partial G}{\partial \nu}(x, y) &= i \frac{\omega}{c} G(x, y) \quad x \in \partial\Omega \end{aligned}$$

Classical inverse problem: *Given $G(x, y)$ for all $x, y \in \partial\Omega$ for a finite number of ω , find $\rho(x)$, $c(x)$ for $x \in \Omega$*

I. Motivation – Classical inverse problem

- Green's function for the acoustic equation:

$$\begin{aligned}-\nabla_x \left(\frac{1}{\rho} \nabla_x G(x, y) \right) - \frac{\omega^2}{\rho c^2} G(x, y) &= \delta_y(x) \quad x \in \Omega, \\ \frac{\partial G}{\partial \nu}(x, y) &= i \frac{\omega}{c} G(x, y) \quad x \in \partial\Omega\end{aligned}$$

Classical inverse problem: *Given $G(x, y)$ for all $x, y \in \partial\Omega$ for a finite number of ω , find $\rho(x)$, $c(x)$ for $x \in \Omega$*


I. Motivation – Classical inverse problem

- Green's function for the acoustic equation:

$$\begin{aligned} -\nabla_x \left(\frac{1}{\rho} \nabla_x G(x, y) \right) - \frac{\omega^2}{\rho c^2} G(x, y) &= \delta_y(x) \quad x \in \Omega, \\ \frac{\partial G}{\partial \nu}(x, y) &= i \frac{\omega}{c} G(x, y) \quad x \in \partial\Omega \end{aligned}$$

Classical inverse problem: *Given $G(x, y)$ for all $x, y \in \partial\Omega$ for a finite number of ω , find $\rho(x)$, $c(x)$ for $x \in \Omega$*

I. Motivation – Passive inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Random sources of waves in $\overline{\Omega}$

Passive inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in helioseismology [5], passive ocean tomography [3], ultrasonics [11], seismology [10]. . .

I. Motivation – Passive inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Random sources of waves in $\overline{\Omega}$

Passive inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in helioseismology [5], passive ocean tomography [3], ultrasonics [11], seismology [10]. . .

I. Motivation – Passive inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Random sources of waves in $\overline{\Omega}$

Passive inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in helioseismology [5], passive ocean tomography [3], ultrasonics [11], seismology [10]. . .

I. Motivation – Passive inverse problem




- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Random sources of waves in $\overline{\Omega}$

Passive inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in helioseismology [5], passive ocean tomography [3], ultrasonics [11], seismology [10]. . .

I. Motivation – Passive inverse problem



- Unknown scatterer in a bounded domain $\Omega \subset \mathbb{R}^d$
- Random sources of waves in $\overline{\Omega}$

Passive inverse problem: *Recover scatterer parameters from boundary measurements of scattered waves*

- Applications in helioseismology [5], passive ocean tomography [3], ultrasonics [11], seismology [10]. . .

I. Motivation – Passive inverse problem

- Random acoustic noise field:

$$\begin{aligned} -\nabla\left(\frac{1}{\rho}\nabla u(x)\right) - \frac{\omega^2}{\rho c^2}u(x) &= q(x) \quad x \in \Omega, \\ \frac{\partial u}{\partial \nu}(x) &= i\frac{\omega}{c}u(x) \quad x \in \partial\Omega, \end{aligned}$$

where $q(x)$ is a random source field in $\overline{\Omega}$

Passive inverse problem: *Given $\langle u(x_1)\overline{u(x_2)} \rangle$ for all $x_1, x_2 \in \partial\Omega$ for a finite number of ω , find $\rho(x), c(x)$ for $x \in \Omega$*

I. Motivation – Passive inverse problem

- Random acoustic noise field:

$$\begin{aligned}-\nabla\left(\frac{1}{\rho}\nabla u(x)\right)-\frac{\omega^2}{\rho c^2}u(x)&=q(x) \quad x\in\Omega, \\ \frac{\partial u}{\partial \nu}(x)&=i\frac{\omega}{c}u(x) \quad x\in\partial\Omega,\end{aligned}$$

where $q(x)$ is a random source field in $\overline{\Omega}$

Passive inverse problem: *Given $\langle u(x_1)\overline{u(x_2)}\rangle$ for all $x_1, x_2 \in \partial\Omega$ for a finite number of ω , find $\rho(x), c(x)$ for $x \in \Omega$*

I. Motivation – Passive inverse problem

- Random acoustic noise field:

$$\begin{aligned}-\nabla\left(\frac{1}{\rho}\nabla u(x)\right)-\frac{\omega^2}{\rho c^2}u(x)&=q(x) \quad x\in\Omega, \\ \frac{\partial u}{\partial\nu}(x)&=i\frac{\omega}{c}u(x) \quad x\in\partial\Omega,\end{aligned}$$

where $q(x)$ is a random source field in $\overline{\Omega}$

Passive inverse problem: *Given $\langle u(x_1)\overline{u(x_2)}\rangle$ for all $x_1, x_2 \in \partial\Omega$ for a finite number of ω , find $\rho(x), c(x)$ for $x \in \Omega$*

I. Motivation – Passive inverse problem


- Random acoustic noise field:

$$\begin{aligned}-\nabla\left(\frac{1}{\rho}\nabla u(x)\right)-\frac{\omega^2}{\rho c^2}u(x)&=q(x) \quad x\in\Omega, \\ \frac{\partial u}{\partial\nu}(x)&=i\frac{\omega}{c}u(x) \quad x\in\partial\Omega,\end{aligned}$$

where $q(x)$ is a random source field in $\overline{\Omega}$

Passive inverse problem: *Given $\langle u(x_1)\overline{u(x_2)}\rangle$ for all $x_1, x_2 \in \partial\Omega$ for a finite number of ω , find $\rho(x), c(x)$ for $x \in \Omega$*

I. Motivation – Local helioseismology




Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Local helioseismology




Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Local helioseismology




Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Local helioseismology




Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Local helioseismology




Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Local helioseismology



Solar Dopplergram
(<https://sdo.gsfc.nasa.gov>)

- Turbulent convection excites acoustic waves
- Acoustic waves travel and scatter inside of the Sun
- Acoustic field on the surface manifests as surface oscillations

Local helioseismology: *Recover the Sun parameters from the measurements of the acoustic field on its surface*

- L. Gizon et al. (*Astronomy & Astrophysics* 600, 2017)

I. Motivation – Classical vs. passive

random sources	x	point sources
$-\nabla(\frac{1}{\rho}\nabla u) - \frac{\omega^2}{\rho c^2}u = q$	Ω	$-\nabla(\frac{1}{\rho}\nabla G(\cdot, y)) - \frac{\omega^2}{\rho c^2}G(\cdot, y) = \delta_y$
$\frac{\partial u}{\partial \nu} = i\frac{\omega}{c}u$	$\partial\Omega$	$\frac{\partial G}{\partial \nu}(\cdot, y) = i\frac{\omega}{c}G(\cdot, y)$

- Assumption:

$$\langle q(x_1)\overline{q(x_2)} \rangle = \frac{|S(\omega)|^2}{\rho c} \delta_{\partial\Omega}(x_1 - x_2)$$

Theorem ([9], \approx [4, 8, 11, 5]). *The following formula holds:*

$$\operatorname{Im} G(x_1, x_2) = \frac{\omega}{|S(\omega)|^2} \langle u(x_1)\overline{u(x_2)} \rangle$$

I. Motivation – Classical vs. passive

random sources	x	point sources
$-\nabla(\frac{1}{\rho}\nabla u) - \frac{\omega^2}{\rho c^2}u = q$ $\frac{\partial u}{\partial \nu} = i\frac{\omega}{c}u$	Ω $\partial\Omega$	$-\nabla(\frac{1}{\rho}\nabla G(\cdot, y)) - \frac{\omega^2}{\rho c^2}G(\cdot, y) = \delta_y$ $\frac{\partial G}{\partial \nu}(\cdot, y) = i\frac{\omega}{c}G(\cdot, y)$

- Assumption:

$$\langle q(x_1)\overline{q(x_2)} \rangle = \frac{|S(\omega)|^2}{\rho c} \delta_{\partial\Omega}(x_1 - x_2)$$

Theorem ([9], \approx [4, 8, 11, 5]). *The following formula holds:*

$$\operatorname{Im} G(x_1, x_2) = \frac{\omega}{|S(\omega)|^2} \langle u(x_1)\overline{u(x_2)} \rangle$$

I. Motivation – Classical vs. passive

random sources	x	point sources
$-\nabla(\frac{1}{\rho}\nabla u) - \frac{\omega^2}{\rho c^2}u = q$	Ω	$-\nabla(\frac{1}{\rho}\nabla G(\cdot, y)) - \frac{\omega^2}{\rho c^2}G(\cdot, y) = \delta_y$
$\frac{\partial u}{\partial \nu} = i\frac{\omega}{c}u$	$\partial\Omega$	$\frac{\partial G}{\partial \nu}(\cdot, y) = i\frac{\omega}{c}G(\cdot, y)$

- Assumption:

$$\langle q(x_1)\overline{q(x_2)} \rangle = \frac{|S(\omega)|^2}{\rho c} \delta_{\partial\Omega}(x_1 - x_2)$$

Theorem ([9], \approx [4, 8, 11, 5]). *The following formula holds:*

$$\operatorname{Im} G(x_1, x_2) = \frac{\omega}{|S(\omega)|^2} \langle u(x_1)\overline{u(x_2)} \rangle$$

I. Motivation – Classical vs. passive

random sources	x	point sources
$-\nabla(\frac{1}{\rho}\nabla u) - \frac{\omega^2}{\rho c^2}u = q$	Ω	$-\nabla(\frac{1}{\rho}\nabla G(\cdot, y)) - \frac{\omega^2}{\rho c^2}G(\cdot, y) = \delta_y$
$\frac{\partial u}{\partial \nu} = i\frac{\omega}{c}u$	$\partial\Omega$	$\frac{\partial G}{\partial \nu}(\cdot, y) = i\frac{\omega}{c}G(\cdot, y)$

- Assumption:

$$\langle q(x_1)\overline{q(x_2)} \rangle = \frac{|S(\omega)|^2}{\rho c} \delta_{\partial\Omega}(x_1 - x_2)$$

Theorem ([9], \approx [4, 8, 11, 5]). *The following formula holds:*

$$\operatorname{Im} G(x_1, x_2) = \frac{\omega}{|S(\omega)|^2} \langle u(x_1)\overline{u(x_2)} \rangle$$

II. Mathematical setup

- $G_n^+ = G_n^+(x, y)$ is the outgoing Green function

$$\begin{aligned}(-\Delta_x - k^2 n(x))G_n^+(x, y) &= \delta_y(x) \quad x \in \mathbb{R}^d, \\ \left(\frac{\partial}{\partial |x|} - ik\right)G_n^+(x, y) &= o(|x|^{\frac{1-d}{2}}), \quad |x| \rightarrow +\infty.\end{aligned}$$

- $n(x)$ real, $n(x) = 1$ outside a bounded domain Ω

Inverse problem: Given $\text{Im } G_n^+(x_1, x_2)$ at fixed $k > 0$ for all $x_1, x_2 \in \partial\Omega$, find $n(x)$ for $x \in \Omega$

Focus of this talk: How to find $\text{Re } G_n^+(x_1, x_2)$ if $\text{Im } G_n^+(x_1, x_2)$ is given at one fixed k

II. Mathematical setup

- $G_n^+ = G_n^+(x, y)$ is the outgoing Green function

$$\begin{aligned}(-\Delta_x - k^2 n(x))G_n^+(x, y) &= \delta_y(x) \quad x \in \mathbb{R}^d, \\ \left(\frac{\partial}{\partial |x|} - ik\right)G_n^+(x, y) &= o(|x|^{\frac{1-d}{2}}), \quad |x| \rightarrow +\infty.\end{aligned}$$

- $n(x)$ real, $n(x) = 1$ outside a bounded domain Ω

Inverse problem: Given $\text{Im } G_n^+(x_1, x_2)$ at fixed $k > 0$ for all $x_1, x_2 \in \partial\Omega$, find $n(x)$ for $x \in \Omega$

Focus of this talk: How to find $\text{Re } G_n^+(x_1, x_2)$ if $\text{Im } G_n^+(x_1, x_2)$ is given at one fixed k

II. Mathematical setup

- $G_n^+ = G_n^+(x, y)$ is the outgoing Green function

$$\begin{aligned}(-\Delta_x - k^2 n(x)) G_n^+(x, y) &= \delta_y(x) \quad x \in \mathbb{R}^d, \\ \left(\frac{\partial}{\partial |x|} - ik\right) G_n^+(x, y) &= o(|x|^{\frac{1-d}{2}}), \quad |x| \rightarrow +\infty.\end{aligned}$$

- $n(x)$ real, $n(x) = 1$ outside a bounded domain Ω

Inverse problem: *Given $\text{Im } G_n^+(x_1, x_2)$ at fixed $k > 0$ for all $x_1, x_2 \in \partial\Omega$, find $n(x)$ for $x \in \Omega$*

Focus of this talk: *How to find $\text{Re } G_n^+(x_1, x_2)$ if $\text{Im } G_n^+(x_1, x_2)$ is given at one fixed k*

II. Mathematical setup

- $G_n^+ = G_n^+(x, y)$ is the outgoing Green function

$$\begin{aligned}(-\Delta_x - k^2 n(x)) G_n^+(x, y) &= \delta_y(x) \quad x \in \mathbb{R}^d, \\ \left(\frac{\partial}{\partial |x|} - ik\right) G_n^+(x, y) &= o(|x|^{\frac{1-d}{2}}), \quad |x| \rightarrow +\infty.\end{aligned}$$

- $n(x)$ real, $n(x) = 1$ outside a bounded domain Ω

Inverse problem: *Given $\text{Im } G_n^+(x_1, x_2)$ at fixed $k > 0$ for all $x_1, x_2 \in \partial\Omega$, find $n(x)$ for $x \in \Omega$*

Focus of this talk: *How to find $\text{Re } G_n^+(x_1, x_2)$ if $\text{Im } G_n^+(x_1, x_2)$ is given at one fixed k*

II. Mathematical setup

- $G_n^+ = G_n^+(x, y)$ is the outgoing Green function

$$\begin{aligned}(-\Delta_x - k^2 n(x)) G_n^+(x, y) &= \delta_y(x) \quad x \in \mathbb{R}^d, \\ \left(\frac{\partial}{\partial |x|} - ik\right) G_n^+(x, y) &= o(|x|^{\frac{1-d}{2}}), \quad |x| \rightarrow +\infty.\end{aligned}$$

- $n(x)$ real, $n(x) = 1$ outside a bounded domain Ω

Inverse problem: *Given $\operatorname{Im} G_n^+(x_1, x_2)$ at fixed $k > 0$ for all $x_1, x_2 \in \partial\Omega$, find $n(x)$ for $x \in \Omega$*

Focus of this talk: *How to find $\operatorname{Re} G_n^+(x_1, x_2)$ if $\operatorname{Im} G_n^+(x_1, x_2)$ is given at one fixed k*

Two important linear operators:

- Single layer potential $\mathcal{G}_n: H^{-\frac{1}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$:

$$\mathcal{G}_n f(x) = \int_{\partial\Omega} G_n^+(x, y) f(y) dy, \quad x \in \partial\Omega$$

- Dirichlet-to-Neumann map $\Phi_n: H^{\frac{1}{2}}(\partial\Omega) \rightarrow H^{-\frac{1}{2}}(\partial\Omega)$:
 $\Phi_n u_0 = \frac{\partial u}{\partial \nu},$

$$\begin{cases} \Delta u + k^2 n(x)u = 0 & x \in \Omega, \\ u|_{\partial\Omega} = u_0. \end{cases}$$

Two important linear operators:

- Single layer potential $\mathcal{G}_n: H^{-\frac{1}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$:

$$\mathcal{G}_n f(x) = \int_{\partial\Omega} G_n^+(x, y) f(y) dy, \quad x \in \partial\Omega$$

- Dirichlet-to-Neumann map $\Phi_n: H^{\frac{1}{2}}(\partial\Omega) \rightarrow H^{-\frac{1}{2}}(\partial\Omega)$:
 $\Phi_n u_0 = \frac{\partial u}{\partial \nu},$

$$\begin{cases} \Delta u + k^2 n(x)u = 0 & x \in \Omega, \\ u|_{\partial\Omega} = u_0. \end{cases}$$

Two important linear operators:

- Single layer potential $\mathcal{G}_n: H^{-\frac{1}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$:

$$\mathcal{G}_n f(x) = \int_{\partial\Omega} G_n^+(x, y) f(y) dy, \quad x \in \partial\Omega$$

- Dirichlet-to-Neumann map $\Phi_n: H^{\frac{1}{2}}(\partial\Omega) \rightarrow H^{-\frac{1}{2}}(\partial\Omega)$:
 $\Phi_n u_0 = \frac{\partial u}{\partial \nu},$

$$\begin{cases} \Delta u + k^2 n(x)u = 0 & x \in \Omega, \\ u|_{\partial\Omega} = u_0. \end{cases}$$

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{array}{l} AQA + BQB = -B \\ AQB = BQA \end{array}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

- The following relations hold:

$$\mathcal{G}_1 - \mathcal{G}_n = \mathcal{G}_n(\Phi_n - \Phi_1)\mathcal{G}_1 \quad [\text{Nachman : 88}]$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \Phi_n - \Phi_1 \quad \& \quad \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}} = \Phi_n - \Phi_1$$

$$\Rightarrow \mathcal{G}_n^{-1} - \mathcal{G}_1^{-1} = \overline{\mathcal{G}_n^{-1}} - \overline{\mathcal{G}_1^{-1}}$$

- Put $A = \operatorname{Re} \mathcal{G}_n$, $B = \operatorname{Im} \mathcal{G}_n$, $Q = \frac{1}{2}(\mathcal{G}_1^{-1} - \overline{\mathcal{G}_1^{-1}})$

$$\Rightarrow \begin{cases} AQA + BQB = -B \\ AQB = BQA \end{cases}$$

- $Q = -T^*T$, where $T: L^2(\partial\Omega) \rightarrow L^2(S^{d-1})$ is a compact, injective, dense range operator (*Dirichlet-to-Farfield map*).

III. Algebraic relations – Derivation

It follows that with $\tilde{A} = TAT^*$, $\tilde{B} = TBT^*$ one has

$$\begin{aligned}\tilde{A}^2 &= \tilde{B} - \tilde{B}^2 \\ \tilde{A}\tilde{B} &= \tilde{B}\tilde{A}\end{aligned}$$

Corollaries:

- If $\tilde{A} > 0$, then

$$\begin{aligned}A &= T^{-1}(\tilde{B} - \tilde{B}^2)^{1/2}(T^*)^{-1} \\ \tilde{B} &= TBT^*\end{aligned}$$

- Spectrum of $\tilde{F} = \tilde{A} - i\tilde{B}$ lies on $|z + \frac{i}{2}| = \frac{1}{2}$ (it is a *farfield operator*).

III. Algebraic relations – Derivation

It follows that with $\tilde{A} = TAT^*$, $\tilde{B} = TBT^*$ one has

$$\begin{aligned}\tilde{A}^2 &= \tilde{B} - \tilde{B}^2 \\ \tilde{A}\tilde{B} &= \tilde{B}\tilde{A}\end{aligned}$$

Corollaries:

- If $\tilde{A} > 0$, then

$$\begin{aligned}A &= T^{-1}(\tilde{B} - \tilde{B}^2)^{1/2}(T^*)^{-1} \\ \tilde{B} &= TBT^*\end{aligned}$$

- Spectrum of $\tilde{F} = \tilde{A} - i\tilde{B}$ lies on $|z + \frac{i}{2}| = \frac{1}{2}$ (it is a *farfield operator*).

III. Algebraic relations – Derivation

It follows that with $\tilde{A} = TAT^*$, $\tilde{B} = TBT^*$ one has

$$\begin{aligned}\tilde{A}^2 &= \tilde{B} - \tilde{B}^2 \\ \tilde{A}\tilde{B} &= \tilde{B}\tilde{A}\end{aligned}$$


Corollaries:

- If $\tilde{A} > 0$, then

$$\begin{aligned}A &= T^{-1}(\tilde{B} - \tilde{B}^2)^{1/2}(T^*)^{-1} \\ \tilde{B} &= TBT^*\end{aligned}$$

- Spectrum of $\tilde{F} = \tilde{A} - i\tilde{B}$ lies on $|z + \frac{i}{2}| = \frac{1}{2}$ (it is a *farfield operator*).


IV. Uniqueness theorems



Classical inverse problem. G_n at fixed k determines $n - 1 \in L^\infty(\Omega, \mathbb{R})$ uniquely if k^2 is not a Dirichlet eigenvalue

- R.G. Novikov (*Funct. Anal. Appl.* 22, 1988) and Y.M. Berezanskii (*Tr. Mosk. Mat. Obschch.* 7, 1958)
- A.I. Nachman (*Ann. of Math.*, 1988)


IV. Uniqueness theorems



Classical inverse problem. \mathcal{G}_n at fixed k determines $n - 1 \in L^\infty(\Omega, \mathbb{R})$ uniquely if k^2 is not a Dirichlet eigenvalue

- R.G. Novikov (*Funct. Anal. Appl.* 22, 1988) and Y.M. Berezanskii (*Tr. Mosk. Mat. Obschch.* 7, 1958)
- A.I. Nachman (*Ann. of Math.*, 1988)


IV. Uniqueness theorems



Classical inverse problem. \mathcal{G}_n at fixed k determines $n - 1 \in L^\infty(\Omega, \mathbb{R})$ uniquely if k^2 is not a Dirichlet eigenvalue

- R.G. Novikov (*Funct. Anal. Appl.* 22, 1988) and Y.M. Berezanskii (*Tr. Mosk. Mat. Obschch.* 7, 1958)
- A.I. Nachman (*Ann. of Math.*, 1988)

IV. Uniqueness theorems



Classical inverse problem. \mathcal{G}_n at fixed k determines $n - 1 \in L^\infty(\Omega, \mathbb{R})$ uniquely if k^2 is not a Dirichlet eigenvalue

- R.G. Novikov (*Funct. Anal. Appl.* 22, 1988) and Y.M. Berezanskii (*Tr. Mosk. Mat. Obschch.* 7, 1958)
- A.I. Nachman (*Ann. of Math.*, 1988)

IV. Uniqueness theorems – Helmholtz equation

Lemma. $\forall M > 0 \exists \varepsilon: \|n\|_{\infty} \leq M \text{ and } k \in (0, \varepsilon) \implies \tilde{A} > 0$

Theorem 1. *For any $M > 0$ there exists ε such that if $k \in (0, \varepsilon)$, $\|n_{1,2}\|_{\infty} \leq M$ and $\operatorname{Im} \mathcal{G}_1(k) = \operatorname{Im} \mathcal{G}_2(k)$ then $n_1 = n_2$.*


IV. Uniqueness theorems – Helmholtz equation

Lemma. $\forall M > 0 \exists \varepsilon: \|n\|_\infty \leq M \text{ and } k \in (0, \varepsilon) \implies \tilde{A} > 0$

Theorem 1. *For any $M > 0$ there exists ε such that if $k \in (0, \varepsilon)$, $\|n_{1,2}\|_\infty \leq M$ and $\text{Im } \mathcal{G}_1(k) = \text{Im } \mathcal{G}_2(k)$ then $n_1 = n_2$.*

IV. Uniqueness theorems – Schrödinger equation

$$(-\Delta_{x_1} + v(x_1) - k^2) G_v^+(x_1, x_2) = \delta(x_2)$$




(*) Let $\tilde{F} = \tilde{A} - i\tilde{B}$. If $\tilde{F}f_1 = \lambda_1 f_1$, $\tilde{F}f_2 = \lambda_2 f_2$ are such that $\text{Im } \lambda_1 = \text{Im } \lambda_2$ then $\text{Re } \lambda_1 = \text{Re } \lambda_2$... is generic, at least, for small k , v and starlike Ω

Theorem 2. Suppose that $\text{Re } \mathcal{G}_{v_0}(k)$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*). Then $\exists \delta$ such that if $\|v_{1,2} - v_0\|_\infty < \delta$ and $\text{Im } \mathcal{G}_{v_1}(k) = \text{Im } \mathcal{G}_{v_2}(k)$, then $v_1 = v_2$.

IV. Uniqueness theorems – Schrödinger equation

$$(-\Delta_{x_1} + v(x_1) - k^2) G_v^+(x_1, x_2) = \delta(x_2)$$




(*) Let $\tilde{F} = \tilde{A} - i\tilde{B}$. If $\tilde{F}f_1 = \lambda_1 f_1$,
 $\tilde{F}f_2 = \lambda_2 f_2$ are such that
 $\text{Im } \lambda_1 = \text{Im } \lambda_2$ **then** $\text{Re } \lambda_1 = \text{Re } \lambda_2$
... is generic, at least, for small k , v
and starlike Ω

Theorem 2. Suppose that $\text{Re } \mathcal{G}_{v_0}(k)$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*).
Then $\exists \delta$ such that if $\|v_{1,2} - v_0\|_\infty < \delta$ and $\text{Im } \mathcal{G}_{v_1}(k) = \text{Im } \mathcal{G}_{v_2}(k)$,
then $v_1 = v_2$.

IV. Uniqueness theorems – Schrödinger equation

$$(-\Delta_{x_1} + v(x_1) - k^2) G_v^+(x_1, x_2) = \delta(x_2)$$




(*) Let $\tilde{F} = \tilde{A} - i\tilde{B}$. If $\tilde{F}f_1 = \lambda_1 f_1$, $\tilde{F}f_2 = \lambda_2 f_2$ are such that $\text{Im } \lambda_1 = \text{Im } \lambda_2$ then $\text{Re } \lambda_1 = \text{Re } \lambda_2$... is generic, at least, for small k, v and starlike Ω

Theorem 2. Suppose that $\text{Re } \mathcal{G}_{v_0}(k)$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*). Then $\exists \delta$ such that if $\|v_{1,2} - v_0\|_\infty < \delta$ and $\text{Im } \mathcal{G}_{v_1}(k) = \text{Im } \mathcal{G}_{v_2}(k)$, then $v_1 = v_2$.

IV. Uniqueness theorems – Schrödinger equation

$$(-\Delta_{x_1} + v(x_1) - k^2) G_v^+(x_1, x_2) = \delta(x_2)$$



(*) Let $\tilde{F} = \tilde{A} - i\tilde{B}$. If $\tilde{F}f_1 = \lambda_1 f_1$, $\tilde{F}f_2 = \lambda_2 f_2$ are such that $\text{Im } \lambda_1 = \text{Im } \lambda_2$ then $\text{Re } \lambda_1 = \text{Re } \lambda_2$... is generic, at least, for small k, v and starlike Ω

Theorem 2. Suppose that $\text{Re } \mathcal{G}_{v_0}(k)$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*). Then $\exists \delta$ such that if $\|v_{1,2} - v_0\|_\infty < \delta$ and $\text{Im } \mathcal{G}_{v_1}(k) = \text{Im } \mathcal{G}_{v_2}(k)$, then $v_1 = v_2$.

IV. Uniqueness theorems – Acoustic equation

- Acoustic equation:

$$-\nabla \cdot \left(\frac{1}{\rho_j} \nabla G_j^+(x, y) \right) - \frac{\omega^2}{\rho_j c_j^2} G_j^+(x, y) = \delta_y(x),$$

+ Sommerfeld radiation condition

Theorem 3. *For any $\omega_{1,2}$, ρ_0 , c_0 such that $\mathcal{G}_0(\omega_{1,2})$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*) there exists δ such that if $\|\rho_{1,2} - \rho_0\| + \|c_{1,2} - c_0\| < \delta$ and $\text{Im } \mathcal{G}_1(\omega_{1,2}) = \text{Im } \mathcal{G}_2(\omega_{1,2})$ then $\rho_1 = \rho_2$, $c_1 = c_2$.*

IV. Uniqueness theorems – Acoustic equation

- Acoustic equation:

$$-\nabla \cdot \left(\frac{1}{\rho_j} \nabla G_j^+(x, y) \right) - \frac{\omega^2}{\rho_j c_j^2} G_j^+(x, y) = \delta_y(x),$$

+ Sommerfeld radiation condition

Theorem 3. *For any $\omega_{1,2}$, ρ_0 , c_0 such that $\mathcal{G}_0(\omega_{1,2})$ is injective in $H^{-\frac{1}{2}}(\partial\Omega)$, (*) there exists δ such that if $\|\rho_{1,2} - \rho_0\| + \|c_{1,2} - c_0\| < \delta$ and $\text{Im } \mathcal{G}_1(\omega_{1,2}) = \text{Im } \mathcal{G}_2(\omega_{1,2})$ then $\rho_1 = \rho_2$, $c_1 = c_2$.*

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\text{Re } \mathcal{G}_n$ and $\text{Im } \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\text{Re } \mathcal{G}_n$ and $\text{Im } \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

V. Conclusion

- Established results:
 - $\operatorname{Re} \mathcal{G}_n$ and $\operatorname{Im} \mathcal{G}_n$ are algebraically related if n is real-valued
 - Global uniqueness for PIP for Helm. equation at one small k
 - Local uniqueness for PIP for Schr. equation at one k
 - Local uniqueness for PIP for Acoust. equation at two k
- Limitations of the approach:
 - Absence of absorption
- What's next?
 - Other wave equations (more complete model of helioseismology)
 - Stability and reconstruction
 - Partial data

References

- [1] A. D. Agaltsov, T. Hohage, and R. G. Novikov. "Monochromatic Identities for the Green Function and Uniqueness Results for Passive Imaging". In: *SIAM Journal on Applied Mathematics* 78.5 (Oct. 2018), pp. 2865–2890. DOI: 10.1137/18m1182218.
- [2] Y. M. Berezanskii. "The uniqueness theorem in the inverse problem of spectral analysis for the Schrödinger equation". In: *Tr. Mosk. Mat. Obshch.* 7 (1958), pp. 3–62. DOI: 10.1090/trans2/035/07.
- [3] V. A. Burov, S. N. Sergeev, and A. S. Shurup. "The use of low-frequency noise in passive tomography of the ocean". In: *Acoustical Physics* 54.1 (Jan. 2008), pp. 42–51. ISSN: 1562-6865. DOI: 10.1134/S1063771008010077.
- [4] J. Garnier and G. Papanicolaou. "Passive sensor imaging using cross correlations of noisy signals in a scattering medium". In: *SIAM J. Imaging Sci.* 2.2 (2009), pp. 396–437. URL: <https://doi.org/10.1137/080723454>.
- [5] L. Gizon et al. "Computational helioseismology in the frequency domain: acoustic waves in axisymmetric solar models with flows". In: *Astronomy & Astrophysics* 600 (2017), A35. DOI: 10.1051/0004-6361/201629470.
- [6] A. I. Nachman. "Reconstructions from boundary measurements". In: *Ann. of Math. (2)* 128.3 (1988), pp. 531–576. DOI: 10.2307/1971435.
- [7] R. G. Novikov. "A multidimensional inverse spectral problem for the equation $-\Delta\psi + (v(x) - Eu(x))\psi = 0$ ". In: *Funktsional. Anal. i Prilozhen.* 22.4 (1988), pp. 11–22, 96. DOI: 10.1007/BF01077418.
- [8] P. Roux et al. "Ambient noise cross correlation in free space: Theoretical approach". In: *The Journal of the Acoustical Society of America* 117.1 (Jan. 2005), pp. 79–84. DOI: 10.1121/1.1830673.
- [9] R. Snieder, K. Wapenaar, and U. Wegler. "Unified Green's function retrieval by cross-correlation; connection with energy principles". In: *Physical Review E* 75.3 (Mar. 2007). DOI: 10.1103/physreve.75.036103.
- [10] R. Snieder et al. "A Comparison of Strategies for Seismic Interferometry". In: *Surveys in Geophysics* 30.4 (Oct. 2009), pp. 503–523. DOI: 10.1007/s10712-009-9069-z.
- [11] R. L. Weaver and O. I. Lobkis. "Ultrasonics without a Source: Thermal Fluctuation Correlations at MHz Frequencies". In: *Physical Review Letters* 87.13 (Sept. 2001), p. 134301. DOI: 10.1103/physrevlett.87.134301.