# The complex Grothendieck constant

New bounds, machine-checked in Lean

Alin Galatan, with several AI research systems

https://github.com/agalatan/grothendieck-lean

October 2026

$$1.401695322458 \;<\; K_G^{\mathbb{C}} \;<\; 1.404598161566$$

Both bounds are theorems of a Lean 4/Mathlib development, resting on explicitly stated analytic and computer-assisted inputs (Section 3).

With **no hypothesis at all**, every step proved in Lean: $K_G^{\mathbb{C}} \le 1.404909543$.

**Abstract**

Grothendieck's inequality bounds how much a semidefinite (vector) relaxation of a bilinear optimisation problem can exceed the true optimum. The best constant in the complex case, $K_G^{\mathbb{C}}$, has been known since 1987 to lie roughly in $[1.338, 1.405]$; its first two decimals were settled only in September 2026. We improve both bounds, to an interval of width 0.0029, and verify the arguments in Lean. This note explains the problem, the two constructions, exactly what the computer checks, the new inequalities and obstructions found along the way, and why the remaining gap is hard: the two sides are dual optimisation problems, and the obstruction on both is the nonlinear (cubic and higher) structure of bounded functions of Gaussian vectors.

## 1 The question

Let $A = (a_{ij})$ be an $m \times n$ complex matrix. Compare

$$C(A) = \sup_{|\varepsilon_i|=|\delta_j|=1} \Big|\sum_{i,j} a_{ij}\,\overline{\varepsilon_i}\,\delta_j\Big|, \qquad V(A) = \sup_{\|x_i\|=\|y_j\|=1} \Big|\sum_{i,j} a_{ij}\,\langle x_i, y_j\rangle\Big|,$$

the first over unimodular scalars, the second over unit vectors in any complex Hilbert space. Clearly $C(A) \le V(A)$. Grothendieck proved in 1953 that $V(A) \le K\,C(A)$ for a constant $K$ independent of $m$, $n$ and the dimension. The smallest such $K$ is the *complex Grothendieck constant* $K_G^{\mathbb{C}}$.

**Why it matters.**

- *Optimisation.* $V(A)$ is a semidefinite program; $C(A)$ is a hard combinatorial problem. $K_G^{\mathbb{C}}$ is the best approximation ratio of the semidefinite relaxation, and rounding algorithms for it are exactly upper-bound proofs.
- *Functional analysis.* $K_G$ is the norm of the identity between the injective and projective tensor norms on $\ell_\infty \otimes \ell_\infty$, and controls absolutely summing operators (Grothendieck's original setting).
- *Quantum information.* The real constant $K_G^{\mathbb{R}}$ is the largest ratio between quantum and classical values of two-player XOR games (Tsirelson); the finite-order constants $K_G(d)$ govern the nonlocality of qubit states.

**History.**

| year | who | lower bound | upper bound |
|---|---|---|---|
| 1953 | Grothendieck | | $K < \infty$ |
| 1984 | Davie | 1.33807 | |
| 1987 | Haagerup | | 1.40490913… (Gaussian phase rounding) |
| Sep 2026 | Guo, Fang, Lu | 1.35584631827 | |
| Sep 2026 | Heilman, Jones, Malavolta | 1.4 | 1.404898554746 |
| Oct 2026 | **this work** | **1.401695322458** | **1.404598161566** |
| Oct 2026 | this work, fully formal | | 1.404909543 (no hypothesis) |

Haagerup's integral $1/\gamma^\dagger = 1.404575934\ldots$, $\gamma^\dagger = \int_0^{\pi/2} \cos^2 t\,(1+\sin^2 t)^{-1/2} dt$, has been proposed as the exact value. It lies inside our interval, $2.22 \cdot 10^{-5}$ below the upper bound. For the real constant the corresponding recent results are $1.773 \le K_G^{\mathbb{R}} \le 1.7799$ (Heilman–Jones–Malavolta, October 2026).

## 2 Two dual constructions

Both bounds use the same Gaussian picture. Let $Z = (z_1, \ldots, z_n)$ be a standard complex Gaussian vector and decompose $L^2(\mathbb{C}^n, \gamma)$ into the *Wick chaoses* $H_{p,q}$ (polynomials of bidegree $(p,q)$ in $z, \bar{z}$, orthogonalised). Under the rotation $z \mapsto e^{i\theta} z$ every relevant object has *charge one*, so only the bidegrees $(1,0), (2,1), (3,2), \ldots$ (degrees $1, 3, 5, \ldots$) appear.

### 2.1 Upper bounds: rounding schemes

Given unit vectors, project a Gaussian onto them and round: $\varepsilon_i = f(\langle G, x_i\rangle)$ with $|f| \le 1$. The correlation of two rounded values depends only on $t = \langle x_i, y_j\rangle$,

$$\mathbb{E}\big[\overline{f(\cdot)}\, g(\cdot)\big] = H(t) = \sum_{k \ge 0} c_k\, t|t|^{2k}.$$

If $H(t) = c\,t + r(t)$ with $\|r\|_A = \sum_k |c_k - c\,\delta_{k0}| < c$, a tensor-power argument (Krivine's trick) gives $K_G^{\mathbb{C}} \le 1/(c - \|r\|_A)$. Haagerup's choice $f = \text{phase}$ gives 1.40491. Lean proves Haagerup's identity for $f = \text{phase}$ from scratch (Euler and Beta integrals, the law of orthonormal Gaussian coordinates); with a degree-29 preprocessor, 1024 exact terms and a signed resolvent tail certificate this gives the unconditional bound $K_G^{\mathbb{C}} \le 1.404909543$ (`haagerup_bound`), which needs no hypothesis. Our sharp rounding is a smoothed two-coordinate phase rounding, $F_0 = \text{phase}(z + \sqrt{1+a^2}\, v(|z|^2)\, w)$ with a twelve-term Laguerre profile $v$, multiplied by the cubic Taylor polynomial of a damped polynomial phase $\varphi$: $T = F_0 P_3(\varphi)$, $P_3(x) = 1 + ix - x^2/2 - ix^3/6$. The identity $|P_3(x)|^2 = 1 - x^4(3-x^2)/36$ and the proved bound $\varphi^2 \le 3$ give $|T| \le 1$, and a disk-valued rounding is as good as a unimodular one. Its kernel is $H(t) = \sum E[n,m]\, p(t)^n (-\tau(t))^m$ with an odd polynomial $p$ of degree 41 and $\tau$ of degree 15 with zero linear coefficient (any odd preprocessors of Wiener norm $\le 1$ are realised by tensor powers). Compared with $c\,t$, $c = 7119483103461/10^{13}$, the residual is at most $\delta = 458437859009/(5 \cdot 10^{17}) \approx 9.16876 \cdot 10^{-7}$:

$$K_G^{\mathbb{C}} \le \frac{1}{c-\delta} = \frac{5000000000000000}{355973696735190991} < 1.404598161566.$$

The whole difficulty is to keep *every* coefficient $c_k$, $k \ge 1$, small at once: improving the cubic coefficient $c_1$ tends to spoil $c_2, c_3, \ldots$, and the infinite tail has to be bounded too.

**Selected-spectrum comparison.** The finite head only sees $S = \{(n, m) : n + 3m \le 501\}$. Write its approximation error as $G_T - X = \sum_r D_r$, where $D_r$ is supported in degree band $B_r$ with $\|D_r\|_2 \le e_r$. Let $w_p \in [0, 1]$ bound the retained absolute coefficient sum of the preprocessed block $p$. For nonnegative radii $A_r^2 \ge \sum_{p \in S \cap B_r} w_p^2 E_X(p)$, Lean proves

$$\sum_{p \in S} w_p |E_{G_T}(p) - E_X(p)| \le 2 \sum_r A_r e_r + \Big(\sum_r e_r\Big)^2.$$

All pieces may share coordinates. Lean proves each coefficient cost by outward interval polynomial arithmetic; native arithmetic checks eighteen weighted radii from the stored table and its error allowance. The Gaussian component-error enclosures and their identification with the pieces remain explicit inputs.

## 2.2 Lower bounds: Hermite projection games

A lower bound needs an operator $T$ whose *vector* value is close to 1 while its *scalar* value is small. We use

$$T = P_{1,0} - \epsilon\, P_{2,1}, \qquad \epsilon = \tfrac{149}{200},$$

the projection onto linear functions minus a multiple of the projection onto the cubic chaos.

- *Vector value.* With $u_\ell = z_\ell / \|z\|$, $\sum_\ell \langle u_\ell, T u_\ell \rangle \ge 1 - (1 + \epsilon)/(n + 1) \to 1$.
- *Scalar value.* For all $|f|, |g| \le 1$, $|\langle f, Tg \rangle| \le B = 0.7134218$.

A finite-partition transfer turns this into matrices, so $K_G^{\mathbb{C}} \ge 1/B > 1.401695322458$.

The scalar cap is the hard part. Writing $h = \frac{1}{2}(f + g)$, $k = \frac{1}{2}(f - g)$,

$$\Re\langle f, Tg \rangle = t^2 - \epsilon \|P_{2,1} h\|^2 - \|P_{1,0} k\|^2 + \epsilon \|P_{2,1} k\|^2, \qquad t = \|P_{1,0} h\|.$$

Only the last term is positive: we must show that a *bounded* function cannot have too large a cubic part without paying elsewhere. After a rotation $P_{1,0} h = t\, z_1$, and the cubic part of $k$ splits into pieces that involve $z_1$ (*visible* sectors) and a *remote* residual $q \in H_{2,1}$ in the other coordinates. The proof then has three layers.

1. **Row reduction.** For each of 19 profiles $r$, the contribution of the visible sectors is the supremum of an explicit finite-dimensional Gaussian integral, a convex one-variable *potential* $\Psi_r$, evaluated at the scale of the remote residual. One gate variable is shared across rows.
2. **The law of the remote cubic.** What remains is a statement about the distribution of $U = |q|^2 / \mathbb{E}|q|^2$ for an arbitrary cubic Wick polynomial in arbitrarily many variables. We use exact identities and inequalities valid for *every* such cubic: the fourth-moment cap $\mathbb{E}U^2 \le 62$; the sixth-moment bound $\mathbb{E}U^3 \le 9m - 12 + 40.45313\,(m - 2)^{3/2}$, $m = \mathbb{E}U^2$, which rests on thirteen tensor-contraction facts including the Wick expansion of the sixth cumulant; a first-moment bound; and families of *Stein cuts* $\mathbb{E}\, G(U) \ge 0$, obtained by Gaussian integration by parts and completing squares against spectral budgets, a covariance order and *contraction-Gram budgets* (positive semidefinite Gram matrices of the coefficient tensor).
3. **A finite certificate.** The parameters are covered by 8,756 boxes in $(t, \tau, x)$; on each box a positive combination of rows and node certificates (351 mixed nodes, 9,237 row nodes, each a polynomial majorant checked in exact rational arithmetic) bounds the scalar value by $B$.

### 2.3 The two problems are one

For the real constant, Heilman, Jones and Malavolta proved an exact minimax: over rounding schemes $\mu$ with kernel coefficients $c_m(\mu)$ and Hermite projection games $Q = \sum_m q_m \Pi_m$,

$$\frac{1}{K_G} = \sup_{\mu} \Big[ c_1(\mu) - \sum_{m \geq 3} |c_m(\mu)| \Big] = \inf_{Q} \ \mathrm{OPT}(Q),$$

and the same duality holds in the complex case on charge-one bidegrees. An upper bound is a good point of the convex set of achievable kernels; a lower bound is a supporting hyperplane of that set. There is *no intrinsic gap*: the gap between our bounds is entirely the gap between restricted constructions and the cost of certifying them.

## 3 What exactly the computer checks

| **Proved in Lean** | **Assumed (certified outside Lean)** |
|---|---|
| *Lower bound.* `lower_bound_of_nodeCaps : NodeCaps → IsGrothendieckBound K → 1.401695322458 < K` | |
| the witness $T$ on $L^2$ of the Gaussian measure and its vector value; the transfer to matrices; the polarisation identity; the per-row Gaussian reduction of $\langle f, Tg \rangle$ for every unimodular pair (frames, fibre decomposition, Parseval, gates); all moment, Stein, spectral and contraction-Gram inequalities for every circular cubic, including the sixth-cumulant Wick identity; the finite certificate (13 single-threaded native checks) | `NodeCaps`: the 19 potentials $\Psi_r$ (defined in Lean) lie below their stored rational caps at the tabulated nodes. The separate `GrotheCaps` target reduces this to `CapsRest`: the vertex caps of rows 12–15 ($t = 0.84$–$0.855$) and any excluded-node caps. The checks for fifteen rows have a verification stamp; four rows retain external vertex caps. |
| *Upper bound.* `UpperChaos.fresh_upper_rounder`, for an $L^2$ charge-one truncation $X$ under component-error, table-accuracy and Hardy inputs, proves $K_G^{\mathbb{C}} \leq 500000000000000000/355973696735190991$. | |
| the concrete disk-valued rounding; Gaussian Hermite completeness, Parseval and Mehler, charge selection, summability; the transfer inequality (tensor powers); kernel composition and locality; the head through degree 501 from 21,168 stored energies by a verified interval evaluator (one `native_decide`); the tail bound; the endpoint; the selected-block comparison and native band masses | `ComparisonInputs`: an exact error decomposition, spectral supports and inherited Gaussian component norm caps. Lean derives the selected comparison $13311891/(15625 \cdot 10^9)$ (92.92% of the residual allowance). The input identification remains external. `hardy`: weighted square sum $7/4$ at radius $4149/4000$, with tail $36748/10^{12}$ (4.01%); finite table accuracy $6847009/(5 \cdot 10^{17})$. The native head is $28153/10^{12}$ (3.07%). |
| *Unconditional upper bound.* `haagerup_bound : IsGrothendieckBound (1404909543/1000000000)` | |
| Haagerup's identity for the phase rounding, $\mathbb{E}[\overline{\mathrm{phase}\langle g, x \rangle}\ \mathrm{phase}\langle g, y \rangle] = \sum_k \frac{\pi}{4} \frac{(1/2)_k^2}{k!(k+1)!} z|z|^{2k}$ (Euler integral, Gaussian integrals, Beta integrals, reduction to two complex Gaussians); the transfer through a degree-29 preprocessor $p(t) = tP(t^2)$ with $\|P\|_1 = 1$ and $\|P^4\|_1 \leq .998903$; 1024 exact terms, a finite polynomial resolvent and the full signed tail, checked by one `native_decide` | nothing: this bound has no hypothesis. The sharper conditional bound above uses a different rounding and retains its explicitly listed Gaussian inputs. |

Axioms: `propext`, `Classical.choice`, `Quot.sound`.

The native facts pinned in `scripts/check_axioms.py` check the upper head, selected-band masses, phase cover, Haagerup bound, cap chords and 13 lower-certificate chunks (18 facts in the default target; the report `scratch/Axioms.lean` lists 424 theorems). No `sorry`; CI builds the default

target and checks its axiom report. The 60 additional cap-range checks belong to the separate heavy `GrotheCaps` target, whose source stamp CI checks once recorded. The generic response and quintic-transfer lemmas add no native facts or endpoint assumptions. The lower bound still needs numerical caps; the sharp upper bound retains external Gaussian component-error and Hardy inputs. The weaker upper bound 1.404909543 is fully formal.

# 4 Discoveries along the way

Status labels: **[Lean]** machine-checked in this repository; **[P]** written proof; **[P,C]** proof with an exact or interval-arithmetic certificate replayed independently; **[F]** floating-point evidence only.

## 4.1 Signed cancellation gives a sharper fully formal bound

**[Lean]** Write $A = XP^2$. A finite polynomial $R$ with $\|1 - R(1 - A)\|_1 \le \eta < 1$ bounds every geometric prefix by $2\|RT\|_1/(1 - \eta)$. Abel summation then bounds the full decreasing-weight series while retaining signed cancellation. Coefficient locality makes this an infinite-tail theorem with only finite checks.

For an exact degree-29 preprocessor, the complete 1024-term normalized residual is at most $95 \cdot 10^{-9}$ and the whole infinite tail at most $174 \cdot 10^{-9}$. The approximate inverse has degree 64 and defect at most $10^{-12}$; its short product with $P^{17}$ is checked as one polynomial. The Gaussian phase identity and tensor transfer prove $K_G^{\mathbb{C}} \le 1.404909543$ with no external inputs. The exact data generate the Lean certificate reproducibly; an independent directed replay audits it.

The simpler obstruction is also exact: if $|P(-1)| = \|P\|_1 = 1$, then $\|P^n\|_1 = 1$ for every $n$. Breaking that common boundary phase permits power contraction; the resolvent additionally retains cancellation between successive composed powers.

## 4.2 Common responses, common tails, and an architecture separator

**[P; supporting algebra in Lean]** A symmetry-closed family of actual cubic responses has an exact five-mode first/cubic image and a fixed-size original-noise replica representation. It preserves the shared quadratic drivers: a scalar Gaussian used to describe a cubic's law must retain third-chaos noise correlation $\rho^3$, rather than acquire first-chaos correlation $\rho$.

**[Lean]** All correlation samples of a response pair use the same signed Hermite coefficients. Prices $a_i$ with $|\sum_i a_i \rho_i^{2k+1}| \le 1$ for all $k \ge 2$ bound the complete omitted absolute tail. A finite front and one geometric remainder budget certify every index. The resulting separator is linear in response columns, so it bounds whole convex pools, with explicit integration-error costs. Numerical pool screens still require certified input integrals; a favorable relaxation value is not an achievable rounding score.

**[Lean]** An exchange can retain the incumbent's signed coefficients with one error allowance. If $\sum_k |c_k - \widehat{c}_k| \le \xi$, then every admissible spectral price has error at most $\xi$, and the full reciprocal score satisfies $D(c) \ge D(\widehat{c}) - \xi$. The same coefficient error must not be charged independently at every sampled correlation. The current bounded phase-perturbation searches use this interface for discovery; their Gaussian integrals still require certification before they can improve either endpoint.

**[P,C; polynomial certificate in Lean]** The entire Gaussian-phase cubic-preprocessor family $H_{\mathrm{phys}}(A\rho + B\rho^3)$, $|A| + |B| \le 1$, including arbitrary complex phases and convex mixtures, has full reciprocal score at most 0.71186. This is below the incumbent $0.7119473934\ldots$. The separator is $\Re(c_0 - \frac{9}{25}c_1 - c_2 - c_3)$; positive polynomial certificates handle all complex phases. Thus optimizing this class further cannot improve the current upper bound. General non-Gaussian cubic responses

and nonlinear shared image fields are not excluded. The Gaussian-family coefficient extraction is a written application of the Gaussian phase identity; it is not yet connected end to end in Lean.

### 4.3 Inequalities for cubic Gaussian chaos

Throughout, $q$ is a circular cubic Wick polynomial (bidegree $(2,1)$) in any number of complex Gaussian variables, normalised by $\mathbb{E}|q|^2 = 1$, and $U = |q|^2$.

- **Sharp fourth moment [Lean].** Every real third-chaos variable satisfies $\mathbb{E}F^4 \le 93\,(\mathbb{E}F^2)^2$, with equality for $x^3 - 3x$. Hence $\mathbb{E}U^2 \le 62$.
- **Sixth moment [Lean].** $\mathbb{E}U^3 \le 9m - 12 + C_6(m-2)^{3/2}$ with $m = \mathbb{E}U^2$ and $C_6 = 40.45313$. The Wick monomial mixed with Gaussian noise attains 40.295, so the constant is sharp to 0.4%. The inequality is *false* without circularity: two real cubics $He_3(x_0)/\sqrt{12}$, $He_3(x_1)/\sqrt{12}$ give $m = 47$ and $\mathbb{E}U^3 = 13506 > 12622.55$.
- **The sixth cumulant as a sum over graphs [Lean].** The Wick identity for $\kappa_6$ of a cubic chaos is proved by expanding $\mathbb{E}F^2, \mathbb{E}F^4, \mathbb{E}F^6$ over closed cubic multigraphs, one Gaussian integration by parts per edge ($69 + 15 + 3$ certified steps). The disconnected graphs cancel, and the six connected cubic multigraphs on six vertices give the tensor contractions.
- **Contraction-Gram budgets [Lean].** A new family of universal inequalities between the chaos components of $U$ and of $q^2$. With $X = \Gamma(q,\bar{q})/3 - 1$, $c = \Gamma(q,q)/3$, $r = \Re(c\bar{q}^2)/U$, $P = \mathbb{E}[XU]$, $Q = \mathbb{E}[rU]$, for example

  $$\mathbb{E}X^2 + \tfrac{1}{10}\mathbb{E}r^2 \le \tfrac{34}{45}P - \tfrac{1}{18}Q.$$

  Each budget is certified by two positive semidefinite $2 \times 2$ Gram matrices of contractions of the coefficient tensor.
- **First moment [P,C].** $\mathbb{E}|q| \le 0.97073$, from a two-plane small-ball estimate (a winding argument on $S^3$ and a horizontal Lipschitz bound). The sharp value $\sqrt{\pi}/2 = 0.8862$, which the Gaussian attains, is open.

**A common budget for the same tensor.** The sixth moment and the Stein fluctuations cannot be optimized independently: both depend on the same contraction invariants. We now formalize this connection in Lean, including the real/complex normalization and a finite checker that reserves part of the contraction budget before accepting a Stein cut. Signed reserve coefficients are retained. The endpoint uses 41 such nodes, with each node restricted to its proved kurtosis range; the full checker and analytic bridge are in Lean. Retaining the upper bound on the negative cumulant contraction yields a stronger parameterized moment family. An exact transport identity then upgrades a node's majorant when its budget permits a larger moment-cut weight, without a new half-line search. These two refinements are also formalized in Lean. The moment costs also admit a convex formulation with two $2 \times 2$ PSD blocks and linear constraints. An exact rational checker connects each accepted recipe to the actual cubic moment inequality, replacing the outer parameter grid. Its exact scalar boundary is also proved in Lean. For moment weight $w > 0$ and $s_0$ cost $A > 0$, the minimum $h_0$ cost is

$$B_{\min}(w,A) = \begin{cases} 11781w/32 + 18225w^2/(4A), & 11A \le 360w, \\ 198w + 22500w^2/(A + 40w), & 11A \ge 360w. \end{cases}$$

Both branches have explicit feasible recipes. A three-rational checker for $(w, A, B)$ therefore replaces the auxiliary moment-cost variables, and Lean connects its acceptance to the actual cubic on the checked node range. The contraction reserve and pointwise majorant still need their own exact checks. This is optimality within the certificate family; it does not identify all attainable response laws.

### 4.4 The remote functional

Let $L = z_1(a + b|z_1|^2)$ with the constants of the witness, and for a cubic $G$ put $\Lambda(G) = \mathbb{E}|L+G| + \mathbb{E}|L-G| - \mathbb{E}|G|^2/\epsilon$ and $S = \sup_v\{2\mathbb{E}|L + \sqrt{v}U| - v/\epsilon\}$, $U$ a complex Gaussian. The conjecture $\Lambda(G) \le S$ for all cubics ("Conjecture K") is the analytic heart of the lower bound.

- **A one-function reformulation [P].** Circular symmetry, $\mathbb{E}|W| = (\pi/2)\mathbb{E}|\Re W|$, and the identity $|a+b| + |a-b| = 2\max(|a|,|b|)$ for real $a, b$ give exactly

$$\Lambda(G) = 2\mathbb{E}|L| + \pi\,\mathbb{E}\big(|\Re G| - |\Re L|\big)_+ - \mathbb{E}|G|^2/\epsilon,$$

  so $\sup_G \Lambda - 2\mathbb{E}|L| = \sup_t\big[\frac{\pi^2\epsilon}{4}\|P_{2,1}t\|^2 - \pi\mathbb{E}(|t|\,|\Re L|)\big]$ over real functions $|t| \le 1$. All of the gain sits near the hyperplane $\Re z_1 = 0$.

- **The Gaussian is a strict local maximum [P,C].** At the Gaussian point every direction that involves $z_1$ has negative second variation, certified in interval arithmetic for seven sector coefficients. The Hessian's kernel is the entire $z_1$-independent cubic sector; a negative fourth-order term controls those genuine cubic directions.

- **The independent sector [P,C].** For every cubic $q$ not involving $z_1$, $\Lambda(q) < S + 3.6\cdot 10^{-7}$. Moment inequalities alone stall about $10^{-5}$ above $S$, because $h(x) = 2\mathbb{E}|L + \sqrt{x}|$ is a Bernstein function and the three-moment relaxation is maximised by a two-point law [P]. Going below that floor needs the Stein and spectral cuts.

- **Two natural shortcuts are false [P,C].** $|q|^2$ does not dominate $\mathbb{E}|q|^2 \cdot \mathrm{Exp}(1)$ in convex order, and there is no Gaussian Laplace domination: $Q = X(4(|X|^2-2) + 3i(|Y|^2-1))/\sqrt{41}$ has $\mathbb{E}e^{-20|Q|^2} < 0.0453 < 1/21$.

### 4.5 Actual responses

Instead of bounding $\Lambda$ for all cubics, one can use that the cubic comes from a bounded response. Write $F$ for the cubic part of the response and $\mathcal{N}$ for the value of the witness on it.

- **Exact support [P].** The paired support $\frac{1}{2}(\mathbb{E}|A+B| + \mathbb{E}|A-B|)$ is exactly the pointwise support function of the complex response set $\{(h,k) = (e^{i\alpha}\cos\theta,\, ie^{i\alpha}\sin\theta)\}$, because $(|A+B| + |A-B|)^2 = 4\max_\theta |A\cos\theta - iB\sin\theta|^2$.

- **Concentrated cubics are harmless [P,C].** At $\epsilon = 1891/2500$ on the central rows $0.84 \le t \le 0.851$, $\mathcal{N} < 0.71325 < 1/1.402$ whenever the cubic part of the response concentrates enough energy on a structured subspace:
  - 80% on one diagonal Wick direction;
  - 90% on one mixed monomial $X^2\bar{Y}$, or on a whole slice $\{X^2\bar{A}\}$;
  - 93.5%, 96.2%, 97.6%, 99.2% on any cubic in $2, 3, 4, 6$ remote coordinates;
  - 95.5% to 99.95% on $H_{2,0}(A)H_{0,1}(B)$ with $\dim A \le 24$, via a sharp support theorem whose extremiser has law $R_m G/\sqrt{m(m+1)}$, $R_m \sim \Gamma(m,1)$.

  The remainder of the cubic may be arbitrary and fully coupled.

- **The obstruction to covering [P].** $Q_N = X_1(\sum_{j\le N}|X_j|^2 - N - 1)/\sqrt{N+1}$ keeps an influential direction while its projection onto every fixed-dimensional core tends to zero. So "small core or Gaussian" is not an exhaustive dichotomy.

- **How far from the benchmark structured cubics sit [F].** Remote-only payoffs relative to $S$:

| sum of $N$ slices, $N=$ | 1 | 2 | 4 | 8 | 16 | 24 | $\infty$ (Gaussian) |
|---|---|---|---|---|---|---|---|
| $S - \sup\Lambda$ $(10^{-3})$ | 1.68 | 1.42 | 1.08 | 0.73 | 0.43 | 0.32 | 0 |

The tested concentrated cubics leave room of order $10^{-3}$; these diffuse sums approach the benchmark. This evidence does not classify all near-maximizers.

### 4.6 Limits of the witness method

- **Ceilings [F] and [P,C].** The norm of the cubic witness $P_{1,0} - \epsilon P_{2,1}$ is about 0.71314 at its best $\epsilon \approx 0.756$ [F]: no certificate for it exceeds 1.40226. Explicit "chirp" responses prove that games of degree $\leq 5$ cannot exceed 1.4036705, and of degree $\leq 7$ cannot exceed 1.4044895 [P,C].
- **Extending the pattern fails [P,C].** The full alternating game $\sum_{k\geq 0}(-1)^k P_{k+1,k}$ has $\sup_n \|\cdot\|_{\infty\to 1} = 1$, attained by explicit unimodular responses (one in complex dimension 308,835 is replayed). So better high-degree games must be re-optimised against new responses, not extrapolated.
- **No duality gap, but no free lunch [P].** By the minimax of Section 2.3, combining a pool of actual responses truncated at level $D$, value $v_D$, with a witness certified to $v_D + \delta$ and a mixture whose omitted coefficient tail is at most $\tau$, gives $1/(v_D + \delta) \leq K_G^{\mathbb{C}} \leq 1/(v_D - \tau)$. The interval for $1/K_G^{\mathbb{C}}$ has width exactly $\delta + \tau$.
- **A global spherical form, and why it does not certify cheaply [P] and [Lean].** Finite Gaussian Hermite games transfer, in every Gaussian dimension, to compact operators on the sphere with the same bounded responses and the same shared drivers; the numerical bounds are unchanged. The dual operator can then be truncated with a rigorous error: the cubic multiplier tail and the signed cap assembly are proved in Lean (`SphericalTransfer.cubic_tail_32_12`, `global_cap_from_truncation`); the analytic bridge is a written proof (`docs/SPHERICAL_TRANSFER.md`). But the operator has no square-integrable kernel: its multiplicity-weighted squared coefficients grow like $r^7$. So no Hilbert–Schmidt-type certificate can control it, and direct coefficient optimisation is impractical (about $8 \cdot 10^{20}$ coefficients in the tested truncation). A useful norm cap must bound the signed action on bounded responses, exploiting the unitary symmetry.

## 5 Why the gap is hard

**The witness class caps the lower bound.** A game supported on degrees $\{1, 3\}$ sees only the cubic constraint on roundings. Even a perfect evaluation of the norm of $P_{1,0} - \epsilon P_{2,1}$ cannot certify more than about 1.40226 (numerical, best $\epsilon \approx 0.756$). Proved obstructions are known for higher degrees: every game of degree $\leq 5$ is capped at 1.4036705, and every game of degree $\leq 7$ at 1.4044895, both below Haagerup's value. Reaching the upper end of the interval therefore needs games of degree $\geq 9$; in the real case the certified game uses all odd degrees up to 21.

**Certifying higher degree is the real bottleneck.** Two different certificate architectures, gated rows and support functions, both lose about $2 \cdot 10^{-2}$ on the best degree-5 game. The reason is structural: a bounded function with a large cubic part is forced to have a degree-five part too. Through the visible coordinate, the cubic part sees the average of the remote coefficient profile and the quintic part its variation, and boundedness makes the profile vary. A certificate that treats the chaos degrees as independent cannot see this coupling. The coupling is *driver sharing*: in the dimension-limit model the best degree-5 game has norm 0.712464 ($1/N = 1.40358$) when its

degree-5 mixed sectors $(|z_1|^2-1)W$ and $z_1^2\overline{W}$ are driven by the same remote cubic $W$ as the cubic sector, and 0.724753 when they are driven by independent copies; the optimum then switches the degree-5 part off. So a higher-degree certificate must let one remote residual feed several degrees through the visible coordinate. The cubic rows already do this for their degree-3 sectors; extending them to the degree-5 companions of the same residual is the next construction [F].

**Why the complex case is harder than the real one.** For real $\pm 1$ responses, $h=\frac{1}{2}(f+g)$ and $k=\frac{1}{2}(f-g)$ have disjoint supports, so the two sides of a game decouple at every degree. For complex unimodular responses $|h|^2+|k|^2=1$ and $\Re(h\bar{k})=0$ force $f=e^{i\alpha}e^{i\theta}$, $g=e^{i\alpha}e^{-i\theta}$: one shared phase field $\alpha$ drives both sides, and they never separate. The same shared phase is why marginal laws do not determine the relevant nonlinear correlations: $(H(X),X)$ and $(H(Y),X)$, with $H(X)=X(|X|^2-2)/\sqrt{2}$ and $X,Y$ independent, have identical marginals and covariances, but $\mathbb{E}[\bar{r}\,q|q|^2]$ equals $8\sqrt{2}$ in the first case and 0 in the second.

## 6 Open problems

Planned work is tracked as GitHub issues of the repository: epics #12 (unconditional lower bound), #17 (upper inputs), #22 (better lower bound), #29 (build infrastructure) and #34 (research plans).

1. Complete and verify the node caps (`CapsRest` in the separate target), which would make the lower bound fully machine-checked. The caps of rows 12–15 are localised caps: coverage by constrained pieces is checked in Lean, and most pieces pass; the remaining pieces fail the exact positivity check of their S-lemma matrix by about $10^{-5}$ and need re-optimised multipliers.
2. Discharge the sharp upper rounding's numerical comparison, energy-table error and contour tail bounds; its Hermite expansion and structural analytic bridge are formalized.
3. Certify a game of degree $\geq 5$ with a *multi-grade fibre*: one bounded function of the visible coordinate and one remote direction per degree, so that the cross-degree coupling is kept. This is the route to a lower bound above 1.403.
4. Run a complex primal/dual search, alternating linear programs over games of degree $\leq D$ and pools of roundings, to locate $K_G^{\mathbb{C}}$ to about $10^{-4}$ and to test whether Haagerup's value $1/\gamma^{\dagger}$ is the answer.
5. Prove the sharp first-moment bound $\mathbb{E}|q| \leq \frac{\sqrt{\pi}}{2}\|q\|_2$ for cubic Wick polynomials. It is natural, it appears to be open, and it is the largest single lever on the law side.

## How this was made

The constructions and certificates were found by several AI research systems working on a shared manuscript, cross-checking each other's packages by independent replay. The endpoint theorems have the explicit Lean hypothesis lists in Section 3, and their axioms are audited in CI. The discoveries labelled [P] and [C] also include paper proofs and external exact or interval certificates; they are not all formalized in Lean. Selected written proofs and reproducible certificate records accompany the repository; the larger working manuscript is maintained separately.

## References


- A. Grothendieck, *Résumé de la théorie métrique des produits tensoriels topologiques*, 1953.

- U. Haagerup, *A new upper bound for the complex Grothendieck constant*, Israel J. Math. 60 (1987).
- A. M. Davie, *Lower bound for $K_G$*, unpublished note, 1984.
- S. Heilman, C. Jones, G. Malavolta, *Sharper bounds for the complex Grothendieck constant*, arXiv:2609.25462.
- S. Heilman, C. Jones, G. Malavolta, *Primal/dual method for the Grothendieck constant*, arXiv:2610.10477.
- Guo, Fang, Lu, arXiv:2609.07000.
- R. Saha et al., *New lower and upper bounds for the Grothendieck constant*, arXiv:2608.11158.
- A. Naor, O. Regev, *Krivine schemes are optimal*, Proc. AMS 142 (2014), arXiv:1205.6415.
- G. Pisier, *Grothendieck's theorem, past and present*, Bull. AMS 49 (2012), arXiv:1101.4195.
- N. Brunner et al., *Bell nonlocality*, Rev. Mod. Phys. 86 (2014), arXiv:1303.2849.