

MatGeo Assignment Presentation

Agamjot Singh,
EE24BTECH11002,
IIT Hyderabad.

November 6, 2024

Outline

Table of Contents

Problem

Solution

General Equation of Circle and Variable Description

Equation of given circle as a general equation in matrix form

Expressing boundary of D in parametric form

Using Green's theorem to find area using line integral

Calculating the area using line integral

Graph

C Code

Python Code

Problem

Problem Statement

Sketch the region $(x, 0) : y = \sqrt{4 - x^2}$ and x -axis. Find the area of the region using integration.

Solution

General Equation of Circle and Variable Description

The general equation of circle is given by

$$\|\mathbf{x}\|^2 + 2\mathbf{u}^\top \mathbf{x} + f = 0 \quad (3.1)$$

Variable	Description
O	Center of the circle
r	Radius of the circle
u	$-\mathbf{O}$
f	$\ \mathbf{u}\ ^2 - r^2$
D	Region for which the area has to be found
A	Area of the region

Equation of given circle as a general equation in matrix form

The given curve $y = \sqrt{4 - x^2}$ is that of a semicircle, since $y \geq 0$.
The equation of the curve can be written as

$$x^2 + y^2 - 4 = 0, y \geq 0 \quad (3.2)$$

By comparing with equation (3.1), the parameters of the circle are

$$\mathbf{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, f = -4 \implies r = 2, \mathbf{O} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (3.3)$$

Expressing boundary of D in parametric form

Boundary of D is the semicircle of radius r , which we can parameterize (in counter clock-wise orientation) using

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} r \cos t \\ r \sin t \end{pmatrix}, 0 \leq t \leq \pi \quad (3.4)$$

Using Green's theorem to find area using line integral

Green's Theorem:

Let C be a curve in the plane, and D be the region bounded by it. If L and M are the functions of (x, y) defined on an open region containing D , then

$$\oint_C (L dx + M dy) = \int \int_D \left(\frac{\partial M}{\partial x} - \frac{\partial L}{\partial y} \right) dA \quad (3.5)$$

In this case, to find the area of the region D , we substitute $L = -\frac{y}{2}$ and $M = \frac{x}{2}$, then

$$\frac{1}{2} \oint_C (x dy - y dx) = \int \int_D dA = \text{area of region } D \quad (3.6)$$

Calculating the area using line integral

By Green's Theorem (3.6),

$$\text{area of } D = A = \iint dA \quad (3.7)$$

$$= \frac{1}{2} \int_C x \, dy - y \, dx \quad (3.8)$$

$$= \frac{1}{2} \int_0^\pi r^2 (\cos^2 t + \sin^2 t) \, dt \quad (3.9)$$

$$= \frac{r^2}{2} \int_0^\pi dt \quad (3.10)$$

$$= \frac{\pi r^2}{2} \quad (3.11)$$

$$= 2\pi \quad (3.12)$$

Graph




Figure 1: Shaded area representing area of region given

C Code

C function for getting ' n ' number of points to graph a circle of radius r and center (x, y) :

```
float **circleGet(int n, float x, float y, float r) {  
    float **pts = (float **) malloc(sizeof(float *) * n);  
    float theta = 0;  
    for(int i = 0; i < n; i++){  
        pts[i] = (float *) malloc(sizeof(float) * 2 * n);  
        pts[i][0] = x + r*cos(theta);  
        pts[i][1] = y + r*sin(theta);  
        theta += 2*PI/n;  
    }  
    return pts;  
}
```

The python code for generating the graph can be found at:

```
https://github.com/agamjotsingh1/EE1030/blob/main/  
matgeo\_questions/q9/codes/graph.py
```