



planetmath.org

Math for the people, by the people.

syntactic congruence

Canonical name	SyntacticCongruence
Date of creation	2013-03-22 18:52:08
Last modified on	2013-03-22 18:52:08
Owner	Ziosilvio (18733)
Last modified by	Ziosilvio (18733)
Numerical id	6
Author	Ziosilvio (18733)
Entry type	Definition
Classification	msc 68Q70
Classification	msc 20M35
Defines	syntactic semigroup
Defines	syntactic monoid
Defines	syntactic morphism
Defines	maximality property of syntactic congruence

Let S be a semigroup and let $X \subseteq S$. The relation

$$s_1 \equiv_X s_2 \text{ iff } \forall l, r \in S (ls_1r \in X \text{ iff } ls_2r \in X) \quad (1)$$

is called the *syntactic congruence* of X . The quotient S/\equiv_X is called the *syntactic semigroup* of X , and the natural morphism $\phi : S \rightarrow S/\equiv_X$ is called the *syntactic morphism* of X . If S is a monoid, then S/\equiv_X is also a monoid, called the *syntactic monoid* of X .

As an example, if $S = (\mathbb{N}, +)$ and $X = \{n \in \mathbb{N} \mid \exists k \in \mathbb{N} \mid n = 3k\}$, then $m \equiv_X n$ if $m \bmod 3 = n \bmod 3$, and the syntactic monoid is isomorphic to the cyclic group of order three.

It is straightforward that \equiv_X is an equivalence relation and X is union of classes of \equiv_X . To prove that it is a congruence, let $s_1, s_2, t_1, t_2 \in S$ satisfy $s_1 \equiv_X s_2$ and $t_1 \equiv_X t_2$. Let $l, r \in S$ be arbitrary. Then $ls_1t_1r \in X$ iff $ls_2t_1r \in X$ because $s_1 \equiv_X s_2$, and $ls_2t_1r \in X$ iff $ls_2t_2r \in X$ because $t_1 \equiv_X t_2$. Then $s_1t_1 \equiv_X s_2t_2$ since l and r are arbitrary.

The syntactic congruence is both left- and right-invariant, *i.e.*, if $s_1 \equiv_X s_2$, then $ts_1 \equiv_X ts_2$ and $s_1t \equiv_X s_2t$ for any t .

The syntactic congruence is maximal in the following sense:

- if χ is a congruence over S and X is union of classes of χ ,
- then $s\chi t$ implies $s \equiv_X t$.

In fact, let $l, r \in S$: since $s\chi t$ and χ is a congruence, $lsr\chi ltr$. However, X is union of classes of χ , therefore lsr and ltr are either both in X or both outside X . This is true for all $l, r \in S$, thus $s \equiv_X t$.